flusser@utia.cas.cz

Prof. Ing. Jan Flusser, DrSc.

Digitální zpracování obrazu

Lecture 1

Důležité informace

- Rozsah: ZS, 3+0, Zk
- http://zoi.utia.cz/teaching

Navazující předměty

- **NPGR013 (J. Flusser, B. Zitová)**
- **NPGR022 (J. Flusser, B. Zitová)**
- **NPGR029 (F. Šroubek, J. Flusser)**
- **NPGR032 (A. Novozámský, B. Zitová)**

Digitálních fotografií je všude plno ...

Smartphones Cause Photography Boom

Number of digital photos taken worldwide*



Devices used in 2017



@StatistaCharts

* estimates

Source: InfoTrends via Bitkom

statista

Jak vzniknou?



$$\begin{bmatrix} 16 & 11 & 10 & 16 & 24 & 40 & 51 & 61 \\ 12 & 12 & 14 & 19 & 26 & 58 & 60 & 55 \\ 14 & 13 & 16 & 24 & 40 & 57 & 69 & 56 \\ 14 & 17 & 22 & 29 & 51 & 87 & 80 & 62 \\ 18 & 22 & 37 & 56 & 68 & 109 & 103 & 77 \\ 24 & 35 & 55 & 64 & 81 & 104 & 113 & 92 \\ 49 & 64 & 78 & 87 & 103 & 121 & 120 & 101 \\ 72 & 92 & 95 & 98 & 112 & 100 & 103 & 99 \end{bmatrix}$$



Jak se tam dostanou barvy?







Fotografie nebývají ideální





Jak vylepšit nekvalitní foto?

Všichni známe Photoshop ...



... na jednoduché úpravy stačí

Kde Photoshop nestačí



(F1, F2, … , Fn)

2AC 7652

Jak odstranit třes ruky?

Segmentace



Originál



Detekce objektů


Groups
background
Roofs
Road
Grass
Trees
Trail
Water
Shadow

Historie

1921

1929



1964

Aplikační oblasti

- medicína





Aplikační oblasti

- Dálkový průzkum





Aplikační oblasti

- Bezpečnost, doprava



Aplikační oblasti

- Kulturní dědictví

Image Proc. x Comput. Graphics



Digital Image Processing

- Digitization (sampling + quantizing)
- Preprocessing (contrast and brightness changes, denoising, sharpening, …)
- Image analysis (object detection and recognition, scene understanding)
- Image coding (compression)

Mathematical background

- **Convolution**
- **Fourier Transform**

Convolution

- Definition in continuous domain
- Properties
- Delta function
- Discrete convolution, boundary effect

Convolution

$$* : L_1 \times L_1 \to L_1 \qquad (f * g)(x) = \int_{-\infty}^{\infty} f(t)g(x-t)dt$$



Basic properties

$$f * g = g * f$$

$$f * (g * h) = (f * g) * h$$

$$a(f * g) = (af) * g = f * (ag)$$

$$f * (g + h) = (f * g) + (f * h)$$

Dirac delta-function

$$\delta(x) = \begin{cases} \infty, & x = 0 \\ 0, & x \neq 0 \end{cases}$$

$$\int \delta(x)\,\mathrm{d}x = 1$$





Delta-function properties

$$\delta(ax) = \frac{\delta(x)}{|a|}$$

$$f * \delta = f$$

$$(f * \delta)(a) = \int_{-\infty}^{\infty} f(x)\delta(a - x)dx = f(a)$$

$$(f * \delta)(x - a) = f(x - a)$$

Cross-correlation

$$(f \star g)(x) = \int_{-\infty}^{\infty} f(t)g(t-x)dt$$

$$f \star g \neq g \star f$$

Convolution in 2D

- Definition $* : L_1 \times L_1 \rightarrow L_1$

$$(f * g)(x, y) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(s, t) g(x - s, y - t) ds dt$$

Discrete convolution

$$(f * g)(n) = \sum_{m=-\infty}^{\infty} f[m] \cdot g[n-m]$$

Boundary effect

Convolution mask



Type

“valid”

Type

“same”

Type

“full”

Boundary effect treatment

- Zero padding
- Mirror extension
- Periodic extension



Computing complexity

- From definition – O(N^2.M^2)
- For a separable kernel (rank = 1) – O(N^2.M)

$$g(x,y) = g_1(x) \cdot g_2(y)$$

$$(f*g)(x,y) = \int_{-\infty}^{\infty} g_1(x-s)\left(\int_{-\infty}^{\infty} f(s,t)g_2(y-t)dt\right)ds$$

- Hardware acceleration (DSP, graphic cards, ...)

Fourier transform

Recalling Fourier series

$\{\phi_k\}$ – ON basis of a Hilbert space $\mathcal{H}$, $\quad f \in \mathcal{H}$

$c_k = (f, \phi_k)$ – Fourier coefficients

$$f = \sum_{i=1}^{\infty} c_k \phi_k$$

Fourier series in $L_2(-\pi, \pi)$

$$\{\phi_k\} = \{1, \sin nx, \cos mx\}$$

$$f(x) = \frac{a_0}{2} + \sum_{k=1}^{\infty} (a_k \cos kx + b_k \sin kx)$$

$$a_k = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos kx \, \mathrm{d}x \qquad b_k = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin kx \, \mathrm{d}x$$

Fourier transform

$$\{\phi_u(x)\} = \{e^{-2\pi i(ux)}\}$$

$$F(u) = \int_{\mathrm{R}} f(x) e^{-2\pi i u x} \mathrm{d}x$$

$$f(x) = \int_{\mathrm{R}} F(u) e^{2\pi i u x} \mathrm{d}u$$

Properties of the FT

If $f \in L_1$ then FT exists

Generally $F \notin L_1$

FT is "one-to-one" mapping: $f \longleftrightarrow F$

$$F = \mathcal{R}(F) + i\mathcal{I}(F) = |F| \cdot e^{i \cdot ph(F)}$$

Properties of the FT

- **linearity**

$$\mathcal{F}[af(x) + bg(x)] = a\mathcal{F}[f(x)] + b\mathcal{F}[g(x)] = aF(k) + bG(k).$$

- **convolution** **convolution theorem**

$$\mathcal{F}[f]\mathcal{F}[g] \quad = \quad \mathcal{F}[f * g]$$

- **shift** **shift theorem**

$$\mathcal{F}_x[f(x - x_0)](k) = e^{-2\pi i k x_0} F(k).$$

- **rotation**

$$\mathcal{F}(R(f)) = R(\mathcal{F}(f))$$

- **scaling** **similarity theorem**

$$\mathcal{F}_x[f(ax)](k) = |a|^{-1} F\left(\frac{k}{a}\right).$$

scaling
similarity theorem

$$\mathcal{F}_x[f(ax)](k) = |a|^{-1} F\left(\frac{k}{a}\right).$$



Rectangular pulse





Harmonic wave



FT is a delta-function properly shifted

Discrete Fourier Transform

$$F_n \equiv \sum_{k=0}^{N-1} f_k e^{-2\pi i n k/N}.$$

$$f_k = \frac{1}{N} \sum_{n=0}^{N-1} F_n e^{2\pi i n k/N}.$$

sin(*wt*) →

sin(3wt) →

sin(5wt) →



Discrete Fourier Transform

- What are the frequencies in DFT?
- Which frequency is the lowest/highest one?
- Formally DFT exists for any n → periodicity. Only N samples are independent.


1st Harmonic
t
0
N-1
f
0
N/2
(Nyquist)
2nd Harmonic
t
0
N-1
f
0
N/2
(Nyquist)
Nyquist Frequency
t
0
N-1
f
0
N/2
(Nyquist)

DFT calculation

$$F_n \equiv \sum_{k=0}^{N-1} f_k e^{-2\pi i n k/N}$$

Directly - *O(N^2)*

FFT - O(N logN)

(Cooley and Tookey 1965, Gauss 1866)

Compare to the complexity of convolution.

2D Fourier transform

$$\{\phi_{uv}(x,y)\} = \{e^{-2\pi i(ux+vy)}\}$$

$$F(u,v) = \int\int_{\mathrm{R}^2} f(x,y)e^{-2\pi i(ux+vy)}\mathrm{d}x\mathrm{d}y$$

$$f(x,y) = \int\int_{\mathrm{R}^2} F(u,v)e^{2\pi i(ux+vy)}\mathrm{d}u\mathrm{d}v$$

2D Fourier transform is separable

Visualisation



Amplitude

Phase

Visualisation



ampl



log(ampl + 1)

Basis functions of 2D FT



Real part, u=v

Rectangular pulse in 2D







Other important functions



Discrete convolution theorem

Convolution mask

... holds for periodic convolution only!

Discrete convolution via FT



Zero padding to the same size

DFT calculation of both

Multiplication of the spectra

Inverse DFT

Filtering in the Fourier domain

$$\mathcal{F}[f * g] = \mathcal{F}[f]\mathcal{F}[g]$$



high pass



Gaussian high pass

band pass

low pass



Gaussian low pass



directional









Discrete shift theorem





What is more important?

The phase!

Image whitening

$$w(x,y) = \mathcal{F}^{-1}\frac{F(u,v)}{|F(u,v)|}$$

original image

“whitened” image

Díky, pro dnešek

končíme s FT!

Nějaké otázky ?