病 理 学 题 库

感谢收集题目的学长学姐们

如果你刷题时遇到了以下问题：

- 这个字/单词好像打错了
- 这道题的答案好像有问题
- 这道题题目中说有图片，实际上却无图无真相
- 这里忘记加上下标了
- 这里是物种的学名/人类基因名（或其它需要斜体的东西），却没有斜体
- 其它你认为应该修改的问题

请发邮件给yuanshuai0217@yeah.net，我会在下个版本修正错误。

目录

1 细胞和组织的损伤 1

中文题 1

英文题 17

答案 31

2 局部血液及体液循环障碍 32

中文题 32

英文题 48

答案 60

3 炎症 61

中文题 61

英文题 77

答案 89

4 肿瘤 90

中文题 90

英文题 106

答案 121

5 心血管疾病 122

中文题 122

英文题 139

答案 154

6 呼吸系统疾病 155

中文题 155

英文题 171

答案 187

7 消化系统疾病 189

中文题 189

英文题 205

答案 238

8 泌尿系统疾病 239

中文题 239

英文题 256

答案 277

9 生殖系统疾病 278

中文题 278

英文题 294

答案 313

10 淋巴造血系统疾病 314

中文题 314

英文题 323

答案 333

11 内分泌系统疾病 334

中文题 334

英文题 342

答案 347

12 神经系统疾病 348

中文题 348

英文题 356

答案 361

13 传染病 362

中文题 362

英文题 377

答案 378

1　细胞和组织的损伤

中文题

1. 不属于细胞适应性改变的是
   A. atrophy
   B. pypertrophy
   C. hyperplasia
   D. metaplasia
   E. degeneration
2. 属于不可复性损伤的是
   A. fatty degeneration
   B. cell edema
   C. mucoid degeneration
   D. amyloidosis
   E. necrosis
3. 结节性动脉炎血管壁出现的红染、纤维素样物质，称为
   A. apoptosis
   B. necrosis
   C. gangrene
   D. caseous necrosis
   E. fibrinous necrosis
4. 心脏褐色萎缩，心肌细胞内的黄褐色物质是
   A. lipofuscin
   B. melanin
   C. hemosiderin
   D. calcification
   E. bilirubin
5. 黏膜坏死脱落而遗留的深达黏膜下层的缺损，称为
   A. erosion
   B. fistula
   C. sinus
   D. cavity
   E. ulcer
6. 表皮细胞属于
   A. stable cell
   B. labile cell
   C. permanent cell
   D. stroma cell
   E. stem cell
7. 细胞内、血管壁和结缔组织内出现均质、红染、玻璃样物质，称为
   A. fatty degeneration
   B. hyaline degeneration
   C. amyloid degeneration
   D. mucoid degeneration
   E. hydropic degeneration
8. 一种分化成熟组织取代另外一种分化成熟组织，称为
   A. atrophy
   B. hypertrophy
   C. hyperplasia
   D. metaplasia
   E. degeneration
9. 不属于萎缩的是
   A. 80 岁老人的脑
   B. 晚期癌症患者的心脏
   C. 空洞型肾结核的肾实质
   D. 小儿麻痹症的下肢肌肉

E. 恶病质患者的脂肪组织

10. 癌症患者首先发生萎缩的组织是

A. 脂肪组织

B. 骨骼肌

C. 心肌

D. 肝细胞

E. 神经元

11. 符合萎缩的描述是

A. 器官体积小于正常即为萎缩

B. 萎缩的器官间质增生

C. 萎缩细胞功能正常

D. 萎缩器官的细胞数目正常

E. 供血断绝引起萎缩

12. 符合萎缩的病变是

A. 淋巴结结核

B. Di George 综合征的胸腺

C. 老年妇女的子宫内膜

D. 先天性食管闭锁

E. 主动脉夹层

13. 不属于萎缩的是

A. 高血压的左心室

B. 80 岁老年人的脑

C. 慢性高血压的肾实质

D. 脑积水的脑实质

E. 晚期癌症患者的心脏

14. 食管癌患者体内最先萎缩的是

A. 心肌细胞

B. 骨骼肌细胞

C. 肝细胞

D. 神经细胞

E. 脂肪细胞

15. 属于生理性萎缩的是

A. 垂体坏死引起的乳腺退缩

B. 青年男性睾丸生精细胞大量减少

C. 小儿麻痹引起的下肢萎缩

D. 青春期胸腺萎缩

E. 动脉粥样硬化症引起的脑萎缩

16. 不属于萎缩的器官是

A. 高血压病的肾脏

B. 肾切除术后，保留的对侧肾脏

C. 慢性肾小球肾炎的肾脏

D. 肾盂积水的肾脏

E. 动脉粥样硬化症的肾脏

17. 与肝脂肪变性无关的是

A. 酗酒

B. 四氯化碳中毒

C. 摄入过多的脂肪

D. 糖尿病

E. 白喉杆菌感染

18. 不符合脂肪变性的描述是

A. 槟榔肝时，肝小叶周边区脂肪变性最明显

B. 磷中毒时，肝小叶中央区脂肪变性最明显

C. 砷中毒时，可引起心肌脂肪变性

D. 白喉时，可引起“虎斑心”

E. 严重的贫血时，肾近曲管上皮可出现脂肪变性

19. 不符合脂肪变性的是

A. 肝细胞内出现大小不等的脂滴

B. 巨噬细胞内出现大小不等的脂滴

C. 心肌细胞内出现大小不等的脂滴

D. 脂肪肉瘤细胞内出现大小不等的脂滴

E. 近曲小管上皮细胞内出现大小不等的脂滴

20. 不引起肝细胞脂肪变性的是

A. 酒精中毒
B. 四氯化碳中毒
C. 糖尿病
D. 白喉
E. 高脂肪饮食

21. 关于肝脂肪变性的描述，正确的是

A. 肝淤血时，脂肪变性首先发生在小叶周围
B. 磷中毒时，脂肪变性主要发生在小叶中央区
C. 轻度肝脂肪变性时，肉眼下肝可无明显改变
D. 重度肝脂肪变性时，肝脏缩小
E. 脂肪变性的肝细胞的细胞浆疏松，呈空网状

22. 关于心肌脂肪变性的描述，错误的是

A. 白喉外毒素可引起心肌脂肪变性
B. 磷、砷、氯仿可引起心肌脂肪变性
C. 严重贫血可引起心肌脂肪变性
D. 心肌脂肪变性严重影响心脏功能
E. 心肌内出现串珠状、细小脂肪空泡

23. 关于肾脂肪变性的描述，错误的是

A. 与肾小球毛细血管通透性升高有关
B. 脂肪变性主要累及远曲管上皮细胞
C. 可见于中毒
D. 可见于脂性肾病
E. 可见于严重贫血

24. 脂肪变性时，脂滴位于

A. 高尔基器
B. 线粒体
C. 溶酶体
D. 内质网
E. 胞浆基质

25. 纤维素样坏死不见于

A. 风湿病
B. 恶性高血压
C. 动脉粥肿
D. 系统性红斑狼疮
E. 类风湿性关节炎

26. 纤维素样坏死的病变内不出现

A. 粘多糖
B. 免疫球蛋白
C. 纤维蛋白
D. 基底膜样物质
E. 胶原崩解产物

27. 结节性多动脉炎动脉壁的红染颗粒状物质是

A. 水变性
B. 脂肪变性
C. 玻璃样变性
D. 淀粉样变性
E. 纤维素样坏死

28. 纤维组织内颗粒状、团块状、强嗜酸红染的无结构物质，状似纤维素，此病变称为

A. 水变性
B. 脂肪变性
C. 玻璃样变性
D. 纤维素样坏死
E. 钙化

29. 纤维素样坏死可见于

A. 风湿病
B. 良性高血压
C. 动脉粥肿
D. 瘢痕
E. 甲状腺功能低下

30. 纤维素样坏死可见于
A. 良性高血压
B. 恶性高血压
C. 动脉粥肿
D. 瘢痕
E. 甲状腺功能低下

31. 纤维素样坏死可见于
A. 良性高血压
B. 甲状腺功能低下
C. 动脉粥肿
D. 系统性红斑狼疮
E. 瘢痕

32. 纤维素样坏死可见于
A. 良性高血压
B. 瘢痕
C. 动脉粥肿
D. 甲状腺功能低下
E. 类风湿性关节炎

33. 玻璃样变性不发生于
A. 结缔组织
B. 血管壁
C. 浆细胞
D. 肝细胞
E. 肾远曲小管上皮细胞

34. 结缔组织玻璃样变的错误描述是
A. 常见于瘢痕组织
B. 常见于纤维化肾小球
C. 纤维母细胞增多
D. 胶原纤维融合为梁状、带状或片状
E. 常见于粥瘤

35. 血管壁玻璃样变多见于
A. 大动脉
B. 中型动脉
C. 细动脉
D. 大静脉
E. 小静脉

36. 关于血管壁玻璃样变，错误的描述是
A. 常见于高血压病
B. 常见于肾、脑、脾、视网膜细动脉
C. 血管壁变硬、管腔狭窄
D. 可引起组织缺血
E. 玻璃样物质由免疫球蛋白和基底膜样物构成

37. 关于细胞内玻璃样变错误的描述是
A. 肾近曲小管上皮内的玻璃样小滴由重吸收的血浆蛋白构成
B. 近曲管上皮玻璃样变多见于肾盂肾炎
C. 肝细胞内的 Mallory 小体由前角蛋白构成
D. Mallory 小体多见于酒精中毒时
E. 浆细胞内的玻璃样小体由免疫球蛋白构成

38. 女，40 岁，患者因血尿、蛋白尿、管型尿，经治疗无明显好转。四年后，因尿毒症死亡。尸体解剖发现两肾对称性萎缩，表面呈细颗粒状。组织学检查，大量肾小球纤维化，并呈现均匀红染，半透明小团。这些肾小球的病变属于

A. 纤维素样坏死
B. 淀粉样变性
C. 玻璃样变性
D. 脂肪变性
E. 水变性

39. 男，30 岁，酗酒时间约 8 年余，日消费北京二锅头酒约 500ml。近半年来经常出现恶心、呕吐、食欲不振。肝穿刺显示肝细胞广泛脂肪变性，肝细胞灶状坏死。肝细胞内可见大小不等的红染的半透明小体。这些透明小体为
A. 前角蛋白微丝
B. 免疫球蛋白
C. 血浆蛋白
D. 乙型肝炎病毒表面抗原
E. 肝糖原

40. 男，55 岁，高血压病史已 20 余年。手术切除脾脏，切片可见脾中央小动脉管壁增厚，管腔狭窄，管壁内可见均匀红染、半透明物质。这些半透明物质为
A. 淀粉样物质
B. 血浆蛋白和基底膜样物质
C. 胶原纤维
D. 免疫球蛋白
E. 脂蛋白

41. 不符合淀粉样物质的描述是
A. 组织化学反应与淀粉相似
B. 丙种球蛋白沉积而成
C. 电镜下呈纤细丝状
D. 镜下呈淡红色均质状
E. 结合粘多糖的蛋白质

42. 不符合淀粉样物质的描述是
A. 可见于骨髓瘤
B. 可见于结核病
C. 可见于慢性化脓性炎
D. 可见于慢性炎症
E. 可见于结节性多动脉炎

43. 不符合变性的病变是
A. 间质内类粘液的积聚
B. 肝细胞内出现脂滴
C. 血管壁出现半透明、均匀红染的物质
D. 心肌细胞内水分增多
E. 细胞核破碎

44. 粘液样变性常见于
A. 甲状腺功能低下的皮下组织
B. 高血压的心脏
C. 硬癌的间质
D. 颗粒性固缩肾
E. 瘢痕

45. 慢性乙型肝炎时，电镜下毛玻璃细胞胞浆内可见
A. 扩张的光面内质网
B. Russell 小体
C. Mallory 小体
D. 脂质小体
E. 残存小体

46. 细胞内线粒体肿大、内质网扩张常见于
A. 水变性
B. 脂肪变性
C. 玻璃样变性
D. 纤维素样变性
E. 钙化

47. 肝细胞气球样变属于

A. 水变性
B. 脂肪变性
C. 玻璃样变性
D. 淀粉样变性
E. 纤维素样变性

48. 符合变性的描述是
A. 属不可复性损伤
B. 细胞功能丧失
C. 表现为异常物质的出现或正常物质显著增多
D. 细胞体积缩小，伴功能下降
E. 细胞体积增大，伴功能增强

49. 干酪样坏死灶周边部钙盐沉积称为
A. 水变性
B. 脂肪变性
C. 玻璃样变性
D. 纤维素样变性
E. 钙化

50. 形成脂褐素的细胞器是
A. 高尔基器
B. 粗面内质网
C. 光面内质网
D. 溶酶体
E. 线粒体

51. 不符合含铁血黄素的描述是
A. 常见于心肌细胞核两侧
B. 光镜下，呈棕黄色，具有折光性
C. 血红蛋白转化而来
D. 普鲁士蓝反应阳性
E. 常见于出血灶周围

52. 男，60 岁，因胃癌广泛转移而死亡。尸体解剖检查，心脏重 200g。切片检查心肌细胞核两侧可见黄褐色细颗粒状色素，此色素可能是
A. 黑色素
B. 胆色素
C. 含铁血黄素
D. 脂褐素
E. 前角蛋白聚集

53. 脂褐素电镜下为
A. 扩张的光面内质网
B. Russell 小体
C. Mallory 小体
D. 脂质小体
E. 残存小体

54. 干酪样坏死灶周围钙盐沉积，称为
A. 溶解吸收
B. 分离排出
C. 机化
D. 钙化
E. 包裹

55. 硬脑膜出现了骨组织属于
A. 肥大
B. 增生
C. 再生
D. 化生
E. 机化

56. 甲状腺瘤间质钙化灶中有骨组织属于
A. 肥大
B. 增生
C. 再生
D. 化生
E. 机化

57. 肠扭转引起的坏死，称为

A. 凝固性坏死
B. 液化性坏死
C. 气性坏疽
D. 干性坏疽
E. 湿性坏疽

58. 深部开放损伤合并厌气菌感染，称为
A. 凝固性坏死
B. 液化性坏死
C. 气性坏疽
D. 干性坏疽
E. 湿性坏疽

59. 产后子宫感染引起的坏死属于
A. 凝固性坏死
B. 干酪样坏死
C. 液化性坏死
D. 湿性坏疽
E. 脂肪坏死

60. 气性坏疽是指
A. 结核杆菌感染
B. 坏死组织伴腐败菌感染
C. 葡萄球菌感染
D. 深部组织的产气荚膜杆菌、恶性水肿杆菌及腐败弧菌等厌气菌感染
E. 溶血性链球菌感染

61. 后果严重需紧急处理的坏死是
A. 乳房脂肪坏死
B. 下肢干性坏疽
C. 臀部气性坏疽
D. 肾局灶性梗死
E. 脾梗死

62. 不符合湿性坏疽的描述是
A. 合并腐败菌感染
B. 具有恶臭味
C. 常有全身中毒症状
D. 坏疽组织呈污秽黑褐色
E. 坏死组织与健康组织之间界限清楚

63. 男性，23 岁，24 小时前因车祸引起左下肢胫骨及腓骨开放性、粉碎性骨折，入院清除固定后 12 小时，出现左小腿及左足青紫、肿胀，患者出现发热和全身不适。次日肿胀加剧，累及左大腿，伤口流出伴有气泡的的恶臭脓汁，肿胀区按之有捻发音。左小腿的坏死为
A. 干性坏疽
B. 湿性坏疽
C. 气性坏疽
D. 液化性坏死
E. 干酪样坏死

64. 湿性坏疽常发生在
A. 脑、脾、肝
B. 脑、肠、子宫
C. 肺、肠、肝
D. 肺、肠、子宫
E. 肺、肾、脑

65. 不符合干性坏疽的描述是
A. 合并腐败菌感染
B. 常伴有全身中毒症状
C. 坏死组织呈污秽黑褐色
D. 坏死组织与健康组织之间界限清楚
E. 坏死部位动脉受阻，而静脉回流通畅

66. 男，68 岁，糖尿病史 10 年。近一年来左下肢行走后疼痛，休息后好转。近两个月左拇趾变黑，皱缩，知觉消失。此

种坏死是

A. 固缩坏死

B. 干性坏疽

C. 湿性坏疽

D. 气性坏疽

E. 液化性坏死

67. 下列哪个脏器不发生坏疽？

A. 肺

B. 下肢

C. 阑尾

D. 小肠

E. 脑

68. 血栓闭塞性脉管炎累及左下肢，引起的坏死，称为

A. 凝固性坏死

B. 液化性坏死

C. 气性坏疽

D. 干性坏疽

E. 湿性坏疽

69. 属于干性坏疽的是

A. 肠扭转引起的肠坏疽

B. 化脓性感染引起的肺坏疽

C. 坏疽性阑尾炎

D. 糖尿病患者的下肢坏疽

E. 坏疽性子宫内膜炎

70. 属于干性坏疽的是

A. 肠扭转引起的肠坏疽

B. 化脓性感染引起的肺坏疽

C. 坏疽性阑尾炎

D. 糖尿病患者的下肢坏疽

E. 坏疽性子宫内膜炎

71. 不符合坏疽的描述是

A. 常伴有腐败菌感染

B. 呈黑褐色

C. 干性坏疽病变境界较清楚

D. 湿性坏疽常伴全身中毒症状

E. 气性坏疽较常见

72. 符合干性坏疽的描述是

A. 合并腐败菌感染

B. 常伴有全身中毒症状

C. 局部肿胀明显

D. 坏死组织与健康组织之间界限不清楚

E. 坏死部位动脉和静脉血流均受阻

73. 肺结核瘤属于

A. 溶解吸收

B. 分离排出

C. 机化

D. 钙化

E. 包裹

74. 腕关节结核死骨形成后将发生

A. 溶解吸收

B. 分离排出

C. 机化

D. 钙化

E. 包裹

75. 大叶性肺炎的痊愈是通过

A. 溶解吸收

B. 分离排出

C. 机化

D. 钙化

E. 包裹

76. 心肌梗死病灶由纤维组织替代属于

A. 溶解吸收

B. 分离排出
C. 机化
D. 钙化
E. 包裹

77. 病毒性肝炎肝细胞灶状坏死将发生
A. 溶解吸收
B. 分离排出
C. 机化
D. 钙化
E. 包裹

78. 属于机化的病变是
A. 较深的粘膜缺损
B. 结核瘤
C. 肺肉质变
D. 肾空洞
E. 肺钙化灶

79. 不属于机化的是
A. 粘膜糜烂愈合
B. 脾梗死瘢痕形成
C. 血管内血栓再通
D. 纤维素性粘连转变为纤维性粘连
E. 脑软化灶形成胶质瘢痕

80. 窦道是指
A. 皮肤或粘膜坏死组织脱落形成的缺损
B. 深部组织坏死穿出皮肤形成的病理性盲管
C. 空腔脏器与体表之间的病理性管道
D. 内脏器官坏死液化，并经相应管道排出体外，留下的空腔
E. 肉芽组织取代坏死组织的过程

81. 不属于化生的是
A. 支气管粘膜出现鳞状上皮
B. 胃粘膜内出现胰腺组织
C. 胃粘膜内出现肠上皮
D. 食管粘膜内出现胃上皮
E. 胆囊粘膜内出现鳞状上皮

82. 与化生有关的肿瘤是
A. 卵巢甲状腺肿
B. 肺鳞状细胞癌
C. 阴道横纹肌肉瘤
D. 直肠黑色素瘤
E. 子宫颈腺癌

83. 不属于化生的是
A. 胃粘膜内出现肠上皮
B. 食道粘膜内出现胃上皮
C. 骨化性肌炎内出现骨组织
D. 膀胱粘膜内出现鳞状上皮
E. 胆囊肌层内出现腺上皮

84. 不属于化生的病变是
A. 食管粘膜出现柱状上皮
B. 胃粘膜中出现胰腺上皮
C. 膀胱粘膜出现鳞状上皮
D. 胆囊粘膜出现鳞状上皮
E. 子宫颈粘膜出现柱状上皮

85. 男，40 岁，吸烟史 20 年，每日 2～3 包。近日咳嗽加重，X 线显示左上肺阴影。支气管镜粘膜活检，可见鳞状上皮。此种病理变化属于
A. 支气管粘膜萎缩
B. 支气管粘膜鳞状上皮化生
C. 支气管粘膜鳞状上皮增生
D. 支气管鳞状细胞癌
E. 支气管粘膜肥大

86. Barrett 食管内出现的腺上皮，称为
A. 鳞状上皮化生
B. 腺性化生
C. 肠上皮化生
D. 骨化生
E. 髓性化生

87. 乳腺增生时出现大汗腺腺体，称为
A. 鳞状上皮化生
B. 腺性化生
C. 肠上皮化生
D. 骨化生
E. 髓性化生

88. 慢性萎缩性胃炎胃粘膜出现杯状细胞，称为
A. 鳞状上皮化生
B. 腺性化生
C. 肠上皮化生
D. 骨化生
E. 髓性化生

89. 胃溃疡的愈复
A. 肥大
B. 增生
C. 再生
D. 化生
E. 机化

90. 男性，6 岁，在家长未注意时，被铡刀将左手铡断，经断肢再植后，左手功能恢复，下列哪种组织属完全再生？
A. 皮肤
B. 神经
C. 肌腱
D. 动脉
E. 静脉

91. 男性，50 岁，因肝右叶肝癌而行肝叶切除术，患者的肝脏发生的变化是
A. 经过较长时间，恢复原结构
B. 保持手术后剩余肝脏的状态
C. 肝小叶不断增生
D. 形成若干再生结节
E. 纤维组织大量增生

92. 男性，20 岁，剧烈运动时，不慎将左踝骨骨折，符合骨折愈合的过程是
A. 骨折处由软骨代替
B. 骨折处由纤维组织代替
C. 骨折处由骨组织代替，但不能恢复原状
D. 骨折处经血肿形成、纤维性骨痂、骨性骨痂、骨痂改建几个阶段达到完全愈合
E. 骨折处由纤维组织、软骨组织、骨组织混合增生代替

93. 男性，50 岁，右腹股沟疝，经疝修补术后，伤口抗拉力强度达到顶点的时间是
A. 一周
B. 三周
C. 五周
D. 二个月
E. 三个月

94. 在创伤愈合过程中，胶原纤维的分解吸收出现在
A. 创伤后 15 天以内
B. 创伤后一个月之后
C. 早期多，晚期少
D. 早期少，晚期多

E. 以上都不对

95. 创伤伤口的收缩是在

A. 受伤后 2、3 天

B. 受伤后 4、5 天

C. 受伤后 6、7 天

D. 受伤后 8、9 天

E. 受伤后 10 天

96. 属于完全性再生的是

A. 胃溃疡愈合

B. 皮肤溃疡愈合

C. 肺空洞愈合

D. 肝细胞点、灶状坏死愈合

E. 心肌梗死灶的修复

97. 肉芽组织是

A. 含有肉芽肿的组织

B. 含有肌肉再生的嫩芽

C. 含有炎细胞的组织

D. 含有多核巨细胞的组织

E. 含有新生毛细血管和纤维母细胞的组织

98. 不含有多核巨细胞的病变是

A. 伤寒小结

B. 结核肉芽肿

C. 类肉瘤肉芽肿

D. 麻风肉芽肿

E. 异物性肉芽肿

99. 肉芽组织的主要成分是

A. 中性粒细胞

B. 淋巴细胞

C. 毛细血管和纤维母细胞

D. 巨噬细胞

E. 异物巨细胞

100. 肉芽组织不具备的功能有

A. 机化凝血块和坏死组织

B. 包裹异物

C. 抗感染和保护创面

D. 分解、液化坏死物

E. 填补伤口及组织缺损

101. 不符合一期愈合的是

A. 创缘整齐

B. 炎症反应重

C. 凝血块少

D. 5～6 天胶原纤维形成

E. 约 3～6 周完全愈合

102. 符合二期愈合的是

A. 组织缺损少

B. 创缘不整并伴有感染

C. 愈合后留下线状瘢痕

D. 约 3～6 周完全愈合

E. 炎症反应轻

103. 有利于伤口愈合的因素是

A. 感染

B. 异物

C. 电离辐射

D. 老年

E. 营养充分及局部血液循环良好

104. 延缓伤口愈合的因素是

A. 青少年

B. 适量维生素及蛋白质补充

C. 保持创伤处血液循环畅通

D. 电离辐射

E. 清除坏死组织，预防感染

105. 不属于肥大的是

A. 高血压病的左心室

B. 脑积水
C. 前列腺肥大患者的膀胱壁增厚
D. 健美运动员的骨骼肌
E. 一侧肾脏切除后对侧肾脏的增大

106. 授乳期的乳腺增大属于
A. 肥大
B. 增生
C. 再生
D. 化生
E. 机化

107. 属于代偿性肥大的是
A. 妊娠子宫
B. 哺乳期乳腺
C. 甲状腺机能亢进患者的甲状腺肿大
D. 地方性甲状腺肿
E. 一侧肾摘除，对侧肾肥大

108. 增生是指
A. 器官和组织体积缩小
B. 器官和组织体积增大
C. 一种分化组织转化为另一种相似性质的分化组织
D. 由于实质细胞增多而造成器官、组织体积增大
E. 器官和组织先天发育异常

109. 不属于内分泌性增生的是
A. 地方性甲状腺肿
B. 乳腺增生症
C. 肝细胞结节状增生
D. 前列腺增生症
E. 子宫内膜增生症

110. ACTH 分泌过多致肾上腺皮质形成多数小结节
A. 肥大
B. 增生
C. 再生
D. 化生
E. 机化

111. 妊娠的子宫
A. 肥大
B. 增生
C. 再生
D. 化生
E. 机化

112. 男性乳腺发育
A. 肥大
B. 增生
C. 再生
D. 化生
E. 机化

113. 肝细胞可以通过完全性再生修复的病变是
A. 肝细胞桥接坏死
B. 肝细胞大片坏死
C. 肝细胞反复坏死
D. 肝细胞亚大片坏死
E. 肝细胞点状坏死

114. 属于永久性细胞的是
A. 神经细胞
B. 肝细胞
C. 胰腺细胞
D. 表皮细胞
E. 淋巴细胞

115. 属于不稳定细胞的是
A. 神经细胞和心肌细胞

B. 汗腺和皮脂腺的上皮细胞
C. 延髓和内分泌腺的腺上皮细胞
D. 骨母细胞和平滑肌细胞
E. 表皮细胞、呼吸道及消化道粘膜上皮细胞

116. 属于稳定细胞的是
A. 男性及女性生殖管道被覆上皮细胞
B. 淋巴造血细胞
C. 肾小管上皮细胞及肝细胞
D. 神经细胞、心肌细胞和骨骼肌细胞
E. 表皮细胞和间皮细胞

117. 再生力最强的细胞是
A. 骨骼肌细胞
B. 神经节细胞
C. 心肌细胞
D. 神经胶质细胞
E. 软骨细胞

118. 符合组织再生力强弱的描述是
A. 成纤维细胞＞神经细胞＞肝细胞
B. 软骨细胞＞骨细胞＞肾小管上皮细胞
C. 骨细胞＞肾小管上皮细胞＞神经细胞
D. 鳞状上皮细胞＞横纹肌细胞＞神经细胞
E. 肾小管上皮细胞＞骨髓细胞＞脂肪细胞

119. 符合肌肉组织的再生的是
A. 骨骼肌肌原纤维部分受损不能完全再生
B. 骨骼肌肌纤维断裂后仍可完全再生
C. 心肌梗死后由瘢痕修复
D. 肠吻合后，肌层可经再生恢复原结构
E. 肌型动脉吻合后，肌层可经再生恢复原结构

120. 符合间叶组织再生的描述是
A. 纤维母细胞具有很强的再生能力
B. 软骨细胞具有很强的再生能力
C. 骨骼肌细胞具有很强的再生能力
D. 平滑肌细胞具有很强的再生能力
E. 心肌细胞具有很强的再生能力

121. 女性，30 岁，腰椎结核并在背部形成冷脓肿，冷脓肿破溃，破溃处流脓不断，此病变是
A. 空洞
B. 瘘管
C. 溃疡
D. 窦道
E. 糜烂

122. 男性，26 岁，因肛旁脓肿，左肛周形成伤口，局部流脓不断，入院后检查，肛旁伤口为外口，实与直肠相通，次病理管道为
A. 空洞
B. 瘘管
C. 窦道
D. 溃疡
E. 糜烂

123. 大叶性肺炎肉质变属于
A. 肥大
B. 增生
C. 再生
D. 化生

E. 机化

124. 表层肠粘膜坏死脱落后形成

A. 糜烂

B. 溃疡

C. 瘘管

D. 包裹

E. 窦道

125. 烫伤引起的深达皮下脂肪的皮肤缺损，称为

A. 糜烂

B. 溃疡

C. 瘘管

D. 包裹

E. 窦道

126. 子宫颈癌累及膀胱引起尿液经阴道排出，称为

A. 糜烂

B. 溃疡

C. 瘘管

D. 包裹

E. 窦道

127. 肾结核坏死物经尿道排出，所形成的病变为

A. 糜烂

B. 溃疡

C. 瘘管

D. 空洞

E. 窦道

128. 慢性化脓性骨髓炎脓液穿破皮肤形成的病变为

A. 糜烂

B. 溃疡

C. 瘘管

D. 包裹

E. 窦道

129. 病毒性肝炎时，肝细胞嗜酸性小

A. 凝固性坏死

B. 干酪样坏死

C. 液化性坏死

D. 湿性坏疽

E. 凋亡

130. 细胞坏死最具特征的形态学标志是

A. 胞浆内糖原减少

B. 溶酶体破裂

C. 核固缩、碎裂及溶解

D. 胞浆内出现微细结构破坏

E. 细胞体积变小，染色变浅

131. 坏死是指

A. 局部组织、细胞的死亡

B. 机体死亡后组织、细胞出现的形态学改变

C. 活体局部组织、细胞死亡后出现的形态学改变

D. 机体内细胞的程序性死亡

E. 细胞的可复性改变

132. 坏死的主要形态学标志是

A. 胞浆红染

B. 核浓缩、核破碎、核溶解

C. 间质崩解

D. 保留组织结构的轮廓

E. 核内假包含体形成

133. 不符合固缩坏死的描述是

A. 常表现为单个细胞或小灶状坏死

B. 各种损伤性刺激改变遗传信息的转

录和翻译，形成死亡蛋白

C. 核酸内切酶被激活，引起染色质凝集

D. 凋亡小体形成

E. 引起强烈炎症反应

134. 不属于坏死标志的是

A. 核碎裂

B. 组织结构崩解

C. 核浓缩

D. 核内出现假包含体

E. 核溶解

135. 不符合固缩坏死的描述是

A. 多为细胞的生理性死亡

B. 多为单个分散的细胞

C. 常形成凋亡小体

D. 常伴有炎症反应

E. 亦可见于病理过程

136. 符合坏死的描述是

A. 局部组织的死亡

B. 细胞的生理性死亡

C. 活体局部组织死亡所发生的一系列形态改变

D. 死亡后组织发生的一系列形态改变

E. 单一细胞的死亡

137. 不符合干酪样坏死的描述是

A. 多由结核杆菌引起

B. 淡黄色、干燥、易碎

C. 镜下呈红染、颗粒状、无结构的物质

D. 含有多量脂质

E. 可见原有结构影像

138. 属于液化性坏死的是

A. 心肌梗死

B. 脑梗死

C. 肠梗死

D. 淋巴结干酪样坏死

E. 脚坏疽

139. 不属于液化性坏死的是

A. 脾梗死

B. 脑梗死

C. 坏死性胰腺炎

D. 脂肪坏死

E. 脓肿

140. 不符合干酪样坏死的描述是

A. 呈淡黄色，均匀细腻

B. 状似纤维蛋白

C. 可发生钙化

D. 可发生液化

E. 常由结核杆菌引起

141. 液化性坏死常见于

A. 脑和胰腺

B. 肝

C. 脾

D. 肾

E. 小肠

142. 女，24 岁，腋窝淋巴结肿大约半年，手术切除。手术标本为直径约 2cm 的卵圆形淋巴结，切面大部为淡黄色、质软、易碎的坏死灶。镜下检查，坏死灶为红染的、颗粒状、无结构的物质，坏死灶周围可见类上皮细胞和多核巨细胞。以上所描述的坏死是

A. 梗死

B. 干性坏死

C. 气性坏死
D. 干酪样坏死
E. 液化性坏死

143. 心肌梗死属于
A. 凝固性坏死
B. 液化性坏死
C. 气性坏疽
D. 干性坏疽
E. 湿性坏疽

144. 脑梗死属于
A. 凝固性坏死
B. 液化性坏死
C. 气性坏疽
D. 干性坏疽
E. 湿性坏疽

145. 阿米巴病时组织发生
A. 凝固性坏死
B. 干酪样坏死
C. 液化性坏死
D. 固缩坏死
E. 湿性坏疽

146. 结核病时发生
A. 凝固性坏死
B. 干酪样坏死
C. 液化性坏死
D. 湿性坏疽
E. 脂肪坏死

147. 重型病毒性肝炎时，肝细胞大片坏死属于
A. 凝固性坏死
B. 干酪样坏死
C. 液化性坏死
D. 湿性坏疽
E. 脂肪坏死

148. 脑脓肿为
A. 凝固性坏死
B. 干酪样坏死
C. 液化性坏死
D. 湿性坏疽
E. 脂肪坏死

149. 急性坏死性胰腺炎的胰腺坏死属于
A. 凝固性坏死
B. 干酪样坏死
C. 液化性坏死
D. 湿性坏疽
E. 脂肪坏死

150. 肾贫血性梗死灶为
A. 凝固性坏死
B. 干酪样坏死
C. 液化性坏死
D. 湿性坏疽
E. 脂肪坏死

151. 不符合脂肪坏死的描述是
A. 是一种特殊的液化性坏死
B. 酶解性脂肪坏死常见于胰腺炎
C. 外伤性脂肪坏死常见于乳腺
D. 坏死灶内可见多量泡沫细胞
E. 坏死灶内可见干酪样物质

152. 不符合液化性坏死的描述是
A. 局部蛋白质含量少
B. 局部蛋白酶丰富
C. 局部脂质含量多
D. 局部化脓
E. 脂肪坏死部署于液化性坏死

153. degeneration
A. 萎缩
B. 肥大
C. 变性
D. 坏死
E. 化生

154. necrosis
A. 萎缩
B. 肥大
C. 变性
D. 坏死
E. 化生

155. gangrene
A. 坏死
B. 坏疽
C. 瘘管
D. 包裹
E. 窦道

156. atrophy
A. 肥大
B. 增生
C. 萎缩
D. 包裹
E. 窦道

157. hypertrophy
A. 糜烂
B. 溃疡
C. 增生
D. 包裹
E. 肥大

158. hyperplasia
A. 糜烂
B. 化生
C. 增生
D. 包裹
E. 变性

159. metaplasia
A. 增生
B. 肥大
C. 坏死
D. 化生
E. 变性

160. granulation tissue
A. 梗死组织
B. 增生组织
C. 变性组织
D. 肉芽组织
E. 坏死组织

英文题

1. Which of the following cells is most likely to have the highest telomerase activity, which is positively with sustained cellular proliferation?
A. Neurons
B. Neutrophils
C. Endothelial cells
D. Germ cells
E. Erythrocytes

2. The diameter of a young man’s left thigh has decreased after a femoral fracture caused by trauma, which immobilized

him for one month. This change is most likely to result from which of the following alterations in the thigh muscles?

A. Necrosis
B. Hypoplasia
C. Dystrophy
D. Atrophy
E. Hyaline degeration

3. Histologic sections of lung tissue from a 72-year-old woman with congestive heart failure and progressive breathing problems reveal numerous hemosiderin-laden cells within the alveoli. Which of the following is the cell of origin of these“heart failure cells”?

A. Endothelial cells
B. Macrophages
C. Eosinophils
D. Lymphocytes
E. Pneumocytes

4. Which feature is not necessarily associated with the healing of open wounds:

A. the formation of granulation tissue
B. infection
C. migration of the surrounding epithelium
D. giant cell formation
E. contraction

5. The healing of a wound will be inhibited by the following, with the exception of:

A. vitamin K deficiency
B. starvation
C. the administration of glucocorticoids
D. lack of blood supply
E. infection

6. Wound healing is not enhanced by the administration of:

A. cortisol (皮质醇)
B. zinc
C. growth factors
D. oxygen
E. vitamin C

7. An old man suffered a severe stroke (脑卒中), and two months later, his head CT scan showed a large cystic area in the parietal lobe. Which of the following pathologic processes has most likely occurred in his brain?

A. Coagulative necrosis
B. Fat necrosis
C. Apoptosis
D. Karyolysis
E. Liquefactive necrosis

8. The healing of an incised wound is not associated with the following:

A. a lag phase
B. a demolition (破坏) phase
C. a proliferative phase
D. a contractile phase
E. a maturation phase

9. General factors predisposing to wound infection do not include:

A. uncontrolled diabetes
B. hypogammaglobulinaemia (低丙种球

蛋白血症)

C. low platelet count

D. agranulocytopenia (粒细胞缺乏症)

E. eosinophilia

10. A routine H&E histologic section from an irregular white area within the anterior wall of the heart of a 71-year-old man who died secondary to ischemic heart disease reveals the myocytes to be replaced by diffuse red material. This material stains blue with a trichrome stain. Which of the following statements correctly describes this material?

A. It is secreted by endothelial cells and links macromolecules to integrins

B. It is secreted by fibroblasts and has a high content of glycine and hydroxyproline

C. It is secreted by hepatocytes and is mainly responsible for intravascular oncotic pressure

D. It is secreted by monocytes and contains a core protein that is linked to mucopolysaccharides

E. It is secreted by plasma cells and is important in mediating humoral immunity

11. A patient presents with a large wound to his right forearm that is the result of a chain saw accident. You treat his wound appropriately and follow him in your surgery clinic at routine intervals. Initially his wound is filled with granulation tissue, which is composed of proliferating fibroblasts and proliferating new blood vessels (angiogenesis). Which of the following substances is thought to be the most important growth factor involved in angiogenesis?

A. Epidermal growth factor (EGF)

B. Platelet-derived growth factor (PDGF)

C. Transforming growth factor-alpha (TGF-a)

D. Transforming growth factor-beta (TGF-P)

E. Vascular endothelial growth factor (VEGF)

12. Which one of these cells is thought to be the source of hepatic stem cells?

A. Ita cells

B. Limbus cells

C. Oval cells

D. Paneth cells

E. Satellite cells

13. A mouse model is created in which the low density lipoprotein (LDL) receptor gene is not expressed in the liver. For creating such a “knock out” mouse, which of the following cells will be most useful?

A. Embryonic stem cells

B. Hematopoietic stem cells

C. Hepatic oval cells

D. Adult bone marrow mesenchymal pro-

genitor cells

E. Regenerating hepatocytes

14. The enzyme levels (AST (谷草转氨酶) and ALT (谷丙转氨酶) finally returned to normal after a hepatitis A patient has received 1 month of treatment. Which phase of the cell cycle best describes the hepatocytes 1 month after infection?

A. G_0


B. G_1

C. S

D. G_2

E. M

15. A 24-year-old man with acute appendicitis undergoes surgical removal of the inflamed appendix. The incision site is sutured. A trichrome-stained section of the site is shown in the photograph. How long after the surgery would this appearance most likely be seen?



A. 1 day

B. 2 to 3 days

C. 4 to 5 days

D. 2 weeks

E. 1 month

16. After the sutures were removed from the wound 1 week after a surgical operation, the site became disfigured by a prominent raised, nodular scar that continued to develop over the following 2 months. Which of the following terms best describes the process that occurred during this 2-month period?

A. Secondary union

B. Keloid formation

C. Resolution

D. Dehiscence

E. Organization

17. Which of the following molecules synthesized by fibroblasts can nost readily bind to cellular integrins and extracellular collagen and attach epidermal basal cells to basement membrane?


A. Heparin

B. Dermatan sulfate

C. Procollagen

D. Fibronectin

E. Hyaluronic acid



18. A middle-aged man experienced sudden death preceeded by severe abdominal pain with hypotension and shock. From the gross appearance of the mesentery, seen in the figure, which of the following events

has most likely occurred?

A. Acute pancreatitis
B. Small intestinal perforation
C. Tuberculous peritonitis (结核性腹膜炎)
D. Gangrenous cholecystitis
E. Hepatitis B virus infection

19. Which of the following substances is most likely responsible for the interaction between the cytoskeleton and the extracellular matrix in a wound healing experiment?

A. Type IV collagen
B. Fibronectin
C. Epidermal growth factor
D. Platelet-derived growth factor
E. Integrin

20. After two months, a wound healing by secondary intention is only 10% of its original size. Which of the following processes best accounts for the observed decrease in wound size over the past 6 weeks?

A. Resolution of subcutaneous edema
B. Elaboration of adhesive glycoproteins
C. Inhibition of metalloproteinases
D. Myofibroblast contraction
E. Increase in synthesis of collagen

21. For a regular sarean section (剖宫产手术) incision, which of the following statements best describes the wound site at the time of suture removal (around 1 week after the operation)?

A. Type IV collagen predominates
B. No more wound strength will be gained
C. Wound strength is 80% of normal tissue
D. Granulation tissue is still present
E. Collagen degradation exceeds synthesis

22. In an experiment, two groups of test animals are infected, one with viral hepatitis and one with bacterial organisms. Two months later, complete recovery of the normal liver architecture is observed microscopically in the former group of animals, while fibrous scars from resolving abscesses are seen microscopically in the latter group of animal livers. Which of the following factors best explains the different outcomes for these two test groups?

A. Extent of damage to the bile ducts
B. Injury to the connective tissue framework
C. Extent of liver cell injury
D. Location of the lesion
E. Nature of the etiologic agent

23. Which of the following pathologic findings would most likely be seen in a left ventricle, which had undergone a severe myocardial infarction 1 month earlier?

A. Complete resolution
B. Nodular regeneration
C. Coagulative necrosis

D. Fibrous scar
E. Abscess

24. Which of the following steps in the inflammatory-repair response is most likely affected by neutralization of transforming growth factor-β?
A. Production of collagen
B. Migration of epithelial cells
C. Leukocyte extravasation
D. Chemotaxis of lymphocytes
E. Increase in vascular permeability

25. Liquefaction associated with necrosis occurs after infarction of the:
A. heart
B. kidney
C. brain
D. liver
E. spleen

26. By pathologic definition，gangrenous necrosis occur together with:
A. desiccation (脱水)
B. colliguative necrosis
C. involvement of a limb
D. infection of the tissues with Gram-positive organisms
E. putrefaction

27. Amyloid reacts with the following stains:
A. Gram staining
B. fluoroscein isothiocyanate
C. Prussian staining
D. methyl green
E. Congo red

28. The essential constituents of primary amyloid include:
A. immunoglobulin
B. complement
C. albumin
D. starch
E. globulin

29. Which of the following materials can be found microscopically in a atheromatous plaque specimen, in the form of crystals that produce long, cleft like spaces?
A. Immunoglobulin
B. Lipofuscin
C. Cholesterol
D. Glycogen
E. Hemosiderin

30. Amyloid is deposited infrequently in:
A. liver
B. brain
C. spleen
D. lungs
E. kidneys

31. The site in which metastatic calcification occurs is:
A. the kidney
B. the wall of the inferior vena cava
C. old tuberculous lesions
D. atheroma
E. thyroid papillary carcinoma

32. Bence-Jones proteins are:
A. the heavy chains of immunoglobulins
B. found in the urine in multiple

myeloma

C. associated with T cell lymphoma

D. found in the urine in Waldenstrom's macroglobulinemia

E. precipitated by boiling

33. Metaplasia, the transformation of one completely differentiated tissue into another, does not occur in:

A. connective tissue elements

B. the gastrointestinal tract

C. the central nervous system

D. the biliary system

E. the urothelium

34. Starvation is associated with a reduction in size of the organs or tissues with the exception of:

A. fat depots

B. heart

C. skeletal muscle

D. liver

E. bones

35. Old age is specifically associated with atrophic changes in the following tissues except for:

A. bone

B. kidneys

C. bone marrow

D. brain

E. peripheral blood

36. Hypertrophy is associated with

A. An increase in the number of visible mitoses

B. an increase in the bulk of a given tissue

C. an increase in the number of cells in an organ or tissue

D. an absolute decrease in interstitial tissue

E. an decrease in functional capacity

37. Disuse atrophy follows

A. Breakage of the duct of an exocrine gland

B. immobilization of a joint

C. interference with the nerve supply to the muscles controlling joint movement

D. interference with the blood supply

E. the diminished secretion of trophic hormones

38. Secondary amyloidosis occurs in the following conditions

A. Familial Mediterranean fever

B. thalassaemia

C. sickle-cell disease

D. multiple myeloma

E. rheumatoid arthritis

39. A 59-year-old man presents with increasing fatigue, swelling, and bone pain. Laboratory evaluation finds increased serum protein and calcium along with markedly increased amounts of protein in his urine. Subsequently he develops signs of progressive renal failure. Microscopic examination of a renal biopsy reveals deposits of eosinophilic, Congo red-positive mate-

rial in the glomeruli. When viewed under polarized light, this material displays an apple-green birefringence. What is the most likely composition of this material?

A. Light chains
B. Transthyretin
C. Beta-2-protein
D. Beta-2-microglobulin
E. Procalcitonin

40. At autopsy of an old woman WHO died from congestive heart failure, the heart weighed 620 g and showed marked left ventricular hypertrophy and severe coronary arterial atherosclerosis. Which of the following pathologic processes best accounts for the appearance of the aortic valve seen in the figure?

A. Fatty change
B. Amyloidosis
C. Lipofuscin deposition
D. Hemosiderosis
E. Dystrophic calcification



41. Histologic sections of lung tissue from a 68-year-old woman with congestive heart failure and progressive breathing problems reveal numerous hemosiderin-laden cells within the alveoli. Which of the following is the cell of origin of these “heart failure cells”?

A. Endothelial cells
B. Eosinophils
C. Lymphocytes
D. Macrophages
E. Pneumocytes

42. Which one of these statements best describes the mechanism through which Fas (CD95) initiates apoptosis?

A. BCL2 product blocks bax channels
B. Cytochrome c activates Apaf-l
C. FADD stimulates caspase 8
D. TNF inhibits IKB
E. TRADD stimulates FADD

43. Pyogenic bacterial infections and ischemic brain infarcts most characteristically produce which one of these types of necrosis?

A. Caseous necrosis
B. Coagulative necrosis
C. Fat necrosis
D. Fibrinoid necrosis
E. Liquefactive necrosis

44. A 30-year-old woman presents with malaise and increasing fatigue. hysical examination reveals a yellowtinge to her skin and sclera, and laboratory evaluation shows elevated serum liver enzmes. After further workup, a diagnosis of viral hepatitis is made. This workup included

a liver biopsy, which revealed scattered, eosinophilic cells (Councilman bodies), but very little inflammation was present. These Councilman bodies were formed by an active process that characteristically has no inflammatory response and results from the activation of genes that form new enzymes, such as endonucleases. These enzymes subsequently destroy the cell itself. Which of the following terms best describes this process?

A. Apoptosis
B. Autophagy
C. Heterophagy
D. Metaplasia
E. Necrosis

45. A 38-year-old woman presents with intermittent pelvic pain. Physical examination reveals a 3cm mass in the area of her right ovary. Histologic sections from this ovarian mass reveal a papillary tumor with multiple, scattered small, round, laminated calcifications. Which of the following is the basic defect producing these abnormal structures?

A. Bacterial infection
B. Dystrophic calcification
C. Enzymatic necrosis
D. Metastatic calcification
E. Viral infection

46. A 48-year-old man who has a long history of excessive drinking presents with signs of alcoholic hepatitis. Microscopic examination of a biopsy of this patient's liver reveals irregular eosinophilic hyaline inclusions within the cytoplasm of the hepatocytes. These eosinophilic inclusions are composed of which one of the following substances?

A. Immunoglobulin
B. Excess plasma proteins
C. Prekeratin intermediate filaments
D. Basement membrane material
E. Lipofuscin

47. An autopsy is performed on a 64-year-old man who died of congestive heart failure. Sections of the liver reveal yellow-brown granules in the cytoplasm of most of the hepatocytes. Which of the following stains would be most useful for demonstrating (with positive staining) that these yellow-brown cytoplasmic granules are in fact composed of hemosiderin (iron)?

A. Oil red a stain
B. Periodic acid-Schiff stain
C. Prussian blue stain
D. Sudan black B stain
E. Trichrome stain

48. A 54-year-old man develops a thrombus in his left anterior descending coronary artery. The area of myocardium supplied by this vessel is irreversibly injured. The thrombus is destroyed by the infusion of streptokinase, which is a plasminogen

activator, and the injured area is reperfused. The patient, however, develops an arrhythmia and dies. An electron microscopic (EM) picture taken of the irreversibly injured myocardium reveals the presence of large, dark, irregular amorphic densities within mitochondria. What are these abnormal structures?

A. Apoptotic bodies
B. Flocculent densities
C. Myelin figures
D. Psammoma bodies
E. Russell bodies

49. A 55-year-old male alcoholic presents with symptoms of liver disease and is found to have mildly elevated liver enzymes. A liver biopsy examined with a routine hematoxylin and eosin (H&E) stain reveals abnormal clear spaces in the cytoplasm of most of the hepatocytes. Which of the following materials is most likely forming these cytoplasmic spaces?

A. Calcium
B. Cholesterol
C. Hemosiderin
D. Lipofuscin
E. Triglyceride

50. High levels of ultraviolet radiant energy will produce abundant denatured proteins that aggregate in proteasomes. Which of the following substances is most likely to bind to these proteins and will target them for catabolism by cytosolic proteasomes?

A. Caspase
B. Calcium
C. Ubiquitin
D. Adenosine monophosphate
E. Hydrogen peroxide

51. If patient's chronic hypertension remains untreated for years, which of the following cellular alterations will most likely be found in the left ventricle?

A. Hemosiderosis
B. Hyperplasia
C. Hypertrophy
D. Metaplasia
E. Atrophy

52. Which of the following subcellular structures in heart failure cells is most important for the accumulation of hemosiderin?

A. Golgi apparatus
B. Chromosome
C. Ribosome
D. Lysosome
E. Endoplasmic reticulum

53. Which of the following cellular processes contributes most to the increased size of a woman's uterus in pregnancy?

A. Myometrial fibroblast proliferation
B. Myometrial smooth muscle hypertrophy
C. Endometrial stromal hypertrophy
D. Endometrial glandular hyperplasia
E. Vascular endothelial hyperplasia


54. A 37-year-old man had a 2.5cm mass, which was sharply circumscribed with a soft, white center, in his pulmonary wedge resection specimen. Mycobacterium tuberculosis is found afterwards with pathologic special staining. Which of the following pathologic processes has most likely occurred in this nodule?
    A. Fat necrosis
    B. Caseous necrosis
    C. Coagulative necrosis
    D. Apoptosis
    E. Fatty change
55. An experiment introduces a "knockout" gene mutation into a cell line. The frequency of shrunken cells with chromatin clumping and cytoplasmic blebbing is increased, compared with a cell line without the mutation. Overall survival of the mutant cell line is reduced. Which of the following genes is most likely to be affected by this mutation?
    A. p53
    B. FAS
    C. C-MYC
    D. BAX
    E. BCL-2
56. Laboratory findings in a middle-aged man show hypercalcemia and hypophosphatemia (血钙增高和血磷降低) and normal serum albumin. Urine microscopic analysis shows deposition of calcium salts in the renal tubular epithelium. Which of the following processes has most likely produced this change in the kidney?
    A. Cellular aging
    B. Dystrophic calcification
    C. Autophagocytosis
    D. Renal tubular atrophy
    E. Metastatic calcification
57. A pneumonectomy (肺切除术) is performed on an elderly lung cancer patient, and examination of the hilar lymph nodes reveals a uniform, dark-black cut surface. Which of the following factors is most likely to account for the appearance of the lymph nodes?
    A. Metastases
    B. Aging
    C. Liver failure
    D. Bleeding disorder
    E. Smoking
58. At autopsy, the heart of an elderly man weighs only 250g and has small right and left ventricles. The myocardium is firm, with a dark chocolate brown color throughout. The coronary arteries show minimal atherosclerotic changes. An excessive amount of which of the following substances will most likely be found in the myocardial fibers of this heart?
    A. Melanin
    B. Lipofuscin

C. Glycogen

D. Bilirubin

E. Hemosiderin

59. A patient undergoes bone marrow transplantation for leukemia and receives partially mismatched donor marrow. One month later, she has a scaling skin rash. Examination of a skin biopsy specimen reveals the cellular change shown in the figure. This change most likely results from which of the following biochemical reactions?



A. Increase in glycolysis


B. Lipid peroxidation

C. Activation of caspases

D. Activation of lipases

E. Reduction of ATP synthesis

60. The microscopic appearance of an enlarged liver with a yellow cut surface is shown in the preceding figure. Before death, the man's total serum cholesterol and triglyceride levels were normal, but he had a decreased serum albumin concentration and increased prothrombin time (凝血酶原时间). Which of the following activities or conditions most likely led to these findings?



A. Suffering tumor

B. Playing basketball

C. Drinking beer

D. Smoking cigarettes

E. Ingesting aspirin

61. A thrombus has been found in Li's left atrium, who has suffered worsening congestive heart failure for many years mitral stenosis and left atrial dilation. One month later, he experiences left flank pain (左肋腹部疼痛) and notes hematuria (血尿). Laboratory testing shows elevated serum AST (谷草转氨酶). Which of the following patterns of tissue injury is most likely to be present?

A. Gangrenous necrosis

B. Liquefactive necrosis

C. Caseous necrosis

D. Fat necrosis

E. Coagulative necrosis

62. A small portion of the left lobe of a man's liver is removed because of trauma. Several weeks later, a CT scan of the abdomen

shows that the liver has nearly regained its size prior to the injury. Which of the following processes best explains this CT scan finding?

A. Hypertrophy
B. Hyperplasia
C. Fatty change
D. Hydropic change
E. Dysplasia

63. An old man died subbenly and at autopsy multiple tissue sites were sampled for microscopic analysis. Examination of the tissues showed noncrystalline amorphous deposits of calcium salts in the intestinal mucosa, renal interstitium, and alveolar walls of lungs. Which of the following conditions is most likely to explain these findings?

A. Pulmonary emphysema
B. Generalized atherosclerosis
C. Disseminated tuberculosis
D. Chronic hepatitis
E. Chronic glomerulonephritis

64. A renal biopsy is performed in a renal failure patient, and the specimen is examined using electron microscopy. Which of the following morphologic changes most clearly suggests a diagnosis of acute tubular necrosis?

A. Mitochondrial swelling
B. Plasma membrane blebs
C. Chromatin clumping
D. Nuclear fragmentation
E. Ribosomal disaggregation

65. After thrombolytic therapy (溶血栓疗法) with tissue plasminogen activator (t-PA) (血浆纤溶酶原激活剂), the serum creatine kinase (CK，血清肌酸磷酸激酶) level increased in a patient with acute myocardial infarction. Which of the following events most likely occurred following t-PA therapy?

A. Chemical injury
B. Increased synthesis of CK
C. Reperfusion injury
D. Cellular regeneration
E. Myofiber atrophy

66. An old man has had difficulty with urination (排尿) for the past 10 years. A B ultrasound examination reveals that the prostate gland is enlarged to about twice normal size. The microscopic appearance of the prostate biopsy shows nodules of glands with intervening stroma. Which of the following pathologic processes has most likely occurred in the prostate?

A. Metaplasia
B. Dysplasia
C. Fatty change
D. Hypertrophy
E. Hyperplasia

67. An old man experienced an episode of chest pain prior to sudden death. A histologic section of left ventricular my-

ocardium taken at autopsy showed a deeply eosinophilic staining area with loss of nuclei and cross striations in myocardial fibers. There was no hemorrhage or inflammation. Which of the following conditions most likely produced these myocardial changes?

A. Antibodies directed against myocardium
B. Viral infection
C. Blunt chest trauma
D. Coronary artery thrombosis
E. Protein-deficient diet

68. Which of the following findings would most likely appear in a liver biopsy specimen from a congenital anemia patient who has required multiple transfusions of RBCs for many years?

A. Glycogen in hepatocytes
B. Bilirubin in canaliculi
C. Hemosiderin in hepatocytes
D. Amyloid in portal triads
E. Steatosis in hepatocytes

答案

中文题

1.E	2.E	3.E	4.A	5.E	6.B	7.B	8.D	9.C	10.A
11.B	12.C	13.A	14.E	15.D	16.B	17.C	18.B	19.D	20.E
21.C	22.D	23.B	24.D	25.C	26.D	27.E	28.D	29.A	30.B
31.D	32.E	33.E	34.C	35.C	36.E	37.B	38.C	39.A	40.B
41.B	42.E	43.E	44.A	45.A	46.A	47.A	48.C	49.E	50.D
51.A	52.D	53.E	54.D	55.D	56.D	57.E	58.C	59.D	60.D
61.C	62.E	63.C	64.D	65.B	66.B	67.E	68.D	69.D	70.D
71.E	72.A	73.E	74.B	75.A	76.C	77.A	78.C	79.A	80.B
81.B	82.B	83.E	84.B	85.B	86.B	87.B	88.C	89.C	90.B
91.A	92.D	93.E	94.D	95.A	96.D	97.E	98.A	99.C	100.D
101.B	102.B	103.E	104.D	105.B	106.B	107.E	108.D	109.C	110.B
111.A	112.B	113.E	114.A	115.E	116.C	117.D	118.C	119.C	120.A
121.C	122.B	123.E	124.A	125.B	126.C	127.D	128.E	129.E	130.C
131.C	132.B	133.E	134.D	135.D	136.C	137.E	138.B	139.A	140.B
141.A	142.D	143.A	144.B	145.C	146.B	147.C	148.C	149.C	150.A
151.E	152.E	153.C	154.D	155.B	156.C	157.E	158.C	159.D	160.D

英文题

1.D	2.D	3.B	4.D	5.A	6.A	7.E	8.D	9.E	10.B
11.E	12.C	13.A	14.A	15.E	16.B	17.D	18.A	19.E	20.D
21.D	22.B	23.D	24.A	25.C	26.E	27.E	28.A	29.C	30.D
31.E	32.B	33.C	34.E	35.E	36.B	37.B	38.D	39.A	40.E
41.D	42.C	43.E	44.A	45.B	46.C	47.C	48.B	49.E	50.C
51.C	52.D	53.B	54.B	55.E	56.E	57.E	58.B	59.C	60.C
61.E	62.B	63.E	64.D	65.C	66.E	67.D	68.C		

2　局部血液及体液循环障碍

中文题

1. 动脉性充血的主要病理变化是
   A. 器官体积增大
   B. 器官重量增加
   C. 小动脉和毛细血管扩张、充满红细胞
   D. 局部呈红色
   E. 局部温度升高
2. 淤血时扩张充盈的血管主要是
   A. 动脉
   B. 静脉
   C. 小动脉和毛细血管
   D. 小静脉和毛细血管
   E. 毛细血管、小动脉和小静脉
3. 不引起淤血的是
   A. 肝硬化
   B. 左心衰竭
   C. 右心衰竭
   D. 肠套叠
   E. 大隐静脉结扎
4. 不符合静脉性充血的是
   A. 局部紫绀
   B. 小动脉扩张
   C. 局部温度降低
   D. 局部组织肿胀
   E. 间质水肿
5. 男性，40 岁，慢性活动性肝炎伴肝硬化，符合该患者的症状是
   A. 颈静脉怒张
   B. 下肢大隐静脉曲张
   C. 下肢象皮肿
   D. 皮肤粘液性水肿
   E. 脐周静脉高度扩张
6. 不符合静脉性充血的描述是
   A. 可引起出血
   B. 可引起间质增生
   C. 可引起血栓形成
   D. 可引起实质增生
   E. 可引起器官硬化
7. 不符合静脉性充血的描述是
   A. 妊娠子宫可引起下肢淤血
   B. 肝硬化可引起胃肠道淤血
   C. 下肢深静脉血栓形成可引起下肢淤血
   D. 股动脉血栓形成可引起下肢淤血
   E. 右心衰竭可引起全身性淤血
8. 静脉淤血可引起
   A. 器官肥大
   B. 实质细胞增生
   C. 结缔组织增生
   D. 器官功能增强
   E. 多量炎细胞浸润
9. 心力衰竭细胞是指肺淤血时
   A. 肺泡内吞噬粉尘的巨噬细胞
   B. 肺泡内含有多量含铁血黄素的巨噬细胞
   C. 增生的肺泡上皮细胞
   D. 血管内皮细胞
   E. 肺泡腔内的巨噬细胞

10. 关于肺淤血的描述错误的是
    A. 肺泡腔内可见水肿液
    B. 肺泡腔内可见红细胞
    C. 肺泡腔内可见心力衰竭细胞
    D. 肺泡壁毛细血管扩张
    E. 肺泡壁内可见纤维素渗出
11. 不符合肺淤血的描述是
    A. 肺泡壁毛细血管扩张
    B. 肺泡内多量中性粒细胞
    C. 肺泡腔内有水肿液
    D. 可发生漏出性出血
    E. 常可见心力衰竭细胞
12. 可引起慢性肺淤血的有
    A. 肝硬化
    B. 左心衰竭
    C. 右心衰竭
    D. 肺动脉主干血栓栓塞
    E. 大叶性肺炎肉质变
13. 符合肺淤血的描述是
    A. 肺泡壁变薄
    B. 肺泡腔内纤维素渗出
    C. 肺泡腔内可见心力衰竭细胞
    D. 肺大泡形成
    E. 肺肉质变
14. 符合肺淤血的描述是
    A. 肺泡壁毛细血管扩张
    B. 肺泡内多量中性粒细胞
    C. 支气管鳞状上皮化生
    D. 肺泡上皮弥漫增生
    E. 肺小动脉玻璃样变
15. 符合肺淤血的描述是
    A. 肺泡上皮弥漫增生
    B. 肺泡内多量中性粒细胞
    C. 肺泡腔内大量水肿液
    D. 肺小动脉玻璃样变
    E. 肺肉质变
16. 不符合肺淤血的描述是
    A. 可引起肺褐色硬化
    B. 可发生肺肉质变
    C. 肺泡腔内有水肿液
    D. 可引起漏出性出血
    E. 可出现心力衰竭细胞
17. 槟榔肝是指
    A. 肝淤血
    B. 肝淤胆
    C. 肝硬化
    D. 肝脂肪变性
    E. 肝萎缩
18. 不符合肝淤血的病变是
    A. 肉眼下呈槟榔状
    B. 肝窦扩张
    C. 肝细胞萎缩、脂肪变性
    D. 肝细胞大片状坏死
    E. 结缔组织增生
19. 不符合肝淤血的描述是
    A. Kupffer 细胞增生
    B. 中央静脉扩张
    C. 肝窦扩张
    D. 肝细胞萎缩
    E. 肝细胞脂肪变性
20. 女性，60 岁，二尖瓣狭窄约 15 年，近日呼吸困难加重，咳粉红色泡沫痰。不符合患者肺脏的病理变化的是
    A. 肺泡内大量纤维素渗出

B. 肺泡腔内大量蛋白性液体
C. 肺泡腔内红细胞漏出
D. 肺泡壁毛细血管扩张
E. 肺泡腔内可见心力衰竭细胞

21. 男性，60 岁，长时间吸烟，肺原性心脏病已近 3 年的历史，近日咳嗽，气喘，心悸。检查肝脏肿大并有压痛。符合患者肝脏的病理变化的是
A. 肝细胞水肿，气球样变
B. 肝细胞大片坏死
C. 肝细胞碎片坏死
D. 肝细胞萎缩、脂肪变性
E. 肝细胞结节状增生

22. 不符合肝淤血的病理改变是
A. 肝窦扩张
B. 肝细胞萎缩
C. 肝细胞脂肪变性
D. 纤维组织增生
E. 肝内大量含铁血黄素沉积

23. 肝淤血可引起
A. 肝纤维化
B. 肝淤胆
C. 肝腺瘤
D. 肝坏死
E. 肝含铁血黄素沉积

24. 符合肝淤血的病理改变是
A. 小胆管内淤胆
B. 小叶内淋巴细胞灶状浸润
C. 肝细胞脂肪变性
D. 肝细胞结节状增生
E. 肝内大量含铁血黄素沉积

25. 关于血栓形成的描述，错误的是
A. 静脉血栓多于动脉血栓
B. 下肢血栓多于上肢血栓
C. 静脉内血栓多为红色血栓
D. 动脉内血栓多为混合性血栓
E. 毛细血管内血栓多为纤维素样血栓

26. 不符合漏出性出血的是
A. 流行性出血热引起的出血
B. 白血病引起的出血
C. 肺癌引起的咯血
D. 血友病 A 引起的出血
E. DIC 引起的出血

27. 漏出性出血多发生于
A. 毛细血管后静脉、毛细血管及毛细血管前动脉
B. 大动脉
C. 大静脉
D. 小动脉
E. 小静脉

28. 不属于破裂性出血的是
A. 慢性空洞型肺结核引起的大咯血
B. 肝硬化引起的大呕血
C. 十二指肠溃疡引起的大出血
D. 肾小球肾炎引起的血尿
E. 直肠癌引起的便血

29. 男性，50 岁，患轻型血友病 A，患者常在运动后发生关节红、肿、热、痛，多由关节内出血引起。此种出血的原因是
A. 血管破裂
B. 毛细血管内皮变性
C. 变态反应性血管炎
D. 凝血因子缺乏
E. 血小板减少

30. 不属于漏出性出血的是
A. 肺结核空洞引起的出血
B. 维生素 C 缺乏症引起的出血
C. 血小板减少性紫癜引起的出血
D. 肺淤血引起的出血
E. 流行性出血热引起的出血

31. 漏出性出血多发生于
A. 毛细血管
B. 小静脉
C. 小动脉
D. 大静脉
E. 大动脉

32. 男性，30 岁，突发性畏寒、发热；并出现颜面、颈部及上胸部潮红，皮肤瘀点；伴有头痛、腰痛及眼眶痛，并出现肾功能衰竭。确诊为流行性出血热。患者的出血性倾向主要是因为
A. 血管破裂
B. 毛细血管壁损害
C. 血小板减少
D. 血小板功能异常
E. 凝血因子缺乏

33. 不属于血栓形成的描述是
A. 冠状动脉前降支斑块处凝血块
B. 主动脉粥样斑块表面凝血块
C. 心肌梗死处的心内膜表面凝血块
D. 尸体解剖时心脏内凝血块，并与心壁粘连不紧密
E. 克山病心肌坏死处心内膜表面凝血块

34. 与血栓形成无关的因素是
A. 血管内膜损伤
B. 血流缓慢
C. 涡流形成
D. 血小板数量增多或粘性增加
E. 纤维蛋白溶酶增加

35. 不易引起血栓形成的疾病是
A. 大面积烧伤
B. 晚期胃癌
C. 冠状动脉粥样硬化
D. 心房纤颤
E. 肝硬化

36. 不符合血栓形成的描述是
A. 静脉血栓多于动脉血栓
B. 下肢血栓多于上肢
C. 动脉内血栓多为混合血栓
D. 静脉内血栓多为红色血栓
E. 毛细血管内血栓多为纤维素性血栓

37. 符合血栓的描述是
A. 血栓由红细胞构成
B. 血栓位于心血管外
C. 血栓是指死后凝血
D. 血栓不易脱落
E. 血栓常与血管壁粘连紧密，不易剥离

38. 与血栓形成无关的因素是
A. 血管内膜损伤
B. 血流缓慢
C. 血小板数量增多
D. 癌细胞崩解产物
E. 纤维蛋白溶酶增加

39. 在血栓形成过程中起核心作用的是
A. 血小板凝集
B. 血流缓慢

C. 涡流形成
D. 内皮损伤
E. 凝血酶原增多

40. 男性，60 岁，胃癌根治术后 7 天，出现左下肢肿痛。检查时，小腿腓肠肌有压痛、沿股静脉压迫有疼痛，下肢可凹性水肿。诊断为左下肢深静脉血栓形成。与深静脉血栓形成无关的因素是
A. 癌细胞释放促凝因子
B. 手术后血小板增多、粘性增高
C. 下肢血流缓慢
D. 血液中蛋白质 C、蛋白质 S 大量减少
E. 纤维蛋白溶酶系统被激活

41. 透明血栓多见于
A. 大动脉
B. 小动脉
C. 微循环小血管内
D. 小静脉
E. 大静脉

42. 属于白色血栓的是
A. 亚急性细菌性心内膜炎的疣赘物
B. 风湿性心内膜炎的疣状赘生物
C. 心肌栓死时的附壁血栓
D. 静脉内柱状血栓的体部
E. 静脉内柱状血栓的尾部

43. 不属于白色血栓的是
A. 静脉内柱状血栓的头部
B. 动脉内延续性血栓的头部
C. 风湿性心内膜炎的疣状赘生物
D. 心房纤颤时心耳内形成的球状血栓
E. 赘疣性血栓性心内膜炎的内膜赘生物

44. 混合性血栓可见于
A. 心室内附壁血栓
B. 静脉血栓头部
C. 风湿性心内膜炎的疣状血栓
D. 肾小球毛细血管内
E. 动脉血栓头部

45. 下肢深静脉内延续性血栓的尾部是
A. 透明血栓
B. 白色血栓
C. 混合性血栓
D. 红色血栓
E. 肿瘤性栓子

46. 女性，36 岁，因胆囊结石行胆囊切除术，术后 5 天，出现右小腿肿胀、凹陷性水肿，腓肠肌压痛。诊断为右下肢深静脉血栓形成。深静脉血栓是
A. 白色血栓
B. 红色血栓
C. 混合性血栓
D. 延续性血栓
E. 透明血栓

47. 心肌梗死时的附壁血栓属于
A. 透明血栓
B. 白色血栓
C. 混合性血栓
D. 红色血栓
E. 肿瘤性栓子

48. 下肢深静脉内延续性血栓的头部是
A. 透明血栓
B. 白色血栓
C. 混合性血栓

D. 红色血栓
E. 肿瘤性栓子

49. 下肢深静脉血栓形成，可引起
A. 下肢坏疽
B. 门静脉栓塞
C. 肺动脉栓塞
D. 肠系膜动脉栓塞
E. 脑动脉栓塞

50. 上肢深静脉血栓形成，可引起
A. 肠系膜动脉栓塞
B. 冠状动脉栓塞
C. 门静脉栓塞
D. 肺动脉栓塞
E. 脑动脉栓塞

51. 静脉内较大血栓完全机化需要的时间是
A. 1～2 天
B. 1 周
C. 2 周
D. 3 周
E. 4 周

52. 不符合血栓结局的描述是
A. 较小血栓可溶解吸收
B. 较大血栓可由肉芽组织机化
C. 长时间不能机化的血栓可形成静脉石
D. 机化的血栓，可使上下游血流再沟通
E. 再沟通的血管可恢复正常血流

53. 股静脉血栓形成不易发生
A. 栓塞
B. 机化
C. 脱落
D. 钙化
E. 下肢坏疽

54. 不符合血栓机化与再通的描述是
A. 血栓干燥收缩
B. 血管内皮增生
C. 裂隙形成
D. 血流通过
E. 血流恢复

55. 股静脉血栓形成患者需卧床休息的时间为
A. 5 天
B. 10 天
C. 15 天
D. 20 天
E. 25 天

56. 静脉石是指
A. 血栓溶解
B. 血栓机化
C. 血栓脱落
D. 血栓钙化
E. 血栓吸收

57. 不符合血栓对机体影响的描述是
A. 阻塞血管
B. 引起栓塞
C. 引起心瓣膜变形
D. DIC 可引起广泛出血、加重休克
E. 对机体有害无利

58. 关于侧支循环的描述，错误的是
A. 痔静脉丛扩张是门静脉与下腔静脉之间的吻合支开放扩张的结果
B. 脐周腹壁静脉扩张是门静脉与下腔

静脉之间的吻合支开放扩张的结果

C. 食管下静脉扩张是门静脉与上腔静脉之间吻合支开放扩张的结果

D. 侧支循环具有代偿作用

E. 静脉有丰富的吻合支

59. 弥漫性血管内凝血形成的血栓是

A. 透明血栓

B. 白色血栓

C. 混合性血栓

D. 红色血栓

E. 肿瘤性栓子

60. 符合左心室附壁血栓的描述是

A. 阻塞心室血流

B. 加重心肌梗死

C. 诱发心壁穿孔

D. 引起脑栓塞

E. 引起肺栓塞

61. 弥散性血管内凝血（DIC）发生广泛出血的主要原因是

A. 肝凝血酶原合成减少

B. 血管壁广泛损伤

C. 大量血小板及纤维蛋白原消耗

D. 单核巨噬细胞系统功能下降

E. 血浆中缓激肽浓度增高

62. 微血栓的主要成分是

A. 红细胞

B. 中性粒细胞

C. 纤维素

D. 血小板

E. 淋巴细胞

63. 微循环纤维素性血栓又叫

A. 透明血栓

B. 白色血栓

C. 红色血栓

D. 混合血栓

E. 层状血栓

64. 心瓣膜的血栓一般不发生

A. 瓣膜粘连

B. 瓣膜穿孔

C. 血栓脱落

D. 瓣膜变形

E. 完全溶解吸收

65. 最常见的栓子是

A. 血栓栓子

B. 空气栓子

C. 细菌栓子

D. 肿瘤栓子

E. 羊水栓子

66. 大循环的动脉栓塞，栓子主要来自

A. 左心

B. 主动脉瓣

C. 主动脉粥样硬化溃疡

D. 主动脉瘤

E. 先天性心脏病的反常栓塞

67. 栓子是指

A. 活体心血管内血液凝固的过程

B. 人体死亡后，血液凝固的过程

C. 循环血液中不溶于血液，随血流运动并阻塞血管的物质

D. 心瓣膜的疣状赘生物

E. 动脉夹层内的凝血块

68. 右上肢静脉血栓脱落主要栓塞于

A. 肠系膜动脉

B. 脑动脉

C. 肺动脉
D. 肾动脉
E. 心冠状动脉

69. 循环血液中的凝血块随血流运行至相应大小的血管引起管腔阻塞的过程叫做
A. 血栓
B. 血栓形成
C. 血栓栓塞
D. 梗死
E. 血栓栓子

70. 脑动脉发生栓塞，其栓子最可内来自
A. 下肢深静脉血栓
B. 下肢浅静脉血栓
C. 盆腔静脉血栓
D. 左心室附壁血栓
E. 门静脉血栓

71. 交叉性栓塞不发生于
A. 室间隔缺损
B. 房间隔缺损
C. 法乐四联症
D. 动脉导管未闭
E. 主动脉狭窄

72. 交叉性栓塞可发生于
A. 室间隔缺损
B. 股动脉血栓形成
C. 门静脉血栓形成
D. 主动脉瘤
E. 主动脉狭窄

73. 引起肺动脉栓塞的血栓栓子主要来自
A. 二尖瓣
B. 主动脉瓣
C. 三尖瓣
D. 下肢深静脉
E. 肠系膜下静脉

74. 引起患者猝死的肺动脉血栓栓塞是
A. 较小栓子栓塞于肺动脉小分支
B. 较小栓子栓塞于淤血肺的动脉小分支
C. 巨大的血栓栓子栓塞于肺动脉主干
D. 较小栓子反复栓塞于肺动脉小分支
E. 较小栓子反复栓塞于淤血肺的动脉小分支

75. 栓塞最常见的类型为
A. 血栓栓塞
B. 脂肪栓塞
C. 羊水栓塞
D. 空气栓塞
E. 肿瘤细胞栓塞

76. 女性，36 岁，因胆囊结石住院手术，手术后 3 天，患者出现右侧胸痛，伴有咳嗽、气喘。查体：右侧胸膜可闻摩檫音，左小腿肿胀，凹陷性水肿。腓肠肌压痛，X 线检查，右下肺有阴影。临床诊断应为
A. 肺淤血
B. 右肺小叶性肺炎
C. 左下肢深静脉炎
D. 右肺血栓栓塞
E. 左下肢深静脉血栓形成，肺动脉血栓栓塞伴肺梗死

77. 肺动脉栓塞多为
A. 血栓栓塞
B. 空气栓塞

C. 羊水栓塞
D. 细菌栓子栓塞
E. 脂肪栓塞

78. 体循环动脉栓子主要来自
A. 右心
B. 左心
C. 主动脉
D. 大动脉
E. 大静脉

79. 不易引起栓塞的是
A. 风湿性心内膜炎
B. 急性细菌性心内膜炎
C. 亚急性细菌性心内膜炎
D. 心肌梗死伴附壁血栓形成
E. 霉菌性心内膜炎

80. 栓塞不易引起组织梗死的器官是
A. 肾脏
B. 脾
C. 脑
D. 大肠
E. 肝脏

81. 关于脂肪栓塞的描述，正确的是
A. 脂肪栓子全部来源于损伤的脂肪组织
B. 直径小于 20um 的脂滴可栓塞于肺毛细血管
C. 进入肺内脂肪量达 9～20g 时，肺内血循环面积可丧失 3/4，导致窒息和右心衰竭
D. 脑脂肪栓塞，可引起脑点状出血和坏死，是死亡主要原因
E. 肾脂肪栓塞可引起急性肾功能衰竭，是死亡的常见原因

82. 脂肪栓塞患者死亡的常见原因是
A. 急性右心衰竭
B. 急性左心衰竭
C. 脑出血坏死
D. 肾功能衰竭
E. 心源性休克

83. 脂肪栓塞患者一般的死亡原因是
A. 动脉系统栓塞
B. 脂肪分解产生自身中毒
C. 肾小动脉栓塞
D. 肺水肿和心功能不全
E. 脑小动脉栓塞

84. 男性，40 岁，因左股骨粉碎性骨折入院，突然发生昏厥、呼吸困难、胸痛。查体，呼吸增快、紫绀，肺部湿　音。X 线检查，肺不张，胸腔内可见少量积液。肺内的病变是
A. 肺炎
B. 肺动脉血栓栓塞
C. 肺脂肪栓塞
D. 肺气肿
E. 肺不张

85. 股骨粉碎性骨折可以引起
A. 血栓栓塞
B. 空气栓塞
C. 羊水栓塞
D. 细菌栓子栓塞
E. 脂肪栓塞

86. 脑脂肪栓塞可引起
A. 脑梗死
B. 脑水肿

C. 脑大片出血
D. 脑积水
E. 脑萎缩

87. 肺脂肪栓塞不会引起
A. 肺水肿
B. 肺出血
C. 肺不张
D. 肺气肿
E. 右心衰竭

88. 与脂肪栓塞无关的疾病有
A. 粉碎性骨折
B. 脂肪组织挫伤
C. 糖尿病
D. 脂肪肝
E. 甲状腺机能亢进

89. 可引起空气栓塞的是
A. 静脉穿刺输液
B. 锁骨下静脉插管输液
C. 肺气肿伴发自发性气胸
D. 支气管扩张症伴发脓气胸
E. 慢性空洞型肺结核病

90. 不会引起空气栓塞的是
A. 胎盘早期剥离
B. 胸部外伤损伤深静脉
C. 颈部外伤损伤深静脉
D. 锁骨下静脉插管输液
E. 静脉穿刺输液

91. 减压病患者血液中形成的气泡的气体是
A. 氧
B. 二氧化碳
C. 氮
D. 氢
E. 氖

92. 一次多量空气进入静脉可引起猝死，最低进入空气量约为
A. 1ml
B. 5ml
C. 10ml
D. 50ml
E. 100ml

93. 不会发生气体栓塞的有
A. 颈部外伤或手术
B. 胸部外伤或手术
C. 大隐静脉切开输液
D. 胎盘早期剥离
E. 锁骨下静脉插管输液

94. 引起沉箱病的栓塞为
A. 血栓栓塞
B. 气体栓塞
C. 羊水栓塞
D. 脂肪栓塞
E. 寄生虫虫卵栓塞

95. 胸壁及肺创伤损伤深部静脉可以引起
A. 血栓栓塞
B. 空气栓塞
C. 羊水栓塞
D. 细菌栓子栓塞
E. 脂肪栓塞

96. 可以发生气体栓塞的有
A. 手臂静脉穿刺输液
B. 足背静脉穿刺输液
C. 大隐静脉切开输液
D. 小儿头皮静脉穿刺输液

E. 锁骨下静脉插管输液

97. 血吸虫卵主要栓塞于

A. 肺毛细血管

B. 肠壁小静脉

C. 肝内门静脉分支

D. 脑小血管内

E. 脾窦内

98. 羊水栓塞的主要病理诊断依据是

A. 肺毛细血管扩张

B. 肺泡内出血

C. 肺内透明膜形成

D. 肺泡腔内角化上皮、胎便小体

E. 肺毛细血管内角化上皮及其他羊水成分

99. 女性，52 岁，书店经理。因近四周来疲乏无力、嗜睡、体重下降、间歇性发热出汗而入院。查体，左结膜下出血，两下肢多处出血斑。两手甲下可见线状出血，脾脏可触及，心率缓慢，心前区可闻收缩期吹风样杂音。临床诊断为急性细菌性心内膜炎。患者多脏器可发生栓塞，其栓子为

A. 血栓栓子

B. 脂肪栓子

C. 细菌栓子

D. 肿瘤栓子

E. 空气栓塞

100. 男性，25 岁，静脉吸毒者。因突发高热、寒战、皮肤淤点及左前臂注射部位红肿、热、痛二天而住院。查体，患者体弱、消瘦，左前臂红肿，压痛，针眼处可见脓疱，皮肤多处淤点。肝脾大，心前区可闻高调收缩期杂音。血液检查，白细胞总数 25×10^9/L。以中性粒为主。临床诊断为急性感染性心内膜炎。患者二尖瓣疣状赘生物脱落的栓子引起感染为

A. 肺脏感染

B. 肝脏感染

C. 下肢感染

D. 肾脏感染

E. 心、脑、肝、肾、肺等全身多脏器感染

101. 急性细菌性心内膜炎引起的梗死的特点是

A. 梗死灶呈地图状

B. 梗死灶呈锥体形

C. 梗死灶呈节段状

D. 梗死灶发生液化

E. 梗死灶化脓

102. 肝癌肺转移时肺小血管内的瘤细胞团是

A. 透明血栓

B. 白色血栓

C. 混合性血栓

D. 红色血栓

E. 肿瘤性栓子

103. 羊水早破可以引起

A. 血栓栓塞

B. 空气栓塞

C. 羊水栓塞

D. 细菌栓子栓塞

E. 脂肪栓塞

104. 急性细菌性心内膜炎可以引起

A. 血栓栓塞

B. 空气栓塞
C. 羊水栓塞
D. 细菌栓子栓塞
E. 脂肪栓塞

105. 梗死是指
A. 缺血引起的坏死
B. 生理性程序性死亡
C. 厌氧细菌感染而引起的坏死
D. 结核杆菌感染而引起的坏死
E. 胰酶外溢而引起的组织坏死

106. 不会引起梗死的是
A. 冠状动脉主干阻塞
B. 卵巢囊肿蒂扭转
C. 尺动脉断裂
D. 肠套叠
E. 嵌顿性疝

107. 不符合梗死的描述是
A. 动脉痉挛促进梗死形成
B. 梗死多由动脉阻塞引起
C. 有效的侧支循环的建立可防止梗死发生
D. 左肝动脉阻塞可引起肝左叶梗死
E. 双重血液循环的器官不易发生梗死

108. 梗死最常见的原因是
A. 血栓形成
B. 血栓栓塞
C. 血管受压
D. 动脉痉挛
E. 动脉壁肿瘤

109. 不符合梗死的描述是
A. 双重血液循环的器官不易发生梗死
B. 全身血液循环状态对梗死的形成无影响
C. 动脉痉挛促进梗死的形成
D. 有效的侧支循环的建立可防止梗死发生
E. 梗死多由动脉阻塞引起

110. 一般不引起肠梗死的疾病是
A. 肠套叠
B. 肠扭转
C. 嵌顿疝
D. 肠系膜上动脉栓塞
E. 门脉高压

111. 男性，65 岁，长期吸烟，近 3 年出现心绞痛，近半年来，右下肢间歇性疼痛，检查右下肢足背动脉波动弱，右足苍白，发凉。右足变化符合
A. 充血
B. 淤血
C. 缺血
D. 水肿
E. 出血

112. 脑梗死的常见原因是
A. 动脉痉挛
B. 动脉受压
C. 血栓栓塞
D. 血栓形成
E. 动脉瘤形成

113. 容易发生贫血性梗死的脏器是
A. 小肠
B. 大肠
C. 肺
D. 肾
E. 卵巢

114. 不符合贫血性梗死的描述是
A. 发生于组织结构致密的器官
B. 梗死灶呈灰白色
C. 周围可见充血出血带
D. 梗死区正常结构破坏
E. 梗死灶呈锥形

115. 梗死为液化性坏死的器官是
A. 心
B. 肺
C. 脑
D. 肾
E. 肠

116. 动脉阻塞后极少发生梗死的器官为
A. 肾
B. 脾
C. 心
D. 肝
E. 脑

117. 男性，50 岁，警官，近一年来经常出现间歇性胸痛，劳累后出现症状，疼痛向左肩、左臂放射。有近 20 年吸烟史。此次发作持续时间超过四小时，伴有恶心、呕吐，并出现心率失常。心电图显示 T 波倒置。心肌的病变是
A. 心肌梗死
B. 心肌细胞增生
C. 心肌蜡样变性
D. 心肌肥大
E. 心肌萎缩

118. 女性，57 岁，饭馆经理，因帮助服务员抬啤酒箱，突然发作，前胸部剧痛并向左肩放射，伴有恶心、呕吐，发作后半小时入院。查体，患者心率缓慢，皮肤苍白而潮湿，心电图显 ST 段升高。X 线检查，胸部未见异常。诊断为心肌梗死。心肌梗死的常见原因是
A. 冠状动脉血栓栓塞
B. 冠状动脉粥样硬化合并血栓形成
C. 主动脉瓣狭窄
D. 梅毒性心脏病
E. 冠状动脉炎症性疾病

119. 脑梗死的特点是
A. 梗死灶呈地图状
B. 梗死灶呈锥体形
C. 梗死灶呈节段状
D. 梗死灶发生液化
E. 梗死灶化脓

120. 心肌梗死的特点是
A. 梗死灶呈地图状
B. 梗死灶呈锥体形
C. 梗死灶呈节段状
D. 梗死灶发生液化
E. 梗死灶化脓

121. 容易发生出血性梗死的脏器是
A. 脑
B. 肺
C. 心
D. 肾
E. 脾

122. 肺出血性梗死常发生于
A. 肺淤血
B. 肺水肿
C. 肺纤维化
D. 肺气肿

E. 肺不张

123. 肠出血性坏死可发生于

A. 肠淤血

B. 肠水肿

C. 肠纤维化

D. 肠系膜动脉、静脉均受压

E. 肠壁增厚

124. 肺梗死的特点是

A. 梗死灶呈地图状

B. 梗死灶呈锥体形

C. 梗死灶呈节段状

D. 梗死灶发生液化

E. 梗死灶化脓

125. 肠梗死的特点是

A. 梗死灶呈地图状

B. 梗死灶呈锥体形

C. 梗死灶呈节段状

D. 梗死灶发生液化

E. 梗死灶化脓

126. 出血性梗死的特点是

A. 梗死灶呈地图状

B. 梗死灶呈锥体形

C. 梗死灶呈节段状

D. 梗死灶呈暗红色

E. 梗死灶化脓

127. 败血性梗死的特点是

A. 梗死灶呈地图状

B. 梗死灶呈锥体形

C. 梗死灶呈节段状

D. 梗死灶发生液化

E. 梗死灶化脓

128. 败血性梗死常见于

A. 急性细菌性心内膜炎

B. 无菌性心内膜炎

C. 心肌梗死伴附壁血栓形成

D. 克山病伴附壁血栓形成

E. 风湿性心内膜炎

129. 梗死不会引起严重后果的器官是

A. 心

B. 脑

C. 小肠

D. 肺

E. 脾

130. 肾梗死一般不会引起

A. 肾功能衰竭

B. 血尿

C. 腰痛

D. 蛋白尿

E. 管形尿

131. 不符合脑梗死的描述是

A. 可引起偏瘫

B. 可引起死亡

C. 多由脑动脉瘤引起

D. 梗死灶发生液化

E. 小梗死灶可以吸收

132. 不符合心肌梗死的描述是

A. 梗死灶呈地图状

B. 可引起休克

C. 可引起死亡

D. 梗死灶发生液化

E. 可引起心脏破裂

133. 脾梗死灶不会发生

A. 溶解吸收

B. 机化

C. 包裹
D. 钙化
E. 囊肿

134. 梗死对机体的影响取决于
A. 受累的器官
B. 梗死灶的大小
C. 梗死灶的形状
D. 梗死灶的继发变化
E. 受累的器官和梗死灶的大小

135. 肺梗死不会引起
A. 胸痛
B. 咯血
C. 呼吸困难
D. 梗死灶发生液化
E. 胸膜炎

136. 心肌梗死灶的修复方式为
A. 溶解吸收
B. 瘢痕修复
C. 心肌细胞增生修复
D. 钙化
E. 包裹

137. 丝虫病患者下肢出现象皮肿的主要原因是
A. 右心衰竭
B. 血浆胶体渗透压降低
C. 毛细血管内液体静力压升高
D. 毛细血管通透性升高
E. 淋巴回流受阻

138. 急性肾小球肾炎时常引起
A. 眼睑水肿
B. 下肢象皮肿
C. 肺水肿
D. 腹水
E. 皮肤黏液性水肿

139. 炎症水肿的主要原因是
A. 血浆胶体渗透压降低
B. 静脉压升高
C. 毛细血管通透性增高
D. 淋巴管闭塞
E. 肝细胞对醛骨酮、抗利尿素的灭活能力降低

140. 乳腺癌根治术后上肢水肿的主要原因是
A. 血浆胶体渗透压低下
B. 静脉压上升
C. 毛细血管通透性增高
D. 淋巴管闭塞
E. 肝细胞对醛骨酮、抗利尿激素灭活能力降低

141. 急性左心衰竭时常引起
A. 皮肤粘液性水肿
B. 眼睑水肿
C. 下垂性水肿
D. 下肢象皮肿
E. 肺水肿

142. 脑水肿可引起
A. 颅内压增高
B. 脑沟变深
C. 脑回变窄
D. 血管周围间隙变窄
E. 细胞外间隙变窄

143. 肺水肿的主要表现是
A. 肺泡腔内纤维素渗出
B. 肺泡腔内中性粒细胞渗出

C. 肺泡腔内充满水肿液
D. 肺泡腔内透明膜形成
E. 肺泡上皮弥漫增生

144. 皮肤水肿的主要表现是
A. 表皮增厚
B. 真皮结缔组织疏松
C. 毛细血管扩张
D. 红细胞漏出
E. 皮脂腺增多

145. 符合血液循环障碍的描述是
A. 梗死是缺血引起的一种坏死
B. 血管内血液凝固形成血栓
C. 肺梗死属贫血性梗死
D. 梗死灶的形状与血管分布无关
E. 脂肪栓塞最常见

146. 不符合血液循环障碍的描述是
A. 淤血可以引起器官功能增强
B. 血栓形成可以引起梗死
C. 栓塞可以引起患者死亡
D. 梗死可以引起患者死亡
E. 出血可以引起黑便

147. 与淤血无关的是
A. 血栓形成
B. 器官纤维化
C. 出血性梗死
D. 贫血性梗死
E. 水肿

148. 发生出血性梗死的器官是
A. 脾
B. 心脏
C. 脑
D. 卵巢
E. 肝

149. 梗死灶呈节段性的器官是
A. 小肠
B. 心脏
C. 脑
D. 卵巢
E. 肝

150. 梗死灶呈地图状的器官是
A. 小肠
B. 心脏
C. 脑
D. 卵巢
E. 肝

151. 不会引起淤血的是
A. 早期妊娠
B. 肝硬化
C. 血栓形成
D. 心力衰竭
E. 静脉曲张

152. 结局中不伴有肉芽组织增生的病变有
A. 附壁血栓
B. 透明血栓
C. 柱状血栓
D. 出血性梗死
E. 贫血性梗死

153. congestion
A. 出血
B. 充血
C. 淤血
D. 缺血
E. 贫血

154. hemorrhage

A. 出血
B. 充血
C. 淤血
D. 缺血
E. 贫血

155. thrombus
A. 出血
B. 血栓
C. 淤血
D. 缺血
E. 贫血

156. embolism
A. 栓塞
B. 栓子
C. 血栓
D. 梗死
E. 水肿

157. infarct
A. 栓塞
B. 栓子
C. 血栓
D. 梗死
E. 水肿

158. edema
A. 栓塞
B. 栓子
C. 血栓
D. 梗死
E. 水肿

159. phleblith
A. 栓塞
B. 栓子
C. 血栓
D. 静脉石
E. 再通

160. heart failure cell
A. 心肌细胞
B. 心力衰竭细胞
C. 异物巨细胞
D. 单核细胞
E. 中性粒细胞

英文题

1. All of the followings are adverse effects of thrombosis except____
A. Obstruction of arteries and veins
B. Deformity of mitral valve
C. DIC and extensive hemorrhage
D. Wound hemostasis
E. Embolism

2. Which of the followings does not represent a potential outcome of thrombus?
A. Dissolution
B. Recanalization
C. Organization
D. Infarction
E. Embolization

3. Which of the following factors is not associated with thrombosis?
A. Thrombocythemia
B. Vasculitis

C. Fibrin degradation products
D. Atrial fibrillation
E. Prolonged bed rest

4. A 9-year-old boy suddenly develops severe testicular pain. He is taken to the emergency room, where he is evaluated and immediately taken to surgery. There his left testis is found to be markedly hemorrhagic due to testicular torsion. Which of the following mechanisms is primarily involved in producing this type of testicular infarction?
A. Arterial occlusion
B. Septic implantation
C. Decreased collateral blood flow
D. Increased dual blood flow
E. Venous occlusion

5. Which is liquefactive necrosis:
A. Pulmonary infarct
B. Spleenic infarct
C. Intestinal infarct
D. Cerebral infarct
E. Myocardial infarct

6. Pulmonary infarct is always:
A. Wedge shaped
B. Anemic infarction
C. Liquefactive necrosis
D. In the upper lobe
E. Caused by occlusion of bronchial arteria

7. Red/hemorrhagic infarction always occur in:
A. Heart and lung
B. Small intestine and lung
C. Lung and kidney
D. Kidney and heart
E. Spleen and liver

8. A majority of all infarction result from:
A. Local vasospasm
B. Extrinsic compression of a vessel such as by tumor
C. Thrombus or embolism in arteri
D. Swelling of an atheroma secondary to hemorrhage within a plaque
E. Venous thrombosis

9. The commonest type of embolism is :
A. Thromboembolism
B. Fat embolism
C. Air embolism
D. Amniotic fluid embolism
E. Tumor embolism

10. Which organ is the most resistant to infarction if a vessel becomes occluded?
A. Lung
B. Kidney
C. Brain
D. Heart
E. Spleen

11. A 27-year-old man is on a scuba diving trip to the Caribbean and descends to a depth of 50 m in the Blue Hole off the coast of Belize. After 30 minutes, he has a malfunction in his equipment and quickly returns to the boat on the surface.

He soon experiences difficulty breathing, with dyspnea and substernal chest pain, followed by a severe headache and vertigo. About 1 hour later, he develops severe, painful myalgias and arthralgias. These symptoms abate within 24 hours. Which of the following mechanisms is the most likely cause of these symptoms?

A. Disseminated intravascular coagulation
B. Systemic vasodilation
C. Venous thrombosis
D. Tissue nitrogen emboli
E. Fat globules in arterioles

12. A 62-year-old retired man was found to have high blood pressure of (172/100mm Hg) 2 months ago. Sincethen, he hasn't taken any medications. About 5 hours ago, he was admitted to the hospital after suddenly losing consciousness. When he regained consiousness, he was unable to move his right arm and his right leg. Which of the following intracranial pathologic abnormalities is most likely to be present?

A. Metastatic carcinoma
B. Hemorrhage into the putamen
C. Cerebral venous thrombosis
D. Subfrontal meningioma
E. Acute pyogenic meningitis

13. A 49-year-old man is in stable condition after an infarction of the anterior left ventricular wall. He receives therapy with antiarrhythmic and pressor agents. Three days later, he develops severe breathlessness, and an echocardiogram shows a markedly decreased ejection fraction. He dies 2 hours later. At autopsy, which of the following microscopic changes is most likely to be present in the lungs?

A. Congestion of alveolar capillaries with fibrin and neutrophils in alveoli
B. Congestion of alveolar capillaries with transudate in alveoli
C. Fibrosis of alveolar walls with hemosiderin-laden macrophages in alveoli
D. Multiple areas of subpleural hemorrhagic necrosis
E. Purulent exudate in the pleural space

14. A 21-year-old woman sustains multiple injuries, including fractures of the right femur and tibia and the left humerus, in a motor vehicle collision. She is admitted to the hospital, and the fractures are stabilized surgically. In a short time following surgical treatment she is in stable condition. However, 2 days later, she suddenly becomes severely dyspneic. Which of the following complications is the most likely cause of this sudden respiratory problem??

A. Right hemothorax
B. Pulmonary edema

C. Fat embolism
D. Cardiac tamponade
E. Pulmonary infarction

15. Which is the main reason for sudden death caused by pulmonary emboli:
A. Left ventricular failure
B. Right ventricular failure
C. Pulmonary infarction
D. Renal failure
E. DIC

16. The most important characteristic of amniotic fluid embolism is:
A. Pulmonary edema
B. Diffuse alveolar damage
C. Fibrin thrombi in the pulmonary microcirculation
D. The presence of squamous cell shedding lanugo hair, fat and mucin in pulmonary microcirculation
E. The presence of squamous cell sheding lanugo hair, fat and mucin in alveolar spaces

17. Paradoxical/crossed embolism may not be found in patient with:
A. Interatrial septum defect
B. Interventricular septum defect
C. Aortic stenosis
D. Patent Ductus Arteriosus
E. Tetralogy of Fallot

18. Paradoxical/crossed embolism may be found in patients with:
A. Interatrial septum defect
B. Mitral stenosis
C. Aortic aneurysm
D. Pulmonary embolism
E. Atherosclerosis


19. The gas emboli of decompression sickness are composed of:
A. Oxygen
B. Nitrogen
C. Carbon dioxide
D. Chlorine
E. Hydrogen

20. Fat emboli are most likely be found associated with
A. Obstetric procedures
B. Chest wall injury
C. Fractures of long bones
D. Intravenous injection
E. Blood transfusion

21. Phlebothrombosis most likely affects the veins in what area?
A. The upper extremities
B. The lower extremities
C. Ovarian and periuterine veins
D. Portal vein
E. Periprostatic plexus

22. The arterial thromboemoboli of lower limbs most likely arise from:
A. Left ventricular and atria
B. Right ventricular and atria
C. Portal vein
D. Superior mesenteric vein
E. Renal vein

23. A 65-year-old man had rheumatic heart disease 20 years ago. He presents to the emergency room complaining of breathlessness for the past three days. He woke at 3 am feeling short of breath and had to sleep in his recliner for the rest of night. On physical examination, he has swelling in his feet and ankles, distended neck veins with visible cannon waves and his liver is palpable three centimeters below the right costal margin. Based on the gross appearance of the liver seen in the figure, which of the following underling conditions is most likely to be present?



A. Inferior vena cava obstruction
B. Chronic renal failure
C. Pulmonary infection
D. Congestive heart failure
E. Cirrhotic portal hypertension

24. Thrombi of large deep veins of arms may cause emboli in:
A. Coronary artery
B. Mesenteric artery
C. Pulmonary artery
D. Arteria cerebri media
E. Portal vein

25. The thromboemoboli of arteria cerebri media may most likely arise from:
A. Renal vein
B. Femoral artery
C. Portal vein
D. Superior mesenteric vein
E. Left ventricular

26. Most pulmonary thromboemoboli arise from thrombi within:
A. Large deep veins of the lower legs
B. Superficial leg veins
C. Portal vein
D. Superior mesenteric vein
E. Renal vein

27. During the autopsy of a 46-year-old man who died when the motorcycle he was riding was hit by a truck, a 1.2cm red mass is found within a branch of the left pulmonary artery. Grossly this mass is rubbery, gelatinous, and has a "chicken fat" appearance. Histologic sections reveal that this mass is not attached to the wall of the pulmonary artery, and lines of Zahn are not seen. Which of the following statements best describes this intravascular mass?
A. Postmortem blood clot
B. Postmortem hematoma
C. Premortem embolic blood clot
D. Premortem non-embolic thrombus
E. Premortem non-thrombotic embolus

28. A 59-year-old obese woman with a history

of diabetes mellitus had a myocardial infarction 3 months ago. She is now taking low dose aspirin to reduce the risk of arterial thrombosis. On which of the following steps in hemostasis does aspirin has its greatest effect?

A. Adhesion of platelets to collagen
B. Aggregation of platelets
C. Production of tissue factor
D. Synthesis of von Willebrand factor
E. Synthesis of antithrombin III

29. A 73-year-old man was diagnosed 1 year ago with pancreatic adenocarcinoma. He now sees his physician because of a transient ischemic attack. On auscultation of the chest, a heart murmur is heard. Echocardiography shows an 1cm nodular lesion on the superior aspect of the anterior mitral valve leaflet. The valve leaflet appears to be intact. The blood culture is negative. Which of the following terms best describes this mitral valve lesion?

A. Adenocarcinoma
B. Atheroma
C. Mural thrombus
D. Petechial hemorrhage
E. Vegetation

30. A 56-year-old man with a history of diabetes mellitus goes to the emergency department because he has had left-sided chest pain that radiates to the arm for the past 5 hours. Serial measurements of serum creatine kinase-MB levels show elevation 24 hours after the onset of pain. Partial thromboplastin time and prothrombin time are normal. Coronary angiography shows occlusion of the left anterior descending artery. Which of the following mechanisms is the most likely cause of thrombosis in this patient?

A. Stasis of blood flow
B. Damage to endothelium
C. Decreased production of tissue plasminogen activator by intact endothelial cells
D. Decreased antithrombin III level
E. Mutation in factor V gene

31. At autopsy, a dark red solid mass in a vein not attached to the underlying venous wall always is:

A. Postmortem clot
B. Venous thrombus
C. Red thrombus
D. Mixed thrombus
E. Tumor embolus

32. Mural thrombi may be associated with all of the following disease except:

A. Myocardial infarction
B. DIC
C. Mitral valve stenosis
D. Atrial fibrillation
E. Extending cardiomyopathy

33. Mixed thrombus is composed of :

A. Platelets and fibrin

B. Platelets fibrin and red cells
C. Fibrin and red cells
D. Platelets and red cells
E. Platelets fibrin and neutrophils

34. In lung tissue sections in chronic pulmonary congestion, some cells celled "heart failure cells" can be found, and these cells are____
A. Dust cells
B. Lymphocytes
C. Hemosiderin-laden macrophages
D. Langhans cells
E. Neutrophils

35. All of these conditions are highly associated with an increased risk of thrombosis except:
A. Genetic factor V mutations
B. Prolonged bed rest after surgery
C. Myocardial infarction
D. Cancer metastases
E. Leukemia

36. Vegetations on heart valves in bacterial endocarditis are
A. Pale thrombi
B. Mixed thrombi
C. Red thrombi
D. Hyaline thrombi
E. Mural thrombi

37. What may be found in the microcirculation when DIC occurs?
A. Pale thrombi
B. Mixed thrombi
C. Red thrombi
D. Hyaline thrombi
E. Mural thrombi

38. Mural thrombus is
A. Pale thrombus
B. Mixed thrombus
C. Red thrombus
D. Hyaline thrombus
E. Fibrinous thrombus

39. Which kind of thrombus is completely composed of fibrin?
A. Pale thrombus
B. Mixed thrombus
C. Red thrombus
D. Hyaline thrombus
E. Mural thrombus

40. A 78-year-old woman falls in the bathtub and strikes the back of her head. Over the next 24 hours, she becomes increasingly somnolent. A head CT scan shows an accumulation of fluid beneath the dura, compressing the left cerebral hemisphere. Which of the following terms best describes this collection of fluid?
A. Hematoma
B. Purpura
C. Congestion
D. Petechia
E. Ecchymosis

41. Which of the followings is false concerning hemorrhage:
A. May be external or enclosed within a

tissue

B. The clinical significance of hemorrhage depends on the volume, rate and site of blood loss

C. Rupture of an aortic aneurysm can cause sudden death

D. Arteriolar vasoconstriction is the initial response to injury but is transient

E. Loss of up to 20% of the blood volume in healthy adult may result in hemorrhagic shock

42. A 44-year-old man has an acute myocardial infarction. Four days later, his cardiac output is reduced with an ejection fraction is of 25%. The patient becomes increasingly dyspneic, and a chest radiograph shows bilateral pulmonary edema. These pulmonary findings are most likely caused by which of the following mechanisms?

A. Lymphatic obstruction

B. Decreased plasma osmotic pressure

C. Decreased central venous pressure

D. Increased hydrostatic pressure

E. Acute inflammation

43. A 39-year-old woman comes to the physician because she has noticed a lump in her breast. Over the past 2 months, the left breast has become slightly enlarged compared with the right breast. On physical examination, the skin overlying the left breast is thickened, reddish-orange, and pitted. Mammography shows a 3-cm underlying density, and a fine-needle aspirate of the density indicates carcinoma. Which of the following mechanisms best explains the gross appearance of the left breast?

A. Venous thrombosis

B. Lymphatic obstruction

C. Ischemia

D. Chronic passive congestion

E. Chronic inflammation

44. The main cause of severe edema of the arm after breast cancer surgery is

A. Increased hydrostatic pressure

B. Reduced plasma osmotic pressure

C. Lymphatic obstruction

D. Sodium and water retention

E. Increased vascular permeability

45. Which of the following dose not result from chronic congestion?

A. Hemorrhage

B. Proliferation of parenchymal cells

C. Proliferation of interstitial cells

D. Hemosiderin deposition

E. Thrombotic complications

46. A 22-year-old second-year medical student develops a “red” face after being asked a question during lecture. Which of the following statements best describes this vascular reaction?

A. Active hyperemia

B. Acute congestion

C. Nonpalpable purpura
D. Passive hyperemia
E. Petechial hemorrhage

47. For the past week, a 61-year-old man has had increasing levels of serum AST and ALT. On physical examination, he has lower leg swelling with grade 2+ pitting edema from feet to knees and prominent jugular venous distention to the level of the mandible. Which of the following underlying conditions is most likely to be present?
A. Thrombocytopenia
B. Portal vein thrombosis
C. Chronic renal failure
D. Common bile duct obstruction
E. Congestive heart failure

48. Right-sided heart failure can cause congestion of
A. Lung brain and gastrointestinal tract
B. Lung and kidney liver and gastrointestinal tract
C. spleen, liver and gastrointestinal tract
D. spleen, liver and lung
E. spleen, lung and kidney

49. Which of the following is incorrect concerning chronic congestion of the liver:
A. “Nutmeg liver” is found grossly
B. The central vein and sinusoids are istended with blood
C. It is always be caused by right-sided heart failure
D. Fatty degeneration, atrophy and necrosis of hepatocytes are found
E. Will not result in cirrhosis

50. Which of the following is incorrect concerning arterial hyperemia:
A. It is an active process
B. It is found sites of inflammation
C. It is found in skeletal muscles during exercises
D. It is associated with rteriolar dilation
E. The tissue has a blue-red color

51. Which are the main components found in the alveolar spaces during acute pulmonary congestion:
A. Heart failure cells
B. Fibrin
C. Neutrophils
D. Blood-tinged edema fluids
E. Mucus


52. A patient with mitral stenosis for 15 years develops severe dyspnea especially during periods of physical activities or when lying down. Which one of the following states will not be found in the lungs of this patient:
A. The septa become thickened and fibrotic
B. “Heart failure cells” and red blood cells can be found in the alveolar spaces
C. The cut surface of the lung exudes a frothy mixture of surfactant-rich fluid

and blood

D. The alveolar capillaries are congested

E. Hyaline membranes line the alveolar spaces

53. A coronary artery is shown from the autopsy of a 61-year-old patient who had acute chest pain and difficulty breathing. What is most likely to have resulted from the pathologic findings in this artery?



A. Caseous necrosis

B. Coagulative necrosis

C. Apoptosis

D. Fibrinoid necrosis

E. Liquefactive necrosis

54. When going upstairs, a 60-year-old man had severe crushing chest pain, which lasted about 45 minutes and was not relieved by nitroglycerine. A coronary angiogram revealed 75% stenosis of the left anterior descending artery. Despite aggressive resuscitative measures, his condition deteriorated and he died. Which of the following microscopic findings is most likely to be present in his left ventricular myocardium at the time of death?

A. Extensive collagen deposition and abundant capillaries

B. Necrosis with neutrophils and macrophages

C. Edema and loss of cross striation

D. Lymphocytic interstitial infiltrates

E. Perivascular and interstitial amyloid deposition

55. Which is the main cause for chest pain when in pulmonary infarction?

A. Ventilatory disorder

B. Pleuritis in the infarct area

C. Stimulation of the bronchi by necrotic tissue

D. Stimulation of the bronchi by hemorrhage

E. Pulmonary infection

56. Which of these is the most common cause of pulmonary edema?

A. Pulmonary tuberculosis

B. Pneumonia

C. Left ventricular heart failure

D. Renal failure

E. Pulmonary embolism


57. A 72-year-old male has been treated in the hospital for pneumonia for the past several weeks. One day, he suddenly developed dyspnea, palpitations and chest pain. A pulmonary ventilation-perfusion scan reveals a high probability for a perfusion defect involving right segmental pulmonary arterial branch. Which of the followings is the most important factor favoring de-

velopment of these findings?

A. An increased platelet count
B. Altered blood flow with stasis
C. An increased white blood cell count
D. Cirrhosis of the liver
E. Atherosclerosis

58. Which of the following conditions would most likely be the cause of death following surgery for adenocarcinoma of the prostate for in a 70 years old man?

A. Inhalation of inorganic dust
B. Prolonged immobilization
C. Pulmonary contusion
D. Chronic bronchitis
E. Bronchial carcinoid



59. This is the gross appearance of a heart from autopsy of a 49-year-old man who had experienced worsening orthopnea for several months. He became dyspneic with exercise, but had no chest pain or peripheral edema. Which of the following underlying conditions do you most strongly suspect?

A. Dilated cardiomyopathy
B. Viral myocarditis
C. Rheumatic heart disease
D. Hypertension
E. Coronary thrombosis

60. A 16-year-old female presented to the emergency department, following a pedestrian vehicular accident. X-ray imaging revealed multiple fractures including the right lower leg and pelvis. After treatment her condition was stable. However, approximately 2 days after initial presentation, she became tachypnea and confused, and then died. This microscopic appearance was found at autopsy. Which of the followings in the most likely explanation?

A. Bronchopneumonia
B. Fat embolism
C. Metastatic carcinoma
D. Systemic miliary tuberculosis
E. Asthmatic attack

61. A solitary red wedge shaped area is found upon sectioning an organ, which is firm and accompanied by extensive hemorrhage. The base of this lesion is on the surface of the organ. In which of the following situations will this lesion most likely occur?

A. Heart with coronary thromboembolism
B. Lung with pulmonary thromboembolism
C. Liver with congestion
D. Kidney with septic embolus

E. Brain with cerebral hemorrhage

62. A 56-year-old female with a history of rheumatic heart disease and chronic atrial fibrillation presents eith dyspnea. Which of the following vascular complications is most likely to occur in this patient?
    A. Pulmonary embolism from a left ventricular mural thrombus
    B. Pulmonary emboli from thrombosis in a peripheral vein
    C. Systemic emboli from a left atrial mural thrombus
    D. Systemic emboli from a left ventricular mural thrombus
    E. Systemic emboli from a lower extremity deep vein thrombus
63. Which of the followings is not an outcome which results from pulmonary embolism?
    A. Sudden death
    B. Pulmonary infarction
    C. Interstitial pneumonia
    D. Pulmonary artery hypertension
    E. Right atrial enlargement
64. What consequence will be induced when small emboli pass out into the smaller, branching arterioles?
    A. Ischemic infarction
    B. Liquefactive necrosis
    C. Commonly no serious consequence
    D. Coagulative necrosis
    E. Septic infarction
65. After a long train trip, a 45-year-old previously healthy female is found to have swelling and tenderness of her legs. An ultrasound examination reveals findings suggestive of femoral vein thrombosis. Which of the following conditions is most likely to have contributed to this problem?
    A. Pregnancy
    B. Trousseau syndrome
    C. Prolonged inactivity
    D. Hypertension
    E. Chronic renal failure
66. Capillary channels are formed in organized thrombi, which then create conduits and interconnect with the original lumen. This process can partly restore blood flow and is called____
    A. Establishment of collateral circulation
    B. Organization
    C. Atherothrombosis
    D. Recanalization
    E. Propagation
67. Which of the following conditions is unlikely to cause edema?
    A. Congestive heart failure
    B. Liver cirrhosis
    C. Thrombosis
    D. Hemorrhage
    E. Inflammation

答案

中文题

1.C	2.D	3.E	4.B	5.E	6.D	7.D	8.C	9.B	10.E
11.B	12.C	13.C	14.A	15.C	16.B	17.A	18.D	19.A	20.A
21.D	22.E	23.A	24.C	25.C	26.C	27.A	28.D	29.D	30.A
31.A	32.B	33.D	34.E	35.E	36.D	37.E	38.E	39.A	40.E
41.C	42.B	43.D	44.A	45.D	46.D	47.C	48.B	49.C	50.D
51.C	52.E	53.E	54.E	55.B	56.D	57.E	58.B	59.A	60.D
61.C	62.C	63.B	64.E	65.A	66.A	67.C	68.C	69.C	70.D
71.E	72.A	73.D	74.C	75.A	76.E	77.A	78.B	79.A	80.E
81.C	82.A	83.D	84.C	85.E	86.B	87.E	88.E	89.B	90.E
91.C	92.E	93.C	94.B	95.B	96.E	97.C	98.E	99.C	100.E
101.E	102.E	103.C	104.D	105.A	106.C	107.D	108.A	109.B	110.E
111.C	112.D	113.D	114.E	115.C	116.D	117.A	118.B	119.D	120.A
121.B	122.A	123.D	124.B	125.C	126.D	127.E	128.A	129.E	130.A
131.C	132.D	133.E	134.E	135.D	136.B	137.E	138.A	139.C	140.D
141.E	142.A	143.C	144.B	145.A	146.A	147.C	148.D	149.A	150.B
151.A	152.B	153.C	154.A	155.B	156.A	157.D	158.E	159.D	160.B

英文题

1.D	2.D	3.C	4.E	5.D	6.A	7.B	8.C	9.A	10.A
11.D	12.B	13.B	14.C	15.B	16.C	17.C	18.A	19.B	20.C
21.B	22.A	23.D	24.C	25.E	26.A	27.A	28.B	29.E	30.B
31.A	32.B	33.B	34.C	35.E	36.A	37.D	38.B	39.D	40.A
41.E	42.D	43.B	44.C	45.B	46.A	47.E	48.C	49.E	50.E
51.D	52.E	53.B	54.B	55.B	56.C	57.B	58.B	59.D	60.B
61.B	62.C	63.C	64.C	65.C	66.D	67.D			

3　炎症

中文题

1. 关于炎症的阐述，错误的是
   A. 炎症是白细胞吞噬及清除有害因子的反应
   B. 炎症是机体对于损伤因子的防御反应
   C. 炎症是机体损伤及抗损伤过程
   D. 炎症的局部反应为红、肿、热、痛和功能障碍
   E. 炎症的全身反应包括发热、和白细胞计数升高
2. 最常见的致炎因子是
   A. 物理性因子
   B. 化学性因子
   C. 生物性因子
   D. 机械性因子
   E. 免疫性因子
3. 不感染脊髓灰质炎的人群是
   A. 体弱的儿童
   B. 体健的儿童
   C. 不去公共场所的儿童
   D. 接受过计划免疫的儿童
   E. 消化道疾患的儿童
4. 炎症的最重要标志是
   A. 变性
   B. 坏死
   C. 渗出
   D. 充血
   E. 增生
5. 炎症过程中，组织损伤的主要表现是
   A. 渗出
   B. 充血
   C. 变质
   D. 间质增生
   E. 实质增生
6. 属于感染性炎症的是
   A. 物理性炎症
   B. 内源性化学性炎症
   C. 生物性炎症
   D. 免疫性炎症
   E. 外源性化学性炎症
7. 炎症的本质是
   A. 代偿
   B. 修复
   C. 增生
   D. 适应
   E. 防御
8. 不引起肝脏炎症的化学因子有
   A. 异烟肼
   B. 利福平
   C. 胆汁
   D. 尿素
   E. 酒精
9. 属于炎症的变质性病变是
   A. 心外膜的纤维素
   B. 蛛网膜下腔的脓液
   C. 表皮内水泡
   D. 乙型脑炎神经细胞溶解消失
   E. 慢性肝炎汇管区炎细胞

10. 女性，30 岁。十天来厌油腻饮食，恶心、呕吐，皮肤黄染。入院检查，肝肿大、压痛，ALT 增高，HBsAg 和 HbcAg 阳性。不属于患者肝脏变质改变的是
A. 肝细胞疏松化
B. 肝细胞灶状坏死
C. 淋巴细胞浸润
D. 肝细胞嗜酸性变
E. 肝细胞气球样变

11. 不符合炎症性增生的是
A. 实质细胞增生
B. 纤维组织增生
C. 淋巴组织增生
D. 单核巨噬细胞增生
E. 不典型增生

12. 属于炎症性增生的病变是
A. 风湿性动脉炎血管壁粘液变性
B. 风湿性关节炎关节肿胀
C. 风湿性心肌炎时风湿小结形成
D. 风湿性心内膜炎疣状赘生物形成
E. 风湿性心外膜炎纤维素渗出

13. 急性重型肝炎引起肝功能衰竭是由于
A. 肝出血
B. 肝充血
C. 肝内炎细胞浸润
D. 肝细胞大片坏死
E. 小胆管增生

14. 流行性乙型脑炎引起患者智力低下是由于
A. 脑内炎细胞浸润
B. 脑内胶质细胞增生
C. 脑内血管扩张
D. 脑水肿
E. 脑神经细胞变性坏死

15. 不符合炎症性增生的描述是
A. 具有抗损伤的作用
B. 具有修复功能
C. 可引起间质增生
D. 可引起实质增生
E. 对机体有益无害防御

16. 病毒性肝炎时，不属于增生的病变是
A. 汇管区淋巴滤泡形成
B. 出现多量双核肝细胞
C. 出现多量小胆管
D. 出现多量纤维组织
E. 出现多量有脂肪空泡的肝细胞

17. 急性炎症反应中，最先出现的血管变化是
A. 血管扩张
B. 血流缓慢
C. 血流加快
D. 血管收缩
E. 血流停滞

18. 炎症的中，最先出现的血管变化是
A. 血管扩张
B. 血流缓慢
C. 血流加快
D. 血管收缩
E. 血流停滞

19. I 超敏反应引起的血管变化是
A. 速发短暂反应
B. 速发持续反应
C. 迟发持续反应
D. 细动脉短暂收缩

E. 静脉持续淤血

20. 重度烧伤引起的血管变化是

A. 速发短暂反应

B. 速发持续反应

C. 迟发持续反应

D. 细动脉短暂收缩

E. 静脉持续淤血

21. 紫外线晒伤引起的血管变化是

A. 速发短暂反应

B. 速发持续反应

C. 迟发持续反应

D. 细动脉短暂收缩

E. 静脉持续淤血

22. 急性炎症反应中，最先渗出的是

A. 纤维蛋白

B. 球蛋白

C. 白蛋白

D. 中性粒细胞

E. 单核细胞

23. 急性炎症反应中，不符合血管通透性增加的因素是

A. 致炎因子引起的内皮细胞损伤

B. 炎细胞介导的内皮细胞损伤

C. 内皮细胞间隙增大

D. 新生毛细血管的高通透性

E. 内皮细胞胞浆滤过增多

24. 急性炎症反应中，不符合血管通透性增加的因素是

A. 致炎因子引起的内皮细胞损伤

B. 炎细胞介导的内皮细胞损伤

C. 内皮细胞吞饮小泡增多

D. 小静脉的高通透性

E. 内皮细胞间隙增大

25. 急性炎症时，组织肿胀的主要原因是

A. 纤维组织增生

B. 实质细胞变质

C. 肉芽组织增生

D. 实质细胞增生

E. 充血及血液成分渗出

26. 不符合炎症渗出液的描述是

A. 液体浑浊

B. 液体比重高

C. 液体静置后凝固

D. 液体内蛋白含量高

E. 液体内含极少细胞

27. 病毒性肝炎时，属于渗出病变是

A. 汇管区淋巴细胞浸润

B. 肝细胞嗜酸性小体形成

C. 肝细胞灶状坏死

D. 肝细胞水肿

E. 肝细胞嗜酸性变

28. 符合炎性渗出液的描述是

A. 液体比重低

B. 外观清亮

C. 细胞含量少

D. 蛋白含量低

E. 液体静置后可凝固

29. 不符合炎症时液体渗出机制的描述是

A. 微循环内液体静压升高

B. 组织渗透压升高

C. 血管内皮细胞损伤

D. 新生毛细血管增加

E. 小静脉增多

30. 不符合漏出液特点的是

A. 液体比重低
B. 液体静置后不凝固
C. 液体内蛋白含量低
D. 液体内细胞数小于 0.5×10^9/L
E. Rivalta 反应阳性

31. 不符合渗出液作用的描述是
A. 稀释毒素
B. 带来抗体
C. 带来补体
D. 带来氧和营养物
E. 对机体有益无害

32. 符合渗出液作用的描述是
A. 不引起组织粘连
B. 不引起组织水肿
C. 不影响器官功能
D. 不利于病原体扩散
E. 不利于清除病原体

33. 不符合炎症时白细胞渗出的描述是
A. 白细胞靠边
B. 白细胞与内皮细胞粘着
C. 内皮细胞主动收缩，使白细胞得以游出
D. 白细胞通过阿米巴运动游出血管
E. 白细胞向炎症灶定向运动

34. 不符合白细胞渗出的描述是
A. 肿瘤坏死因子（TNF）可促进粘附分子的表达
B. 白细胞以阿米巴运动方式从内皮细胞缝隙游出
C. 炎症的不同阶段，游出的白细胞种类不同
D. 红细胞漏出是主动过程
E. 趋化因子的作用是有特异性的

35. 不符合急性炎症时白细胞渗出的描述是
A. 中性粒细胞进入边流附壁
B. 附壁是内皮细胞和白细胞粘附因子互相识别的结果
C. 游出是指红细胞进入血管外
D. 在趋化因子作用下，到达炎症部位
E. 白细胞三烯具有趋化作用

36. 与白细胞杀菌能力有关的疾病是
A. DiGeorge 综合征
B. Bruton 综合征
C. T 细胞缺陷病
D. 慢性肉芽肿病
E. 丙种球蛋白缺乏

37. 影响白细胞吞噬与脱颗粒的疾病是
A. DiGeorge 综合征
B. Bruton 综合征
C. Chediak-Higashi 综合征
D. 慢性肉芽肿病
E. 丙种球蛋白缺乏

38. 趋化作用是指
A. 白细胞靠边
B. 白细胞附壁
C. 白细胞游出
D. 白细胞定向运动
E. 白细胞吞噬作用

39. 与白细胞杀菌能力无关的过程是
A. 识别和粘着
B. 包围吞入
C. 杀伤降解
D. 吞噬溶酶体的形成

E. 趋化作用

40. 不参与吞噬细胞识别病原体的因子是

A. C3b 受体

B. 溶酶体酶

C. C3b

D. Fc 受体

E. 抗体

41. 病毒感染的病灶内最常见的细胞是

A. 中性粒细胞

B. 嗜酸粒细胞

C. 淋巴细胞、单核细胞

D. 浆细胞

E. 肥大细胞

42. 沙门杆菌感染的炎症反应中最多见的细胞是

A. 淋巴细胞

B. 肥大细胞

C. 嗜酸粒细胞

D. 中性粒细胞

E. 单核细胞

43. 葡萄球菌感染的炎症灶内主要的炎症细胞是

A. 单核细胞

B. 淋巴细胞

C. 嗜酸粒细胞

D. 浆细胞

E. 中性粒细胞

44. 寄生虫感染时，病灶内最多见的炎症细胞是

A. 中性粒细胞

B. 淋巴细胞

C. 嗜碱粒细胞

D. 浆细胞

E. 嗜酸粒细胞

45. 脓细胞是

A. 转化的淋巴细胞

B. 含 Russel 小体的浆细胞

C. 吞噬脂质的巨噬细胞

D. 变性坏死的中性粒细胞

E. 变性坏死的嗜酸粒细胞

46. 慢性炎症组织内侵润的细胞主要是

A. 淋巴细胞

B. 中性粒细胞

C. 嗜酸粒细胞

D. 嗜碱粒细胞

E. 肥大细胞

47. 肉芽肿性炎增生的细胞是

A. 巨噬细胞

B. 淋巴细胞

C. 浆细胞

D. 中性粒细胞

E. 嗜酸性粒细胞

48. 不符合巨噬细胞作用的描述是

A. 参与免疫反应

B. 具有吞噬细菌的作用

C. C3b 增强其吞噬作用

D. 常见于炎症早期

E. 可转化为类上皮细胞、多核巨细胞

49. 不符合嗜酸粒细胞的描述是

A. 来源于骨髓

B. 运动能力弱

C. 无吞噬功能

D. 见于寄生虫病及过敏性疾病

E. 见于某些部位的亚急性或慢性炎症

50. 合成免疫球蛋白的细胞是
A. 单核细胞
B. T 淋巴细胞
C. 浆细胞
D. 中性粒细胞
E. 嗜酸粒细胞

51. 不符合浆细胞的描述是
A. 不存在于血液
B. 无吞噬功能
C. 来源于 T 淋巴细胞
D. 椭圆形，核位于胞浆一端，核染色质呈车轮状分布
E. 胞浆内有丰富的粗面内质网

52. 可形成多核巨细胞的细胞是
A. 浆细胞
B. 巨噬细胞
C. 中性粒细胞
D. 嗜酸粒细胞
E. 肥大细胞

53. 出现在新鲜梗死灶边缘的细胞是
A. 浆细胞
B. 单核细胞
C. 中性粒细胞
D. 嗜酸粒细胞
E. 肥大细胞

54. 出现在哮喘反应的支气管壁内的细胞是
A. 浆细胞
B. 单核细胞
C. 中性粒细胞
D. 嗜酸粒细胞
E. 肥大细胞

55. 不符合中性粒细胞的描述是
A. 又称小噬细胞
B. 有活跃的运动能力
C. 常出现于炎症晚期
D. 吞噬细菌和坏死组织
E. 有丰富的中性颗粒

56. 不符合肥大细胞的描述是
A. 胞浆嗜碱性
B. 可产生肝素
C. 可产生组织胺
D. 可产生 5-羟色胺
E. 具有吞噬功能

57. 急性炎症中，引起血管通透性升高的速发反应的炎症介质是
A. 补体
B. 细菌毒素
C. 过敏毒素
D. 组胺
E. 坏死分解产物

58. 具有化学趋化作用的炎性介质是
A. 组胺
B. C5a
C. C3b
D. 缓激肽
E. 活性氧代谢产物

59. 具有调理化作用的炎性介质是
A. 组胺
B. C5a
C. C3b
D. 缓激肽
E. 活性氧代谢产物

60. 引起疼痛的炎症介质是

A. 组胺
B. 溶酶体酶
C. 缓激肽
D. 白细胞三烯
E. 细菌产物

61. 不引起血管通透性增高的炎症介质是
A. 组胺
B. C3a
C. 缓激肽
D. C5a
E. 白细胞三烯

62. 引起发热的炎症介质是
A. 组胺
B. 溶酶体酶
C. 缓激肽
D. 白细胞三烯
E. TNF

63. 不符合补体的描述是
A. 抗原抗体复合物可激活补体
B. 可引起血管壁的通透性增高
C. 具有趋化作用
D. 引起发热
E. C3 和 C5 是最重要的炎症介质

64. 5-羟色胺主要存在于
A. 巨噬细胞
B. 血管内皮细胞
C. 肥大细胞
D. 淋巴细胞
E. 嗜酸粒细胞

65. 不符合炎症介质的描述是
A. 可由体液释放
B. 可由细胞释放
C. 可具有趋化作用
D. 可引起发热和疼痛
E. 不引起组织损伤

66. 不符合组胺的描述是
A. 存在于肥大细胞、嗜碱粒细胞和血小板中
B. 引起血管扩张
C. 对嗜酸粒细胞有趋化作用
D. 引起疼痛和发热
E. 导致血管通透性升高

67. 不符合 TNF 的描述是
A. 由激活的淋巴细胞和单核细胞产生
B. 内毒素等致炎因子可促进 TNF 分泌
C. 可促进白细胞粘附
D. 为中性粒细胞趋化因子
E. 引起疼痛

68. 不符合化学趋化作用的描述是
A. 趋化因子吸引白细胞定向运动
B. 阳离子蛋白质对单核细胞有趋化作用
C. TNF 促进中性粒细胞聚集
D. IL-8 对中性粒细胞有趋化作用
E. 血小板激活因子不具有趋化作用

69. 不符合缓激肽的描述是
A. 可由体液释放
B. 可引起细动脉扩张
C. 可引起内皮细胞收缩
D. 可引起发热
E. 可引起血管通透性增加

70. 不符合补体激活的描述是
A. 抗原抗体结合可激活补体

B. 细菌毒素可激活补体
C. 坏死组织释放的酶可激活补体
D. 中性粒细胞释放的酶可激活补体
E. 纤维蛋白降解产物不激活补体

71. 不符合纤维蛋白溶解系统的描述是
A. 可激活激肽系统
B. 可激活补体系统
C. 可使血管通透性增加
D. 可导致组织损伤
E. 可引起白细胞渗出

72. 不符合炎症介质的描述是
A. 引起血管扩张
B. 导致组织损伤
C. 具有趋化作用
D. 引起发热和疼痛
E. 引起组织增生

73. 不符合炎症全身反应的描述是
A. 发热
B. 末梢血白细胞计数增多
C. 单核巨噬细胞系统功能增强
D. 末梢血出现幼稚的白细胞
E. 实质器官变性、坏死、功能降低

74. 炎症局部发红的原因是
A. 发热
B. 血管扩张、血流加快
C. 毛细血管增生
D. 静脉淤血
E. 淋巴液淤积

75. 炎症局部肿胀的原因是
A. 淤血
B. 纤维增生
C. 毛细血管增生
D. 淋巴液淤积
E. 炎症性充血、渗出

76. 引起炎症局部疼痛的因素是
A. C3a、C5a
B. 溶酶体酶、氧自由基
C. PGE_2、缓激肽
D. TNF、IL-1
E. 白细胞三烯、细菌产物

77. 炎症时，不引起发热的因素是
A. 前列腺素
B. IL-6
C. TNF
D. IL-1
E. 白细胞三烯、细菌产物

78. 可导致迅速死亡的炎症性疾病是
A. 丹毒
B. 淋巴结结核
C. 青霉素过敏
D. 蜂窝织炎性阑尾炎
E. 伤寒

79. 不符合炎症时末梢血变化的描述是
A. 可出现核左移
B. 白细胞计数可增多
C. 白细胞计数可减少
D. 淋巴细胞比例升高
E. 血小板计数升高

80. 不符合炎症的描述是
A. 血管反应是中心环节
B. 有利于清除致病因子
C. 有利于修复损伤
D. 有利于维持器官功能
E. 有利于组织再生

81. 细菌进入血中并大量繁殖，引起全身中毒症状，称之为
A. 毒血症
B. 病毒血症
C. 菌血症
D. 败血症
E. 脓毒败血症

82. 不符合毒血症的描述是
A. 可出现高热、寒战
B. 可出现脾脏及淋巴结肿大
C. 可出现心、肝、肾等实质细胞变性或坏死
D. 可出现中毒性休克
E. 血中可培养出致病菌

83. 不符合败血症的描述是
A. 可出现高热、寒战
B. 可出现脾脏及淋巴结肿大
C. 可出现皮肤粘膜点状出血
D. 可出现中毒性休克
E. 血培养无致病菌

84. 女性，12 岁，发热（39℃），髋关节剧痛、活动受限，伴有皮肤粘膜多发性出血斑、脾脏轻度肿大。查血，白细胞总数 20×10^9/L，分叶核 60%，杆状核 25%。血培养可检出葡萄球菌。诊断符合
A. 毒血症
B. 菌血症
C. 败血症
D. 脓毒败血症
E. 病毒血症

85. 不符合败血症的描述是
A. 肝肿大
B. 高热寒战
C. 多器官的多发性小脓肿
D. 皮肤、粘膜点状出血
E. 脾肿大

86. 符合毒血症的描述是
A. 破伤风患者出现全身肌肉痉挛
B. 血培养细菌阳性，无全身中毒症状
C. 血液中病毒检查阳性
D. 全身中毒症状，血培养细菌阳性
E. 全身器官有多发性脓肿，血培养细菌阳性

87. 符合败血症的描述是
A. 破伤风患者出现全身肌肉痉挛
B. 血培养细菌阳性，无全身中毒症状
C. 血液中病毒检查阳性
D. 全身中毒症状，血培养细菌阳性
E. 全身器官有多发性脓肿，血培养细菌阳性

88. 符合脓毒败血症的描述是
A. 破伤风患者出现全身肌肉痉挛
B. 血培养细菌阳性，无全身中毒症状
C. 血液中病毒检查阳性
D. 全身中毒症状，血培养细菌阳性
E. 全身器官有多发性脓肿，血培养细菌阳性

89. 不符合化脓性炎的结局的描述是
A. 吸收消散
B. 包裹
C. 纤维组织增生修复
D. 转为慢性
E. 引起菌血症

90. 有利于炎症痊愈的因素是
A. 治疗及时
B. 病原体数量多
C. 病原体毒力强
D. 机体抵抗力差
E. 机体变态反应剧烈

91. 不利于炎症痊愈的因素是
A. 病原体数量少
B. 病原体毒力弱
C. 机体抵抗力强
D. 治疗及时
E. 机体变态反应剧烈

92. 男性，20 岁，胆囊切除、胆总管引流术后，引流处拔管后胆汁外流。此病变应诊断为
A. 糜烂
B. 溃疡
C. 机化
D. 瘘管
E. 窦道

93. 男性，28 岁，胫骨慢性骨髓炎伴有皮肤破溃、流脓。病变符合
A. 瘘管
B. 窦道
C. 溃疡
D. 糜烂
E. 机化

94. 腰椎结核患者，背部脓肿破裂，脓液长期流出体表，此病变是
A. 溃疡
B. 窦道
C. 瘘管
D. 空洞
E. 糜烂

95. 属于炎症淋巴道播散引起的疾病是
A. 输尿管结核
B. 干酪性肺炎
C. 化脓性淋巴结炎
D. 丹毒
E. 蜂窝织炎性阑尾炎

96. 不属于炎症淋巴道播散的引起的疾病是
A. 化脓性淋巴结炎
B. 急性淋巴管炎
C. 肺门淋巴结结核
D. 腰椎结核
E. 颈部淋巴结结核

97. 风湿性关节炎属于
A. 浆液性炎
B. 蜂窝织炎
C. 脓肿
D. 纤维素性炎
E. 肉芽肿性炎

98. 急性炎症反应中出血的机制是
A. 淤血
B. 纤维蛋白原合成减少
C. 纤维蛋白溶解
D. 血管壁的损伤
E. 凝血因子缺乏

99. 不符合浆液性炎的描述是
A. 多见于疏松组织、浆膜和粘膜
B. 在疏松组织形成积脓
C. 在浆膜腔形成积液
D. 在粘膜形成“卡他性炎”

E. 渗出液中含有少量白细胞和纤维素

100. 男性，5 岁，左腿烫伤后红肿、疼痛，数小时后起水疱。其病变为

A. 出血性炎

B. 变质性炎

C. 炎性水肿

D. 炎性积液

E. 蜂窝织炎

101. 属于出血性炎的疾病是

A. 细菌性痢疾

B. 肠阿米巴病

C. 流行性脑脊髓膜炎

D. 钩端螺旋体病

E. 丹毒

102. 不属于浆液性炎的病变是

A. 湿性结核性胸膜炎

B. 风湿性关节炎

C. 肾盂积水

D. 单纯疱疹

E. 皮肤 II 度烧伤水疱

103. 鼠疫杆菌感染引起肺部病变属于

A. 浆液性炎

B. 纤维素性炎

C. 脓肿

D. 蜂窝织炎

E. 出血性炎

104. 卡他性炎是指

A. 局灶性化脓性炎

B. 疏松组织的弥漫性化脓性炎

C. 体腔大量脓液蓄积的炎症

D. 粘膜的浆、粘液渗出的炎症

E. 大量纤维蛋白渗出的炎症

105. 有关纤维素性炎的描述，错误的是

A. 多见于粘膜、浆膜和肺

B. 渗出大量纤维蛋白原

C. 发生在浆膜时，形成伪膜性炎

D. 可以发生机化，引起粘连

E. 常伴有中性粒细胞渗出

106. 不属于纤维素性炎的是

A. 绒毛心

B. 假膜性肠炎

C. 大叶性肺炎

D. 干型结核性腹膜炎

E. 湿型结核性腹膜炎

107. 白喉杆菌感染引起

A. 浆液性炎

B. 蜂窝织炎

C. 脓肿

D. 纤维素性炎

E. 肉芽肿性炎

108. 男性，45 岁，四肢小关节疼痛，肿胀、活动不便。诊断为类风湿性关节炎。其关节病变可表现为

A. 大量浆液渗出

B. 大量纤维素渗出

C. 大量中性粒细胞渗出

D. 大量单核细胞渗出

E. 大量嗜酸粒细胞渗出

109. 假膜性炎的特征性的渗出物是

A. 浆液

B. 纤维素

C. 中性粒细胞

D. 巨噬细胞

E. 淋巴细胞

110. 属于假膜性炎的是
A. 发生在腹膜的纤维素性炎
B. 发生在肠粘膜的纤维素性炎
C. 发生在胸膜的纤维素性炎
D. 发生在肺的纤维素性炎
E. 发生在心包膜的纤维素性炎

111. 女性，25 岁。患风湿病 5 年。听诊可以听到心包摩擦音。心包腔的渗出物的主要成分为
A. 淋巴细胞
B. 纤维素
C. 中性粒细胞
D. 单核细胞
E. 红血球

112. 属于纤维素性炎的是
A. 局灶性大量中性粒细胞侵润及组织坏死液化
B. 疏松组织的弥漫性化脓性炎症
C. 体腔大量脓液蓄积
D. 粘膜的浆、粘液渗出
E. 白喉引起的喉和气管的病变

113. 金黄色葡萄球菌感染引起
A. 浆液性炎
B. 蜂窝织炎
C. 脓肿
D. 纤维素性炎
E. 肉芽肿性炎

114. 有关化脓性炎症的描述，不正确的是
A. 有大量中性粒细胞渗出
B. 有脓液形成
C. 有大量纤维素渗出
D. 常伴有不同程度的组织坏死
E. 蜂窝织炎和脓肿均为化脓性炎

115. 蜂窝织炎性阑尾炎最主要的病变特点是
A. 中性粒细胞弥漫浸润阑尾各层组织
B. 中性粒细胞间粘膜表面渗出
C. 阑尾充血、水肿
D. 阑尾壁内脂肪组织增生
E. 阑尾腔内可见粪石

116. 非化脓性炎症性疾病是
A. 急性蜂窝织炎性阑尾炎
B. 病毒性肝炎
C. 丹毒
D. 急性细菌性心内膜炎
E. 肾盂肾炎

117. 男性，30 岁，颈部红肿 8 天，检查时，发现颈部皮肤红肿区约 6×5cm，表面可见多个脓头，压痛明显。病变符合
A. 疖
B. 痈
C. 蜂窝织炎
D. 淋巴管炎
E. 丹毒

118. 患者女，17 岁，转移性右下腹痛已四天，腹痛渐加重，伴有发热（39℃），右下腹有压痛、反跳痛，并可触及一有压痛的肿块。符合患者病情的诊断是
A. 急性单纯性阑尾炎
B. 急性蜂窝织炎性阑尾炎
C. 急性蜂窝织炎性阑尾炎伴阑尾周围脓肿
D. 慢性阑尾炎
E. 阑尾粘液囊肿

119. 体腔内大量脓液蓄积，应称之为
A. 蜂窝织炎
B. 脓肿
C. 积脓
D. 表面化脓
E. 痈

120. 女性，30 岁，分娩后 3 个月。1 月前左乳外上象限红、肿、热、痛，经热敷后病变局限。检查时，左乳外上象限有一直径 3×3cm 的结节，有波动感。穿刺液内可见
A. 大量红细胞
B. 大量单核淋巴细胞
C. 大量中性粒细胞
D. 大量清亮液体
E. 大量浆细胞

121. 关于化脓性炎症的结局，错误的是
A. 吸收消散
B. 机化包裹
C. 形成窦道或瘘管
D. 癌变
E. 血道播散

122. 肛门周围深部脓肿可发生
A. 溃疡
B. 空洞
C. 炎性息肉
D. 瘘管
E. 糜烂

123. 男性，80 岁。7 天来因感冒，咳嗽，病情不断加重。X 光显示双肺散在小片性实变灶，以下肺为重。其肺部的病变属
A. 化脓性炎
B. 纤维素性炎
C. 浆液性炎
D. 肉芽肿性炎
E. 变质性炎

124. 男性，30 岁，手指挤压伤后 2 天，疼痛加剧，红肿并向掌部蔓延，与正常组织无明显界限。这种病变应诊断为
A. 浆液性炎
B. 纤维素性炎
C. 脓肿
D. 蜂窝织炎
E. 坏疽

125. 男性，40 岁，背部红、肿、热、痛 7 天，表面有三、四个破口，流黄色汁液。其病变由
A. 溶血性链球菌感染引起
B. 大肠杆菌感染引起
C. 金黄色葡萄球菌感染引起
D. 变形杆菌感染引起
E. 绿脓杆菌引起

126. 溶血性链球菌感染最常引起
A. 浆液性炎
B. 假膜性炎
C. 出血性炎
D. 蜂窝织炎
E. 脓肿

127. 蜂窝织炎是指
A. 局灶性大量中性粒细胞浸润及组织坏死液化
B. 疏松组织的弥漫性化脓性炎症
C. 体腔大量脓液蓄积
D. 粘膜的浆、粘液渗出

E. 急性痢疾杆菌的典型肠病变

128. 脓肿是指

A. 局灶性大量中性粒细胞浸润及组织坏死液化

B. 疏松组织的弥漫性化脓性炎症

C. 体腔大量脓液蓄积

D. 粘膜的浆、粘液渗出

E. 急性痢疾杆菌的典型肠病变

129. 男性，47 岁，慢性腹泻，脓血便 7 年，临床诊断为慢性溃疡性结肠炎，肠道病变常表现为

A. 炎性息肉形成

B. 假膜形成

C. 肉芽肿形成

D. 瘘管形成

E. 炎性假瘤形成

130. 有关慢性炎症的描述，错误的是

A. 可有实质细胞增生

B. 可有肉芽肿形成

C. 可有纤维组织增生

D. 可形成多发性息肉

E. 可形成肿瘤结节

131. 不符合慢性炎症的病变是

A. 中性粒细胞为主的炎细胞浸润

B. 炎症性息肉

C. 炎性假瘤

D. 多量结缔组织增生

E. 炎性肉芽肿形成

132. 女性，40 岁，体检时发现肺上叶有直径 3cm 的结节。手术切除后，病理诊断为肺的炎性假瘤，不符合炎性假瘤的病变是

A. 纤维组织增生

B. 肉芽肿形成

C. 小血管增生

D. 单核、淋巴细胞浸润

E. 肺泡上皮增生

133. 不属于慢性炎症的病变是

A. 鼻息肉

B. 大肠炎性息肉

C. 大肠腺瘤性息肉

D. 胃炎症性息肉

E. 宫颈息肉

134. 符合慢性炎症的描述是

A. 起病急

B. 病程短

C. 以增生性病变为主

D. 常以粒细胞浸润为主

E. 局部血管通透性增高

135. 有关慢性炎症的描述，错误的是

A. 起病缓慢

B. 病程长

C. 以渗出性病变为主

D. 以巨噬细胞、淋巴细胞浸润为主

E. 致炎因子持续存在

136. 女性，28 岁，有 10 年的吸烟史。近三年来慢性咳嗽、咳白色泡沫痰，每年发病时间 3 个月以上。气管镜粘膜活检的病变是

A. 纤维素性炎

B. 化脓性炎

C. 慢性炎

D. 肉芽肿性炎

E. 变质性炎

137. 女性，46 岁，颈部淋巴结肿大，手术切除。肿大的淋巴结内可见淡黄色、质软的干酪样坏死。符合淋巴结病变的描述是
A. 化脓性炎
B. 纤维素性炎
C. 肉芽肿性炎
D. 浆液性炎
E. 出血性炎

138. 结核性肉芽肿中的多核巨细胞主要来源于
A. 纤维母细胞
B. 中性粒细胞
C. 肥大细胞
D. 单核细胞
E. 浆细胞

139. 有关肉芽肿性炎的描述，错误的是
A. 均为慢性炎症
B. 主要由巨噬细胞增生构成
C. 增生的巨噬细胞可以转变为多核巨细胞
D. 不同的病因可以引起形态不同的肉芽肿
E. 可根据典型的肉芽肿形态作出病因诊断

140. 结核性肉芽肿不具有的成分是
A. Langhans 巨细胞
B. 类上皮细胞
C. 纤维素样坏死
D. 淋巴细胞
E. 纤维母细胞

141. 不引起异物性肉芽肿的是
A. 手术缝线
B. 滑石粉
C. 石棉
D. 血吸虫卵
E. 铍

142. 属于感染性肉芽肿的是
A. 风湿性肉芽肿
B. Crohn 病
C. 结核性肉芽肿
D. 异物肉芽肿
E. 结节病

143. 女性，20 岁，阑尾切除术后，残留在体内的缝线周围的病理改变是
A. 肉芽肿性炎
B. 炎性假瘤
C. 机化
D. 化脓性炎
E. 蜂窝织炎

144. 麻风杆菌感染引起
A. 浆液性炎
B. 蜂窝织炎
C. 脓肿
D. 纤维素性炎
E. 肉芽肿性炎

145. 伤寒小结中的伤寒细胞来源于
A. 中性白细胞
B. 单核细胞
C. 肥大细胞
D. 淋巴细胞
E. 嗜酸粒细胞

146. 原因不明的肉芽肿性炎是
A. 结核性肉芽肿

B. 麻风性肉芽肿
C. 梅毒树胶肿
D. 肉芽瘤病
E. 伤寒肉芽肿

147. 慢性炎症的主要原因是
A. 急性炎症对组织的破坏
B. 细菌毒力小
C. 细菌数量少
D. 自身免疫反应
E. 致炎因子持续存在

148. 慢性炎症中，寿命长的炎细胞是
A. 中性白细胞
B. 单核细胞
C. 肥大细胞
D. 浆细胞
E. 嗜酸粒细胞

149. 慢性炎症中，对纤维母细胞增生无刺激作用的是
A. 表皮生长因子
B. 血小板源性生长因子
C. 转化生长因子
D. 肿瘤坏死因子
E. IL-2

150. 慢性炎症中，对血管内皮细胞无刺激作用的是
A. 表皮生长因子
B. 血小板源性生长因子
C. 转化生长因子
D. 肿瘤坏死因子
E. IL-2

151. 慢性炎症中，对毛细血管形成无刺激作用的是
A. 表皮生长因子
B. 血小板源性生长因子
C. 转化生长因子
D. 肿瘤坏死因子
E. 血管内皮生长因子

152. 慢性炎症中，对胶原纤维形成无刺激作用的是
A. 表皮生长因子
B. 血小板源性生长因子
C. 转化生长因子
D. 肿瘤坏死因子
E. 纤维母细胞生长因子

153. exudation
A. 变质
B. 渗出
C. 增生
D. 浸润
E. 感染

154. inflammatory mediator
A. 趋化因子
B. 渗出液
C. 吞噬作用
D. 炎症介质
E. 炎症反应

155. granuloma
A. 肉芽组织
B. 肉芽肿
C. 炎性息肉
D. 脓肿
E. 积脓

156. suppurative inflammation
A. 纤维素性炎

B. 蜂窝织炎
C. 浆液性炎
D. 化脓性炎
E. 增生性炎

157. septicemia
A. 菌血症
B. 败血症
C. 毒血症
D. 病毒血症
E. 脓毒血症

158. chemotasis
A. 趋化因子
B. 趋化作用
C. 白细胞游出
D. 白细胞附壁
E. 白细胞靠边

159. inflammatory pseudotumor
A. 炎性息肉
B. 炎性肉芽组织
C. 炎性肉芽肿
D. 炎性假瘤
E. 炎性增生

160. abscess
A. 积脓
B. 积液
C. 脓肿
D. 瘘管
E. 空洞

英文题

1. A 36-year-old man has had a chronic cough with fever and weight loss for the past month. A chest radiograph reveals multiple nodules of 1 to 4 cm, some of which demonstrate cavitation in the upper lobes. A sputum sample reveals the presence of acid fast bacilli. Which of the following cells is the most important in the development his lung lesions?
A. Macrophage
B. Fibroblast
C. Neutrophil
D. Mast cell
E. Platelet

2. After two weeks in the hospital following a fall in which he incurred a fracture of his left femoral trochanter, a 78-year-old man now has a left leg that is swollen, particularly the lower leg below the knee. He experiences pain on movement of this leg, and there is tenderness to palpation. Which of the following complications is most likely to occur next after these events?
A. Gangrenous necrosis of the foot
B. Pulmonary thromboembolism
C. Disseminated intravascular coagulation
D. Hematoma of the thigh
E. Soft tissue sarcoma

3. A 62-year-old man has had increasing dyspnea for 6 years. He has no cough

or fever. He had inhaled silica dust for many years in his job. A chest x-ray now shows increased interstitial markings and parenchymal nodules of 1 to 3cm. His pulmonary problems are most likely due to which of the following inflammatory processes?


A. Neutrophilic infiltration with release of leukotrienes
B. Foreign body giant cell formation
C. Plasma cell production of immunoglobulin
D. Release of growth factors by macrophages
E. Histamine release by mast cells

4. A 4-year-old child reaches up to the stove and touches a pot of boiling soup. Within several hours, there is marked erythema of the skin of fingers on the child's right hand, and small blisters appear on the finger pads. Which of the following terms best describes this process?

A. Fibrinous inflammation
B. Serous inflammation
C. Purulent inflammation
D. Ulceration
E. Granulomatous inflammation

5. Patients have pharyngeal infections. The most typical clinical course averages 3 days from the time of onset until the patient sees the physician. In most instances the patients experience fever and chills. On physical examination, the most common finding was a pharyngeal purulent exudate. Which of the following types of inflammation did these patients most likely have?

A. Granulomatous inflammation
B. Abscess formation
C. Acute inflammation
D. Resolution of inflammation
E. Chronic inflammation



6. A 48-year-old man has experienced midabdominal pain for several weeks. He is afebrile. There is mild upper abdominal tenderness on palpation, and bowel sounds are present. Stool is positive for occult blood. An upper gastrointestinal endoscopy is performed and biopsies are taken. Microscopic examination of a biopsy specimen of a duodenal lesion is shown. Which of the following terms best describes this lesion?

A. Abscess
B. Caseating granuloma
C. Chronic inflammation
D. Purulent exudate
E. Ulceration

7. One month after an appendectomy, a 34-year-old man palpates a small nodule beneath the skin at the site of the healed right lower quadrant incision. The nodule is excised, and microscopic examination shows macrophages, collagen, a few small lymphocytes, and multinucleated giant cells. Polarizable, refractile material is seen in the nodule. Which of the following complications of the surgery best accounts for these findings?
   A. Chronic inflammation
   B. Abscess formation
   C. Edema
   D. Ulceration
   E. Suture granuloma
8. The most characteristic feature of granulomatous inflammation is
   A. Aggregations of epithelioid macrophages
   B. Necrosis
   C. Multinucleated giant cell
   D. Langhans'giant cells
   E. Eosinophils
9. All of the following are associated with chronic inflammation EXCEPT:
   A. Preponderance of lymphocytes, macrophages, and plasma cells
   B. Arises de novo or from acute inflammation
   C. Lasts months to years
   D. Macrophages aggregated through macrophage inhibiton factor (MIF)
   E. Usually related to higher levels of tissue damage than seen in acute inflammation
10. Which one of the following mediators of acute inflammation is chemotactic?
   A. Serotonin
   B. Complement 3a
   C. Prostaglandins
   D. Lysosomal proteases
   E. Oxygen radicals
11. An exudate is best described as:
   A. A fluid low in protein content with a specific gravity of less than 1.020
   B. A fluid high in protein content with a specific gravity of less than 1.020
   C. A fluid high in protein content with a specific gravity of more than 1.020
   D. A fluid with large amounts of neutrophils and necrotic material
   E. Happens before transudate due to increased vascular permeability
12. The vascular response in acute inflammation
   A. Is initially a vasodilatation followed by vasoconstriction
   B. Involves the emigration of neutrophils into the interestial tissues before other inflammatory cells
   C. Allows leukocytes to move from the periphery to the center of the column of blood in a vessel
   D. Results in a transudate

E. Results in leukocyte pavementing the adventitia of blood vessels

13. All of following are true of neutrophils EXCEPT:
    A. Phagocytize bacteria
    B. Contain oxidases and proteases
    C. First on scene after tissue injury
    D. Release histamine
    E. Life expectancy in tissue is around 4 days
14. A 56-year-old man has had increasing difficulty breathing for the past week. On physical examination he is afebrile. Auscultation of the chest reveals diminished breath soundsand dullness to percussion bilaterally. There is 2+ pitting edema present to the level of the thighs. A chest radiograph reveals bilateral pleural effusions. Which of the following laboratory test findings is he most likely to have?
    A. Hypoalbuminemia
    B. Glucosuria
    C. Neutrophilia
    D. Anemia
    E. Hypernatremia
15. A 58-year-old woman had a cough with fever for 3 days. A chest radiograph revealed infiltrates in the right lower lobe. A sputum culture grew Streptococcus pneumoniae. The clearance of these organisms from the lung parenchyma would be most effectively accomplished through generation of which of the following substances by the major inflammatory cell type responding to this infection?
    A. Platelet activating factor
    B. Prostaglandins
    C. Kallikrein
    D. Hydrogen peroxide
    E. Leukotriene
16. In an experiment, a lung tissue preparation is exposed to Mycobacterium tuberculosis organisms. Over the next week, it is observed that granulomas form in the lung. Within the granuloma there are inflammatory cells expressing class II MHC antigens. These cells elaborate cytokines that promote fibroblastic production of collagen within the granulomas. From which of the following peripheral blood leukocytes are these cells bearing class II antigen most likely to be derived?
    A. Neutrophils
    B. B cells
    C. Monocytes
    D. NK cells
    E. Basophils
17. A 52-year-old woman with no major medical problems takes a long airplane flight. Upon arrival at Sydney's Kingsford Smith airport in New South Wales following the flight from Los Angeles, California, she cannot put her shoes back on. Which of the following is the most likely explana-

tion for this phenomenon?

A. Activation of Hageman factor has led to bradykinin production
B. A lot of drinks were served in the first class section.
C. She didn't take any aspirin
D. A cellulitis developed in her legs
E. Venous hydrostatic pressure was increased

18. A 45-year-old man has had a fever and dry cough for 3 days, and now has difficulty breathing and a cough productive of sputum. On physical examination his temperature is 38.5 ℃ . Diffuse rales are auscultated over the lower lung fields. A chest radiograph shows a right pleural effusion. A right thoracentesis is performed. The fluid obtained has a cloudy appearance with a cell count showing 5500 leukocytes per microliter, 98% of which are neutrophils. Which of the following terms best describes his pleural process?

A. Serous inflammation
B. Purulent inflammation
C. Fibrinous inflammation
D. Chronic inflammation
E. Granulomatous inflammation

19. A 19-year-old man incurs a stab wound to the chest. The wound is treated in the emergency room. Two months later there is a firm, 3×2cm nodular mass with intact overlying epithelium in the region of the wound. On examination the scar is firm, but not tender, with no erythema. This mass is excised and microscopically shows fibroblasts with abundant collagen. Which of the following mechanisms has most likely produced this series of events?

A. Hypertrophic scar (keloid) formation
B. Development of a fibrosarcoma
C. Poor wound healing from diabetes mellitus
D. Foreign body response from suturing
E. Staphyloccocal wound infection

20. A 24-year-old primigravida is late in the second trimester of pregnancy. She experiences the sudden onset of some cramping lower abdominal pain. This is immediately followed by passage of some clear fluid from the vagina along with a foul-smelling discharge. The fetus is stillborn two days later. Examination of the placenta demonstrates extensive neutrophilic infiltrates in the chorion and amnion. Which of the following organisms is most likely to be responsible for these findings?

A. Mycobacterium tuberculosis
B. Herpes simplex virus
C. Escherichia coli
D. Treponema pallidum
E. Toxoplasma gondii

21. A 20-year-old woman sustains an injury to her right calf in a mountain biking ac-

cident near Moab. On physical examination she has a 5cm long laceration on the right lateral aspect of her lower leg. This wound is closed with sutures. Wound healing proceeds over the following week. Which of the following factors will be most likely to aid and not inhibit wound healing in this patient?

A. Secondary wound infection
B. Presence of sutures
C. Poor tissue perfusion
D. Hypoalbuminemia
E. Corticosteroid therapy

22. A 65-year-old woman has had a fever for the past 1 day. On physical examination her temperature is 39℃ and blood pressure is 90/50 mm Hg with heart rate of 106/minute. Laboratory studies include a CBC with WBC count of 12,220/microliter and WBC differential count of 78 neutrophils, 8 eosinophils and basophilic cells, 11 lymphs, and 3 monos. A blood culture is positive for Escherichia coli. Her central line pressure falls markedly. She goes into hypovolemic shock as a result of the widespread inappropriate release of a chemical mediator derived from macrophages. She develops multiple organ failure. Which of the following mediators is most likely to produce these findings?

A. Nitric oxide
B. Bradykinin
C. Histamine
D. Prostacyclin
E. Complement C3a

23. A 9-year-old girl sustains a small 0.5 cm long laceration to her right index finger while playing “Queen of Swords” with a letter opener. Which of the following substances, upon contact with injured vascular basement membrane activates both the coagulation sequence and the kinin system as an initial response to this injury?

A. Thromboxane
B. Plasmin
C. Hageman factor
D. Platelet activating factor
E. Histamine

24. A 31-year-old woman has a laparotomy performed for removal of an ovarian cyst. She recovers uneventfully, with no complications. At the time of surgery, a 12 cm long midline abdominal incision was made. When you advise this patient of the outcome of the healing process, you indicate that nearly all of the tensile strength to be obtained in wound healing in the skin will most likely be achieved within which of the following time periods?

A. One week
B. One month
C. Three months
D. Six months

E. One year

25. In a clinical study, patients undergoing laparoscopic cholecystectomy are followed to document the post-surgical wound healing process. The small incisions are closed with sutures. Over the 4 weeks following surgery, the wounds are observed to regain tensile strength and there is re-epithelialization. Of the following substances, which is most likely found to function intracellularly in cells involved in this wound healing process?
    A. Tyrosine kinase
    B. Laminin
    C. Fibronectin
    D. Hyaluronic acid
    E. Collagen
26. A 45-year-old man incurs a burn injury to his hands and arms while working on a propane furnace. Over the next 3 weeks, the burned skin heals without need for skin grafting. The most critical factor in determining whether the skin in the region of the burn will regenerate following such an injury is the presence of which of the following factors?
    A. Good cardiac output
    B. Minimal edema and erythema
    C. Underlying connective tissue
    D. Skin appendages
    E. Granulation tissue
27. A 55-year-old man with a history of ischemic heart disease has worsening congestive heart failure. He has noted increasing dyspnea and orthopnea for the past 2 months. On physical examination there is dullness to percussion at the lung bases. A chest x-ray shows bilateral pleural effusions. A left thoracentesis is performed, and 500 ml of fluid is obtained. Which of the following characteristics of this fluid would most likely indicate that it is a transudate?
    A. Cloudy appearance
    B. < 3 lymphocytes/microliter
    C. High protein content
    D. The presence of fibrin
    E. Large size of the effusion
28. In an experiment, surgical wound sites are observed following suturing. An ingrowth of new capillaries is observed to occur within the first week. A substance elaborated by macrophages is found at the wound site stimulating this capillary proliferation. Which of the following substances is most likely to have this function?
    A. Platelet-derived growth factor
    B. Phospholipase C-gamma
    C. Fibroblast growth factor
    D. Fibronectin
    E. Epidermal growth factor
29. In an experiment, lymphatic channels are observed in normal soft tissue prepara-

tions. Staphylococcus aureus organisms are innoculated into the tissues and the immunologic response is observed over the next 24 hours. Which of the following functions is most likely to be served by these lymphatics to produce a specific immune response to these organisms?

A. Carry lymphocytes to peripheral tissue sites
B. Transport antigen presenting cells
C. Remove extravascular tissue fluid
D. Serve as a route for dissemination of infection
E. Provide an emigration point for neutrophils

30. A 43-year-old woman has had nausea with vomiting persisting for the past 5 weeks. On physical examination there are no abnormal findings. She undergoes an upper GI endoscopy and gastric biopsies are obtained. The microscopic appearance of these biopsies shows mucosal infiltration by lymphocytes, macrophages, and plasma cells. Which of the following conditions is this woman most likely to have?

A. Staphylococcus aureus septicemia
B. Ingestion of chili peppers
C. Diabetes mellitus
D. Tuberculosis
E. Infection with Helicobacter pylori

31. In an experiment, Enterobacter cloacae organisms are added to a solution containing leukocytes. Engulfment and phagocytosis of the microbes is observed. Next a substance is added which enhances engulfment. Which of the following substances is most likely to produce this effect?

A. Complement C3b
B. Glutathione peroxidase
C. Immunoglobulin M
D. P-selectin
E. NADPH oxidase

32. A 72-year-old woman did not get a “flu” shot in the fall as recommended for older persons. In the wintertime, she became ill, as did many people in her community, with a respiratory illness that lasted for several weeks. During this illness, she had a fever with a non-productive cough, mild chest pain, myalgias, and headache. What was her chest radiograph most likely to have shown during this illness?

A. 6cm left hilar mass
B. Bilateral interstitial infiltrates
C. Marked hilar lymphadenopathy
D. Left upper lobe consolidation
E. Extensive pleural thickening with pleural effusions

33. Within minutes following a bee sting, a 37-year-old man develops marked respiratory stridor with dyspnea and wheezing. He also develops swelling and erythema in his arms and legs. An injec-

tion of epinephrine helps to reverse these events and he recovers within minutes. Which of the following chemical mediators is most important in the pathogenesis of this man's condition?

A. Bradykinin
B. Complement C5a
C. Nitric oxide
D. Histamine
E. Tumor necrosis factor

34. A 45-year-old man has been working hard all day long carrying loads of bricks to build a wall. He can use a non-steroidal anti-inflammatory drug such as ibuprofen in order to try and reduce which of the following processes in his arms?

A. Thrombosis
B. Liquefactive necrosis
C. Pain
D. Fibrinolysis
E. Scar formation

35. A 15-year-old girl has had episodes of sneezing with watery eyes and runny nose for the past 2 weeks. On physical examination she has red, swollen nasal mucosal surfaces. She has had similar episodes each spring and summer when the amount of ragweed pollen in the air is high. Her symptoms are most likely to be mediated by the release of which of the following chemical mediators?

A. Complement C3b
B. Platelet activating factor (PAF)
C. Tumor necrosis factor (TNF)
D. Histamine
E. Immunoglobulin G

36. A 55-year-old man has a history of hypercholesterolemia with coronary artery disease and suffered a myocardial infarction 2 years ago. He now presents with crushing substernal chest pain. Which of the following laboratory tests will be most useful in this patient?

A. Increased white blood cell count
B. Elevated sedimentation rate
C. Decreased serum complement
D. Increased serum creatine kinase
E. Decreased platelet count

37. A 36-year-old woman had bilateral silicone breast implants placed two years ago. After that time, she noted increased firmness with slight deformity of the breast on the left. The implants were removed, and there was evidence for leakage of the implant contents on the left. Which of the following cell types is most likely to be characteristic of the inflammatory response in this situation?

A. Mast cell
B. Eosinophil
C. Plasma cell
D. Neutrophil
E. Giant cell

38. A 45-year-old woman has had a chronic,

non-productive cough for 3 months, together with intermittent fever. She has a chestradiograph that reveals multiple small parenchymal nodules along with hilar and cervical lymphadenopathy. A cervical lymph node biopsy is performed. Microscopic examination of the biopsy shows noncaseating granulomatous inflammation. Cultures for bacterial, fungal, and mycobacterial organisms are negative. Which of the following chemical mediators is most important in the development of this inflammation?

A. Interferon gamma
B. Bradykinin
C. Complement C5a
D. Histamine
E. Prostaglandin E_2

39. A 19-year-old woman is in the early second trimester at 16 weeks gestation with her first pregnancy. On a routine prenatal visit, the nurse cannot detect fetal heart tones and an ultrasound reveals no fetal cardiac activity or fetal movement. The mother has noted no recent illness or other factors to account for this fetal demise. She is given a pharmacologic agent to induce abortion. By which of the following mechanisms is this induction most likely to be accomplished?

A. Leukotriene-induced neutrophil aggregation
B. Complement-induced hemolysis
C. Histamine-induced edema
D. Prostaglandin-induced muscular contraction
E. Endotoxin-induced disseminated intravascular coagulation

40. A 17-year-old driving a pickup truck is involved in a collision. He incurs blunt force abdominal trauma. In response to this injury, cells in tissues of the abdomen are stimulated to enter the G_1 phase of the cell cycle from the G_0 phase. Which of the following cell types is most likely to remain in G_0 following this injury?

A. Smooth muscle
B. Skeletal muscle
C. Endothelium
D. Fibroblast
E. Hepatocyte

41. A 37-year-old man has had nausea and vomiting for 5 weeks. He experienced an episode of hematemesis yesterday. On physical examination he has no abnormal findings. Upper GI endoscopy is performed, and there is a 1.5 cm diameter lesion in the gastric antrum which appears to be an area with loss of the epithelial surface. These findings are most typical for which of the following pathologic processes?

A. Abscess
B. Serositis

C. Granuloma
D. Gangrene
E. Ulcer

42. An inflammatory process that has continued for 3 months includes the transformation of tissue macrophages to epithelioid cells. There are also lymphocytes present. Over time, fibroblasts lay down collagen as the focus of inflammation heals. These events are most likely to occur as an inflammatory response to which of the following infectious agents?
A. Pseudomonas aeruginosa
B. Mycobacterium leprae
C. Cytomegalovirus
D. Giardia lamblia
E. Treponema pallidum

43. A small sliver of wood becomes embedded in the finger of a 25-year-old man. He does not remove it, and over the next 3 days the area around the sliver becomes red, swollen, and tender. Neutrophils migrate into the injured tissue. Expression of which of the following substances on endothelial cells is most instrumental in promoting this inflammatory reaction?
A. Interferon gamma
B. Hageman factor
C. E-selectin
D. Lysozyme
E. Prostacyclin

44. A 35-year-old woman has been taking acetylsalicylic acid (aspirin) for arthritis for the past 4 years. Her joint pain is temporarily reduced via this therapy. However, radiographs of her knees reveal continuing joint destruction with loss of articular cartilage and joint space narrowing. This pain reduction is most likely to be the result of diminishing which of the following inflammatory responses?
A. Neutrophil chemotaxis by leukotriene B_4
B. Prostaglandin-mediated vasodilation
C. Fever resulting from interleukin-1 release
D. Pain resulting from bradykinin generation
E. Anticoagulation by Hageman factor inhibition

45. A 90-year-old woman has developed a fever and cough over the past 2 days. Staphylococcus aureus is cultured from her sputum. She receives a course of antibiotic therapy. Two weeks later she no longer has a productive cough, but she still has a fever. A chest radiograph reveals a 3 cm rounded density in the right lower lobe whose liquefied contents form a central air-fluid level. There are no surrounding infiltrates. Which of the following is the best description for this outcome of her pneumonia?
A. Hypertrophic scar

B. Bronchogenic carcinoma
C. Regeneration
D. Abscess formation
E. Progression to chronic inflammation

46. An episode of marked chest pain lasting 4 hours brings a 51-year-old man to the emergency room. He is found to have an elevated serum creatine kinase. An angiogram reveals complete blockage of the left circumflex artery 2 cm from its origin. Which of the following substances would you most expect to be elaborated around the region of tissue damage in the first 3 days as an initial response which will increase myocardial function and promote healing?
A. Histamine
B. Vascular endothelial growth factor
C. Complement C3b
D. Leukotriene B_4
E. Immunogloblulin G

47. A 25-year-old man experienced painful urination for 4 days following spring break. Urethritis is suspected, and Neisseria gonorrheae is cultured. Numerous neutrophils are present in a smear of the exudate from the penile urethra. These neutrophils are most likely to have undergone diapedesis to reach the organisms as a consequence of release of which of the following chemical mediators?
A. Complement C5a
B. Prostaglandin
C. Hageman factor
D. Bradykinin
E. Histamine

48. A 42-year-old man has developed marked right lower quadrant abdominal pain over the past 1 day. On physical examination there is rebound tenderness to palpation over the right lower quadrant. Laparoscopic surgery is performed, and the appendix is swollen, erythematous, and partly covered by yellowish exudate. It is resected, and a microscopic section shows infiltration with numerous neutrophils. The pain experienced by this patient is predominantly the result of expression of which of the following two chemical mediators?
A. Prostaglandin and bradykinin
B. Interleukin-1 and tumor necrosis factor
C. Histamine and serotonin
D. Complement C3b and IgG
E. Leukotriene and HPETE

答案

中文题

1.A	2.C	3.D	4.C	5.C	6.C	7.E	8.D	9.D	10.C
11.E	12.C	13.D	14.E	15.E	16.E	17.D	18.D	19.A	20.B
21.C	22.C	23.E	24.D	25.E	26.E	27.A	28.E	29.E	30.E
31.E	32.D	33.C	34.D	35.C	36.C	37.C	38.D	39.E	40.B
41.C	42.E	43.E	44.E	45.D	46.A	47.A	48.D	49.C	50.C
51.C	52.B	53.C	54.D	55.C	56.E	57.D	58.B	59.C	60.C
61.C	62.E	63.D	64.C	65.E	66.D	67.E	68.E	69.D	70.E
71.D	72.E	73.D	74.B	75.E	76.C	77.E	78.C	79.E	80.D
81.D	82.E	83.E	84.C	85.C	86.A	87.D	88.E	89.B	90.A
91.E	92.D	93.B	94.B	95.C	96.D	97.A	98.D	99.B	100.D
101.D	102.C	103.E	104.D	105.C	106.E	107.D	108.B	109.B	110.B
111.B	112.E	113.C	114.C	115.A	116.B	117.B	118.C	119.C	120.C
121.D	122.D	123.A	124.D	125.C	126.D	127.B	128.A	129.A	130.E
131.A	132.B	133.C	134.C	135.C	136.C	137.C	138.D	139.A	140.C
141.D	142.C	143.A	144.E	145.B	146.D	147.E	148.B	149.E	150.A
151.B	152.B	153.B	154.D	155.B	156.D	157.B	158.B	159.D	160.C

英文题

1.A	2.B	3.D	4.B	5.C	6.E	7.E	8.A	9.E	10.C
11.C	12.B	13.D	14.A	15.D	16.C	17.E	18.B	19.A	20.C
21.B	22.A	23.C	24.C	25.A	26.D	27.B	28.C	29.B	30.E
31.A	32.B	33.D	34.C	35.D	36.D	37.E	38.A	39.D	40.B
41.E	42.B	43.C	44.B	45.D	46.B	47.A	48.A		

4 肿瘤

中文题

1. 不符合肿瘤性增生的是
   A. 增生的细胞具有异型性
   B. 需致瘤因素的持续存在
   C. 细胞代谢旺盛
   D. 可不形成肿块
   E. 相对无止境生长
2. 肿瘤性增生区别于炎症性增生的特点是
   A. 增生组织不分化不成熟
   B. 血管增生
   C. 纤维组织增生
   D. 器官的实质细胞增生
   E. 炎症细胞浸润
3. 大肠癌的肿瘤标志是
   A. CEA
   B. AFP
   C. CA19-9
   D. 酸性磷酸酶
   E. 碱性磷酸酶
4. 前列腺癌的肿瘤标志是
   A. CEA
   B. AFP
   C. CA19-9
   D. 酸性磷酸酶
   E. 碱性磷酸酶
5. 骨肉瘤的肿瘤标志是
   A. CEA
   B. AFP
   C. CA19-9
   D. 酸性磷酸酶
   E. 碱性磷酸酶
6. 肝细胞癌的肿瘤标志是
   A. CEA
   B. AFP
   C. CA19-9
   D. 酸性磷酸酶
   E. 碱性磷酸酶
7. 胰腺癌的肿瘤标志是
   A. CEA
   B. AFP
   C. CA19-9
   D. 酸性磷酸酶
   E. 碱性磷酸酶
8. 符合肿瘤防治现状的描述是
   A. 恶性肿瘤早期症状明显
   B. 大多数中晚期癌的治疗效果满意
   C. 早期发现、早期诊断、早期治疗具有重要意义
   D. 公民对癌的防治意识已大幅度提高
   E. 癌症的发病率不断下降
9. 符合癌的肉眼形态是
   A. 包膜完整
   B. 火山口状
   C. 分叶状
   D. 质地软
   E. 灰红色
10. 弥漫型胃癌肉眼形态呈
    A. 息肉状

B. 溃疡状
C. 弥漫浸润
D. 结节状
E. 囊状

11. 诊断恶性肿瘤的组织学依据主要是
A. 细胞浆呈嗜碱性
B. 细胞核大
C. 细胞异型性明显
D. 核仁明显
E. 可见核分裂像

12. 女性，26 岁，右乳外上象限有一直径约 2cm 大小的圆形肿块，边界清楚，表面光滑，可移动，不疼。此肿块最可能是
A. 乳腺癌
B. 乳腺增生症
C. 乳腺纤维瘤
D. 乳腺纤维腺瘤
E. 乳腺炎症性肿块

13. 恶性肿瘤异型性主要表现在
A. 肿瘤细胞胞浆嗜碱性
B. 可见核分裂
C. 肿瘤细胞核的多形性
D. 肿瘤细胞分泌减少
E. 组织结构紊乱

14. 良性肿瘤异型性主要表现在
A. 细胞形态不一
B. 核分裂像增多
C. 组织结构紊乱
D. 细胞间变明显
E. 细胞分泌减少

15. 不符合肿瘤形态特点的描述的是
A. 体积巨大的肿瘤多为恶性
B. 可发生多原发癌
C. 可形成微小癌
D. 可发生重复癌
E. 多为单发

16. 不符合肿瘤代谢特点的是
A. DNA、RNA 合成代谢旺盛
B. 蛋白质合成代谢旺盛
C. 糖代谢以无氧酵解为主
D. 可产生肿瘤蛋白
E. 肿瘤标志物仅见于肿瘤组织

17. 癌转移至淋巴结时，首先出现在
A. 边缘窦
B. 髓窦
C. 淋巴滤泡内
D. 副皮质区
E. 淋巴结门部

18. 淋巴结转移性癌的确切诊断依据是
A. 淋巴结肿大
B. 淋巴结质地变硬
C. 淋巴结疼痛
D. 淋巴结内出现上皮细胞
E. 淋巴结内出现癌巢

19. 在恶性肿瘤患者的周围血中查见恶性肿瘤细胞，说明该患者
A. 已发生了血道转移
B. 已发生了淋巴道转移
C. 已是恶性肿瘤晚期
D. 并发白血病
E. 有可能发生血道转移

20. 左锁骨上淋巴结转移性鳞癌的原发部位最可能是
A. 甲状腺

B. 胃
C. 食管
D. 乳腺
E. 肝

21. 男性，60 岁，上腹痛半年，因上消化道大出血，急诊入院。手术标本胃窦部粘膜可见一个巨大的火山口状溃疡。切面质硬，磁白色，侵入肌层。肝内可见数个小结节，手术切除一同送病理，支持胃癌肝转移的病理变化是
A. 小结节由肝细胞构成
B. 小结节由印戒细胞癌构成
C. 小结节由假小叶构成
D. 小结节由鳞癌细胞构成
E. 小结节由毛细血管构成

22. 肿瘤血道转移的确切依据是
A. 肿瘤细胞侵入静脉
B. 肿瘤细胞侵入动脉
C. 肿瘤细胞出现在血液中
D. 肿瘤细胞栓塞于远隔器官
E. 在远隔器官形成与原发瘤同一组织学类型的肿瘤

23. 胃内两个癌结节，支持胃内癌转移最可靠的依据是
A. 两个癌结节不相连
B. 两个癌结节组织学类型不同
C. 血管、淋巴管内可见癌栓
D. 两个癌结节组织学类型相同
E. 两个癌结节均发生在粘膜层

24. 不符合癌转移的特点是
A. 可发生淋巴道转移
B. 可发生血道转移
C. 可发生种植性转移
D. 可发生跳跃转移
E. 不发生逆行转移

25. 不易发生癌转移的器官是
A. 脑
B. 骨
C. 肺
D. 肝
E. 心

26. 恶性肿瘤最本质的表现是
A. 浸润性生长
B. 生长迅速
C. 出现转移
D. 异型性明显
E. 有出血坏死

27. 良性肿瘤的最显著的特点是
A. 异型性小
B. 生长缓慢
C. 不浸润周围组织
D. 出血坏死少见
E. 不转移

28. 良性肿瘤对机体的影响最主要决定于
A. 肿瘤的病程
B. 肿瘤的大小
C. 肿瘤组织的来源
D. 肿瘤发生的部位
E. 肿瘤出现继发性变化

29. 不符合良性肿瘤对机体的影响的描述是
A. 可发生癌变
B. 可引起内分泌紊乱
C. 可发生出血囊性变

D. 可浸润周围组织
E. 可发生转移

30. 副肿瘤综合征是指
A. 恶性肿瘤引起的发热
B. 恶性肿瘤引起的贫血
C. 恶性肿瘤引起的营养不良
D. 恶性肿瘤引起的内分泌紊乱
E. 恶性肿瘤引起的神经功能紊乱

31. 恶性肿瘤患者的主要死亡原因是
A. 继发大出血
B. 继发感染
C. 广泛转移
D. 疼痛
E. 营养不良

32. 不符合副肿瘤综合征的描述是
A. 可发生于肝癌
B. 可发生于肺癌
C. 可发生于胰腺癌
D. 可发生于结肠癌
E. 不发生于肉瘤

33. 癌前病变不包括
A. 大肠腺瘤
B. 甲状腺腺瘤
C. 膀胱乳头状瘤
D. 阴茎乳头状瘤
E. 家族性腺瘤性息肉病

34. 原位癌的主要特征是
A. 发生于粘膜、表皮和腺体
B. 癌细胞占据上皮全层，但基底膜仍完整
C. 可以治愈
D. 上皮内出现异型性的细胞
E. 胃粘膜内癌是一种原位癌

35. 不符合非典型增生的描述是
A. 属癌前病变
B. 增生的细胞大小形态不一、核深染、核分裂增多
C. 轻度和中度非典型增生是不可逆转的
D. 重度非典型增生需积极治疗
E. 可发展为原位癌

36. 不属于癌前病变的是
A. 粘膜白斑
B. 乳腺纤维腺瘤
C. 结肠腺瘤性息肉病
D. 慢性萎缩性胃炎
E. 乳腺纤维囊性病

37. 符合原位癌的是
A. 早期子宫颈癌
B. 乳腺导管内癌
C. 大肠粘膜内癌
D. 早期食管癌
E. 胃粘膜内癌

38. 癌前病变中，需要积极治疗的病变是
A. 纤维组织增生
B. 上皮增生
C. 轻度非典型增生
D. 中度非典型增生
E. 重度非典型增生

39. 符合癌前病变的描述是
A. 一类较高癌变潜能的病变
B. 一类已发生癌变的病变
C. 一类必然癌变的病变
D. 一类需积极手术治疗的病变

E. 一类交界性病变

40. 上皮非典型增生是指

A. 上皮细胞单纯增生

B. 上皮细胞层次增多

C. 上皮细胞化生

D. 增生的上皮细胞具有异型性

E. 无癌变倾向

41. 非间叶组织来源的肿瘤是

A. 骨肉瘤

B. 血管肉瘤

C. 精原细胞瘤

D. 恶性纤维组织细胞瘤

E. 平滑肌肉瘤

42. 癌肉瘤是

A. 既有癌又有肉瘤的肿瘤

B. 腺上皮发生的恶性肿瘤

C. 色素细胞发生的恶性肿瘤

D. 间叶组织发生的恶性肿瘤

E. 鳞状上皮发生的恶性肿瘤

43. 黑色素瘤

A. 既有癌又有肉瘤的肿瘤

B. 腺上皮发生的恶性肿瘤

C. 色素细胞发生的恶性肿瘤

D. 恶性间叶性肿瘤

E. 恶性上皮性肿瘤

44. 不属于恶性肿瘤的是

A. 黑色素瘤

B. 淋巴瘤

C. 尤文瘤

D. 多形性腺瘤

E. 骨髓瘤

45. 交界性肿瘤是指

A. 有两个胚层的组织构成的肿瘤

B. 介于良性与恶性肿瘤之间的肿瘤

C. 具有癌和肉瘤结构的肿瘤

D. 发生于表皮与真皮交界处的肿瘤

E. 既有鳞癌又有腺癌结构的肿瘤

46. 腺鳞癌是指

A. 伴有鳞状上皮化生的腺癌

B. 既有上皮成分，又有间质成分的肿瘤

C. 介于良性与恶性之间的肿瘤

D. 同时具有癌和肉瘤结构的肿瘤

E. 肿瘤中既有鳞状细胞癌成分，又有腺癌成分

47. 属于恶性肿瘤的是

A. 纤维瘤

B. 神经母细胞瘤

C. 骨瘤

D. 乳头状瘤

E. 腺瘤

48. 发生于儿童的恶性肿瘤是

A. 肾母细胞瘤

B. 黑色素瘤

C. 骨髓瘤

D. 鳞状细胞癌

E. 腺癌

49. 男性，70 岁，1 年来常有便血，肠镜检查，发现距肛门 8cm 的肠粘膜有一 2cm 大小的肿物，中心有溃疡。活检诊断为粘液腺癌，组织学改变是

A. 癌细胞形成腺腔

B. 癌细胞大小较一致

C. 癌细胞分泌大量粘液

D. 癌细胞周围间质丰富
E. 癌细胞周围血管丰富

50. 鳞状细胞癌的主要诊断依据是
A. 发生于鳞状上皮的覆盖部位
B. 呈浸润性生长
C. 癌细胞形成角化珠和细胞间桥
D. 肿瘤呈灰白色、质硬
E. 经淋巴道转移

51. 胃幽门部一般不会出现
A. 腺癌
B. 鳞状细胞癌
C. 硬癌
D. 胶样癌
E. 髓样癌

52. 不符合乳头状瘤特征是
A. 可发生于移行上皮
B. 可发生于皮肤表面
C. 可发生于胃肠道粘膜
D. 呈外生性生长
E. 极少发生恶变

53. 有关移行细胞癌的描述，错误的是
A. III 级预后最好
B. 好发于膀胱和肾盂等处
C. 组织学上可有不同程度的移行上皮的特点
D. 呈浸润性生长
E. 可由乳头状瘤恶变而来

54. 诊断腺癌的组织学依据是
A. 肿瘤细胞多形性明显
B. 出现病理核分裂
C. 侵犯血管
D. 细胞异型性明显，呈腺样排列
E. 核深染

55. 不符合腺瘤特征的是
A. 具有分泌功能
B. 形成导管和小叶结构
C. 细胞异型性不明显
D. 浸润性生长
E. 形成腺样结构

56. 不符合髓样癌特征的是
A. 来源于腺上皮
B. 癌巢呈实体性，无腺腔形成
C. 癌细胞多
D. 可见淋巴细胞浸润
E. 纤维间质丰富

57. 不符合乳腺纤维腺瘤特征的是
A. 是一种良性肿瘤
B. 好发于青年女性
C. 切面灰白、质韧，呈编织状
D. 包膜完整
E. 由分化良好的纤维组织构成

58. 男性，60 岁，1 年来进行性吞咽困难，日渐加重。胃镜检查，食管下段可见一溃疡型肿物，质地较硬，溃疡周边隆起，底部凹凸不平。活检的病理诊断最可能是
A. 食管小细胞癌
B. 食管腺癌
C. 食管鳞状细胞癌
D. Barrett 食管
E. 食管平滑肌瘤

59. 女性，58 岁，停经 6 年后，阴道不规则出血 2 个月。妇科检查，子宫体积增大，如孕两个月大。刮宫取子宫内膜

送病理，诊断为中分化子宫内膜腺癌。不符合其组织学改变的是

A. 腺体密集、背靠背

B. 出现鳞状上皮

C. 腺上皮细胞核深染

D. 腺上皮细胞排列紊乱

E. 可见病理核分裂

60. 女性，40 岁，偶然发现右乳有一肿块，黄豆大，质硬，手术切除，其组织学改变符合导管内癌的是

A. 导管上皮乳头状增生

B. 导管上皮与纤维组织混合增生

C. 导管扩张

D. 导管内细胞异型性明显

E. 导管上皮及小叶上皮增生

61. 男性，60 岁，无痛性血尿半年余。膀胱镜检查，膀胱三角区可见一乳头状肿物，有蒂。活检报告为乳头状瘤。符合此诊断的病变是

A. 可见鳞状细胞团

B. 可见腺体

C. 乳头被覆分化好的移行细胞

D. 乳头被覆分化好的腺上皮

E. 乳头被覆分化好的鳞状上皮

62. 不符合皮肤基底细胞癌的描述是

A. 好发于面部

B. 多见于老年人

C. 可形成溃疡

D. 与紫外线照射有关

E. 易发生转移

63. 不符合硬癌特征的是

A. 发生于腺上皮

B. 癌细胞无腺腔形成

C. 癌巢小而少，间质纤维组织多

D. 可弥漫性浸润

E. 分化较好

64. 腮腺多形性腺瘤的特点是

A. 由上皮和肌上皮细胞发生

B. 由腺上皮发生

C. 由肌上皮发生

D. 包膜完整

E. 切除后不易复发

65. 不符合毛细血管瘤特征的是

A. 儿童常见

B. 多为先天性发生

C. 多见于皮肤

D. 成年后可停止发展，甚至可以自行消退

E. 膨胀性生长

66. 不符合横纹肌肉瘤特征的是

A. 多见于婴幼儿和青少年

B. 肿瘤由不同分化阶段的横纹肌母细胞构成

C. 切面呈鱼肉状

D. 恶性度较高，主要发生血道转移

E. 常发生于横纹肌的部位

67. 不符合骨瘤特点的是

A. 良性肿瘤

B. 好发于头面骨及颌骨

C. 由成熟的骨质组成

D. 骨母细胞异型性明显

E. 可引起压迫症状

68. 有关瘤样纤维组织增生的描述，错误的是

A. 是一种良性肿瘤
B. 呈浸润性生长
C. 可复发
D. 不转移
E. 肿瘤无包膜

69. 不符合平滑肌瘤特点的是
A. 好发于子宫、胃肠道
B. 肿瘤有包膜
C. 切面灰白、质韧，呈编织状
D. 不见核分裂像
E. 不转移

70. 有关脂肪肉瘤的描述，错误的是
A. 多由脂肪瘤恶变而来
B. 好发于大腿、腹膜后等部位
C. 多见于 40 岁以上的患者
D. 肿瘤呈结节状或分叶状
E. 常有假包膜

71. 平滑肌肉瘤的诊断依据中，最重要的是
A. 肿瘤细胞丰富
B. 肿瘤粘液变性
C. 核分裂像的多少
D. 细胞的多形性明显
E. 肿瘤出血坏死

72. 女性，50 岁，发现背部肿瘤 5 年，生长缓慢。检查时，肿瘤直径约 5cm 大小，可活动，境界清楚。手术切除，肿瘤呈淡黄色，分叶状，包膜完整，质地较软。此结节最可能是
A. 纤维瘤
B. 血管瘤
C. 平滑肌瘤
D. 脂肪瘤
E. 滑膜瘤

73. 男性，30 岁，右胫骨上端病理性骨折，局部被囊性肿物破坏。手术后，病理诊断为骨巨细胞瘤。符合此诊断的描述是
A. 良性肿瘤
B. 巨细胞是肿瘤的主要细胞
C. 间质细胞是肿瘤的主要细胞
D. 瘤细胞形成骨组织
E. 瘤细胞形成软骨组织

74. 女性，47 岁，因子宫多发性结节，子宫切除。病理检查，符合平滑肌肉瘤的病变是
A. 肿瘤细胞分化良好
B. 肿瘤细胞密集
C. 肿瘤细胞异型性明显
D. 肿瘤细胞核分裂像多见
E. 肿瘤细胞粘液变性

75. 骨肉瘤的主要诊断依据是
A. 好发于青少年
B. 血道转移
C. 发生于长骨干骺端
D. 恶性肿瘤细胞产生骨组织
E. 发生病理性骨折

76. 葡萄状肉瘤属于
A. 血管肉瘤
B. 脂肪肉瘤
C. 纤维肉瘤
D. 横纹肌肉瘤
E. 纤维肉瘤

77. 不符合软骨肉瘤的描述是
A. 青少年多见
B. 肿瘤呈分叶状

C. 浸润性生长
D. 肿瘤细胞产生软骨基质
E. 可发生出血坏死

78. 不符合恶性纤维组织细胞瘤的描述是
A. 可发生与内脏器官
B. 由原始间叶组织发生
C. 可见泡沫细胞和多核巨细胞
D. 可见异型性平滑肌细胞
E. 恶性度高

79. 纤维肉瘤的主要诊断依据是
A. 发生于深部组织
B. 鱼肉状
C. 异型性明显的纤维母细胞
D. 出血坏死
E. 假包膜形成

80. 混合型血管瘤的组织学特点是
A. 由毛细血管构成
B. 由血窦构成
C. 由小静脉构成
D. 由小动脉构成
E. 由毛细血管和血窦构成

81. 不符合精原细胞瘤的特征的是
A. 恶性
B. 多发于婴儿
C. 起源于原始生殖细胞
D. 由较一致的大圆形细胞组成
E. 间质可见淋巴细胞侵润

82. 不符合皮样囊肿的特征的描述是
A. 属于良性肿瘤
B. 多见于皮肤
C. 囊内充满油脂和毛发
D. 囊壁内被覆鳞状上皮
E. 囊壁内可见皮肤附属器

83. 符合成熟型畸胎瘤的特征的描述是
A. 良性肿瘤
B. 不发生于胸腔
C. 多为实性肿物
D. 不形成甲状腺组织
E. 不形成脑组织

84. 卵巢成熟性畸胎瘤多呈
A. 分叶状
B. 乳头状
C. 弥漫浸润
D. 结节状
E. 囊状

85. 有关畸胎瘤的描述，错误的是
A. 好发于卵巢、睾丸
B. 由两个胚层以上的组织构成
C. 未成熟型可见原始神经组织
D. 多呈囊性，囊内充满油脂、毛发
E. 肿瘤起源于生发上皮

86. 关于皮肤黑痣的描述，错误的是
A. 皮内痣易恶变
B. 来源于黑色素细胞
C. 可恶变为黑色素瘤
D. 溃破、发炎是恶变的象征
E. 痣细胞分泌色素

87. 有关黑色素瘤的描述，错误的是
A. 可由黑色素痣恶变而来
B. 恶性度较高，易发生血道转移
C. 均可见色素颗粒
D. 好发于足底、外阴、肛周的皮肤
E. 瘤细胞核仁清楚、核分裂像多见

88. 女性，22 岁，发现右下腹包块 3 个月，

手术发现右卵巢被破坏，被一囊性肿物所取代，囊内充满脂性物质、毛发，囊壁可见牙齿。应诊断为

A. 无性细胞瘤

B. 皮样囊肿

C. 错构瘤

D. 迷离瘤

E. 畸胎瘤

89. 不属于致癌因子的是

A. 硒

B. 苯胺

C. 亚硝胺

D. 二甲基氨基偶氮苯

E. 3,4—苯并芘

90. 与肺癌发生关系密切的是

A. 3，4-苯并芘

B. 黄曲霉毒素

C. 石棉纤维

D. 紫外线

E. 乙萘胺

91. 与膀胱癌发生关系密切的是

A. 3，4-苯并芘

B. 黄曲霉毒素

C. 石棉纤维

D. 紫外线

E. 乙萘胺

92. 与肝癌发生关系密切的是

A. 3，4-苯并芘

B. 黄曲霉毒素

C. 石棉纤维

D. 紫外线

E. 乙萘胺

93. 不属于致癌物质的是

A. 砷

B. 金

C. 铬

D. 镍

E. 镉

94. 不符合亚硝胺致癌特点的描述是

A. 致癌性强

B. 致癌谱广

C. 自然界广泛存在

D. 其形成与亚硝酸盐有关

E. 其形成与二级胺有关

95. 不符合 3，4-苯并芘致癌特点的描述是

A. 致癌性强

B. 致癌谱广

C. 自然界广泛存在

D. 存在于烟草的烟雾中

E. 具有直接致癌作用

96. 不符合化学致癌的描述是

A. 间接致癌物多见

B. 直接致癌物少见

C. 某些致癌物需要其他物质的协同作用

D. 激发剂是致癌物质

E. 促进剂是致癌物质

97. 与病毒无关的肿瘤是

A. 肝癌

B. 鼻咽癌

C. Burkitt 淋巴瘤

D. 子宫颈癌

E. 胃癌

98. 与肝癌关系密切的病毒是

A. HBV
B. HSV
C. EBV
D. HPV
E. HTLV

99. 艾滋病患者常见的恶性肿瘤是
A. 鼻咽癌
B. Kaposi 肉瘤
C. 食管癌
D. 胃癌
E. 乳腺癌

100. 与 HPV 感染关系密切的肿瘤是
A. 鼻咽癌
B. 食管癌
C. 子宫颈癌
D. 胃癌
E. 乳腺癌

101. 与 HCV 感染关系密切的肿瘤是
A. 鼻咽癌
B. Kaposi 肉瘤
C. 食管癌
D. 肝癌
E. 乳腺癌

102. 与 EBV 感染关系密切的肿瘤是
A. 鼻咽癌
B. Kaposi 肉瘤
C. 食管癌
D. 胃癌
E. 乳腺癌

103. 与 HTLV 感染关系密切的肿瘤是
A. 鼻咽癌
B. Kaposi 肉瘤
C. 食管癌
D. 胃癌
E. T 细胞白血病

104. 与 EBV 感染关系密切的肿瘤是
A. Burkitt 淋巴瘤
B. Kaposi 肉瘤
C. 食管癌
D. 胃癌
E. T 细胞白血病

105. 男性，20 岁，轻度腹泻，伴粘液便 4 年。肠镜检查，结肠粘膜可见密布的米粒到黄豆大的息肉。病变符合
A. 结肠腺瘤
B. 结肠癌
C. 家族性腺瘤性息肉病
D. 神经纤维瘤病
E. 家族性大肠癌

106. 女性，5 岁，双眼均患有视网膜母细胞瘤。生殖细胞发生改变的基因是
A. APC 基因
B. WT-1 基因
C. P53 基因
D. NF-1 基因
E. Rb 基因

107. APC 基因突变可引起
A. 神经纤维瘤病
B. 结肠腺瘤性息肉病
C. 视网膜母细胞瘤
D. 肾母细胞瘤
E. 家族性乳腺癌

108. WT-1 基因突变可引起
A. 神经纤维瘤病

B. 结肠腺瘤性息肉病
C. 视网膜母细胞瘤
D. 肾母细胞瘤
E. 家族性乳腺癌

109. BRCA-1 基因突变可引起
A. 神经纤维瘤病
B. 结肠腺瘤性息肉病
C. 视网膜母细胞瘤
D. 肾母细胞瘤
E. 家族性乳腺癌

110. NF-1 基因突变可引起
A. 神经纤维瘤病
B. 结肠腺瘤性息肉病
C. 视网膜母细胞瘤
D. 肾母细胞瘤
E. 家族性乳腺癌

111. 不符合遗传因素在肿瘤发生中的作用的描述是
A. 在遗传性肿瘤中起决定作用
B. 大多数恶性肿瘤的发生是环境与遗传共同作用的结果
C. 着色性干皮病属于遗传性肿瘤
D. 遗传性肿瘤多由肿瘤抑制基因丢失或灭活引起
E. 遗传性肿瘤也可由 DNA 修复基因突变引起

112. 不属于肿瘤易感性遗传的疾病是
A. 着色性干皮病
B. Bloom 综合征
C. Fanconi 综合征
D. 毛细血管扩张性共济失调
E. 视网膜母细胞瘤

113. 与雌激素水平过高关系密切的肿瘤是
A. 肝癌
B. 肺癌
C. 卵巢癌
D. 乳腺癌
E. 子宫颈癌

114. 有关肿瘤免疫的描述，错误的是
A. 体液免疫起主要作用
B. 癌组织内大量淋巴细胞浸润是预后较好的指标
C. 恶性肿瘤可以破坏宿主的免疫功能
D. 免疫功能低下的患者容易发生恶性肿瘤
E. 艾滋病患者恶性肿瘤的发生率升高

115. 与雄激素关系密切的肿瘤是
A. 精原细胞瘤
B. 阴茎癌
C. 前列腺癌
D. 胚胎性癌
E. 内胚窦瘤

116. 不符合肿瘤免疫的描述是
A. 艾滋病患者更易发生恶性肿瘤
B. 免疫抑制治疗可诱发恶性肿瘤
C. Wilms 瘤的发生与免疫系统不健全有关
D. 老年人肿瘤的发生与免疫系统功能下降有关
E. 先天性免疫缺陷病恶性肿瘤的发病率升高

117. 不符合肿瘤抗原的描述是
A. 肿瘤相关抗原既存在于肿瘤细胞，又存在于某些正常细胞

B. 肿瘤胚胎抗原在正常机体有微量存在
C. 肿瘤特异抗原可以用于肿瘤的诊断
D. 肿瘤胚胎抗原在胚胎发育中曾多量出现
E. 肿瘤胚胎抗原可作为肿瘤标记物

118. 不符合肿瘤免疫逃避的描述是
A. 肿瘤细胞有较强的抗原性
B. 肿瘤细胞 MHC 抗原减少
C. 肿瘤患者的免疫系统受到抑制
D. 肿瘤细胞产生免疫抑制因子
E. 极少数癌症患者的肿瘤自行消退与免疫功能增强有关

119. 与营养因素有关的恶性肿瘤有
A. 乳腺癌
B. 子宫颈癌
C. 食管癌
D. 胃癌
E. 大肠癌

120. 在肿瘤免疫当中重要的效应细胞是
A. 中性粒细胞
B. 浆细胞
C. 巨噬细胞
D. 嗜酸粒细胞
E. B 细胞

121. 与皮肤癌发生关系密切的是
A. 3，4-苯并芘
B. 黄曲霉毒素
C. 石棉纤维
D. 紫外线
E. 乙萘胺

122. 与肺间皮瘤发生关系密切的是
A. 3，4-苯并芘
B. 黄曲霉毒素
C. 石棉纤维
D. 紫外线
E. 乙萘胺

123. 可引起白血病的因素是
A. 3，4-苯并芘
B. 黄曲霉毒素
C. 石棉纤维
D. X 线
E. 乙萘胺

124. 着色性干皮病患者容易发生皮肤癌是由于
A. 3，4-苯并芘
B. 黄曲霉毒素
C. 石棉纤维
D. 紫外线引起的 DNA 损伤
E. 乙萘胺

125. 与血吸虫卵慢性刺激有关的癌是
A. 肺癌
B. 胆管上皮癌
C. 膀胱癌
D. 大肠癌
E. 胃癌

126. 不符合电离辐射致癌的描述是
A. 导致染色体断裂
B. 导致染色体异位
C. 引起点突变
D. 肿瘤发生潜伏期短
E. 激活癌基因、灭活肿瘤抑制基因

127. 不符合紫外线照射致癌的描述是
A. 可引起黑色素瘤

B. 可引起鳞状细胞癌
C. 可引起基底细胞癌
D. 可引起蕈样霉菌病
E. 可导致嘧啶二聚体形成

128. 不符合电离辐射致癌的描述是
A. 致癌谱广
B. 与辐射剂量有关
C. 易防护
D. 放射治疗不会引起肿瘤发生
E. 与化学及生物致癌因素有叠加作用

129. 有关抗癌基因与肿瘤关系的描述，错误的是
A. APC 基因——结肠腺瘤性息肉病
B. NF-1 基因——神经纤维瘤病
C. Rb 基因——视网膜母细胞瘤
D. WT-1 基因——肾母细胞瘤
E. P53 基因——肺癌

130. 有关原癌基因的描述，错误的是
A. 正常细胞的固有基因
B. 对细胞的生长起抑制作用
C. 在正常细胞中以非激活的方式存在
D. 环境和遗传因素均可激活原癌基因
E. 其激活可由点突变、染色体异位、插入、扩增引起

131. 有关肿瘤的发生发展的描述，错误的是
A. 长时间的过程
B. 多个因素参与
C. 多基因突变
D. 多阶段逐步演化
E. 恶性肿瘤不可逆转

132. 不符合肿瘤抑制基因的特点有
A. 正常细胞的固有基因
B. 促进细胞的生长
C. Rb 基因和 P53 基因是两种常见的抗癌基因
D. 抗癌基因突变可能引起细胞的肿瘤性转化
E. 肿瘤的发生需要两个等位基因的突变

133. 不符合原癌基因蛋白质产物的描述是
A. 生长因子
B. 生长因子受体
C. 信号转导蛋白质
D. 核调节蛋白质
E. 激素受体

134. 不符合恶性肿瘤发生发展的描述是
A. 肿瘤是一种基因病
B. 单一基因改变即可引起细胞恶性转化
C. 累及癌基因的改变
D. 累及肿瘤抑制基因的改变
E. 累及 DNA 修复基因的改变

135. 不符合恶性肿瘤发生发展的基因改变的描述是
A. 不同阶段均有相应的基因改变
B. 多种基因受累
C. 多种基因的积累效应为主
D. 多种基因的序贯效应为主
E. 肿瘤抑制基因在生殖细胞的突变是某些遗传性肿瘤的基础

136. 不符合 P53 基因的描述是
A. P53 基因的突变发生于结肠癌、肺癌等实体肿瘤

B. 野生型 P53 蛋白抑制细胞生长
C. 突变型 P53 蛋白促进细胞增生
D. P53 蛋白存在于细胞浆内
E. P53 基因位于 17 号染色体

137. 不符合良性肿瘤的是
A. 生长缓慢
B. 不浸润周围组织
C. 可发生恶变
D. 术后很少复发
E. 不发生转移

138. 诊断恶性肿瘤的主要组织学依据是
A. 结构紊乱
B. 组织坏死
C. 异型性明显
D. 血管丰富
E. 浸润周围组织

139. 区别良性肿瘤与恶性肿瘤的主要组织学依据是
A. 结构紊乱
B. 细胞形态不一
C. 间质内炎细胞浸润
D. 细胞核多形性
E. 纤维组织增生

140. 区别乳腺内良性肿瘤与恶性肿瘤的主要临床依据是
A. 肿瘤的体积
B. 肿瘤的部位
C. 肿瘤的硬度
D. 肿瘤的活动度
E. 肿瘤的浸润情况

141. 区别子宫平滑肌瘤和平滑肌肉瘤的主要病理学依据是
A. 结构紊乱
B. 细胞形态不一
C. 体积大小
D. 核分裂像的多少
E. 细胞核多形性

142. 区别脂肪瘤与脂肪肉瘤的主要病理学依据是
A. 肿瘤的体积
B. 肿瘤的分化情况
C. 肿瘤的硬度
D. 肿瘤的活动度
E. 肿瘤的部位

143. 大肠腺瘤癌变的主要肉眼形态改变是
A. 腺瘤的体积
B. 腺瘤的部位
C. 腺瘤溃疡形成
D. 腺瘤呈分叶状
E. 腺瘤呈绒毛状

144. 区别肺癌与肺炎性假瘤的主要组织学依据是
A. 结构紊乱
B. 肺泡上皮增生
C. 间质内炎细胞浸润
D. 细胞核多形性
E. 纤维组织增生

145. 支持肉瘤的主要诊断依据是
A. 假包膜形成
B. 淋巴道转移
C. 呈鱼肉状
D. 血道转移
E. 肿瘤细胞弥漫散在，与间质混杂

146. 支持癌的主要诊断依据是

A. 恶性肿瘤细胞弥散分布
B. 恶性肿瘤细胞浸润周围组织
C. 坏死灶形成
D. 大片出血
E. 恶性肿瘤细胞形成巢片

147. 符合肉瘤的主要组织学依据是
A. 恶性肿瘤细胞间可见网状纤维
B. 恶性肿瘤细胞浸润周围组织
C. 恶性肿瘤细胞有内分泌功能
D. 淋巴结内出现转移灶
E. 恶性肿瘤细胞形成巢片

148. 确定癌的主要依据是
A. 老年人
B. 浸润性生长
C. 癌巢形成
D. 肿瘤组织的异型性明显
E. 核分裂像多见

149. 不符合癌的分级的描述是
A. 高分化鳞癌形成角化珠
B. 低分化鳞癌恶性度高
C. 高分化腺癌恶性度低
D. 实性癌不形成腺体
E. 未分化癌是腺癌的一种

150. 不符合恶性肿瘤分期的描述是
A. T 表示肿瘤的大小
B. L 表示淋巴结转移情况
C. M 表示脏器转移情况
D. $T_1L_0M_0$ 表示肿瘤处于早期
E. $T_4L_3M_1$ 表示肿瘤处于中期

151. 制定胃癌的治疗方案常依据
A. 脱落细胞学诊断
B. 细针穿刺病理诊断
C. 冰冻切片病理诊断
D. 活体组织检查及手术标本的病理诊断
E. 影像学诊断

152. 不符合腺癌的分级的描述是
A. 高分化腺癌形成腺体
B. 中分化腺癌部分形成腺体
C. 低分化腺癌恶性度低
D. 实性癌不形成腺体
E. 髓样癌是腺癌的一种

153. atypia
A. 间变
B. 多形性
C. 异型性
D. 分化
E. 成熟

154. leiomyosarcoma
A. 骨肉瘤
B. 脂肪肉瘤
C. 横纹肌肉瘤
D. 平滑肌肉瘤
E. 软骨肉瘤

155. teracoma
A. 错构瘤
B. 迷离瘤
C. 假瘤
D. 结核瘤
E. 畸胎瘤

156. adenocarcinoma
A. 鳞癌
B. 移行细胞癌
C. 腺癌

D. 粘液癌
E. 硬癌

157. melanoma
A. 淋巴瘤
B. 骨髓瘤
C. 尤文瘤
D. 黑色素瘤
E. 迷离瘤

158. metastasis
A. 侵袭
B. 转移
C. 扩散
D. 间变
E. 进展

159. cachexia
A. 恶性度
B. 恶病质
C. 恶性转化
D. 恶变
E. 癌变


160. oncogene
A. 原癌基因
B. 病毒癌基因
C. 癌基因
D. 凋亡基因
E. 抗癌基因

英文题

1. Workers in the dye industry are most prone to which of the following tumors?
A. Aarcinoma of the lung
B. Squamous cell carcinoma of the skin
C. Basal cell carcinoma of the skin
D. Carcinoma of the stomach
E. Carcinoma of the urinary bladder

2. Which of the following tumors has been associated most strongly with a possible viral etiology?
A. Carcinoma of the lung
B. Burkitt's lymphoma
C. Carcinoma of the esophagus
D. Follicular lymphoma
E. Fibrosarcoma

3. A 72-year-old female has been feeling fatigued for weeks. She has had a 10kg weight loss in the past couple of months. She has some nausea, but no vomiting, hemoptysis, or hematemesis. She does not have a cough. She is afebrile. The most likely explanation for these findings, from the gross appearance of the liver seen here, is the presence of:



A. Hepatic adenomas
B. Metastatic adenocarcinoma
C. Hepatocellular carcinoma
D. This tumor should express AFP oncofetal protein

E. Micronodular cirrhosis

4. Which one of the following statements is most appropriate for the lesion shown here?



A. This tumor is always well-differentiated
B. The growth pattern for the tumor is local invasion
C. Metastasis may be exist when the tumor is identified
D. The etiology of the tumor is related to specific virus
E. None of above



5. A 40-year-old female has noted some intermenstrual blooding and felt pelvic discomfort. A pelvic examination is performed and the cervical lesion seen here is found and a hysterectomy performed. Which one of the following statements is most appropriate based on the appearance of this lesion?

A. This lesion is the result of smoking
B. This is a benign lesion that will not recur
C. This is a malignant lesion that will be lethal
D. This kind of tumor is not related to HPV
E. Syphilis is the main cause of the lesion

6. A 68-year-old male is found to have occult blood in the stool on physicial examination. A colonoscopy is performed, and the lesion seen here was found and resected. Which of the following statements is most appropriate for this patient?



A. His children have a high risk for a similar lesion
B. This is a benign lesion that will not recur
C. This lesion is the result of smoking
D. A sexually transmitted disease led to this lesion
E. Metastases from this lesion may be present

7. Which of the following is the most common growth pattern for transitional cell carcinoma?

A. Ulcerative
B. Odular

C. Leather like
D. Cystic
E. Papillary

8. Basal cell carcinoma is most common in which location in the elderly?
A. Hands
B. Face
C. Feet
D. Liver
E. Trunk

9. The most common cause of death from cancer in China is
A. Stomach cancer
B. Liver cancer
C. Lung cancer
D. Breast cancer
E. Colon cancer

10. Which one of the following items is most helpful in distinguishing poorly differentiated carcinoma from sarcoma?
A. The growth pattern
B. The manner of spread
C. The age of patient
D. The location of the tumor
E. Reticulum fiber stain

11. Seeding metastasis is most common in which of the following malignancies?
A. Melanoma
B. Gastric signet-ring carcinoma
C. Breast carcinoma
D. Osteosarcoma
E. Squamous carcinoma of penis

12. Which one the following of statements concerning the total cell cycle time in tumors is most appropriate when it is compared with normal cells?
A. 3 times faster than that of normal cells
B. 2 times faster than that of normal cells
C. Equal or even longer than that of normal cells
D. 5 times faster than that of normal cells
E. Much slower than that of normal cells

13. Analysis of cancers which have metastasized reveals a mutation which results in decreased cell surface expression of E-cadherin. Which of the following genes is most likely mutated to produce these findings?
A. Beta-catenin
B. Cyclin D
C. BRCA-2
D. NF-1
E. Rb

14. Patients with long-standing Epstein-Barr virus infection have an increased risk for development of malignant neoplasms in adulthood. Which of the following neoplasms is most likely to be found in these patients?
A. Kaposi sarcoma
B. Lipoma
C. Cervical carcinoma
D. Nasopharyngeal carcinoma
E. Endometrial carcinoma

15. A previously healthy 67-year-old man has been feeling tired for 5 months. A complete physical examination is performed and stool guaiac is positive. A colonoscopy is performed, and a large, sessile 4.0 cm mass with surface ulceration is found in the cecum. A biopsy of this mass microscopically shows irregular glands with crowded, tall columnar cells having marked nuclear hyperchromatism. Which of the following statements about his condition is most accurate?
    A. He has hereditary non-polyposis colon cancer
    B. A p53 mutation is present in the neoplastic cells
    C. There is lack of apoptosis from a faulty Bcl-2 gene
    D. His relatives have hereditary adenomatous polyposis
    E. He has inherited a faulty Rb gene from one parent
16. Cells of urothelial carcinoma demonstrate a single mutation causing cellular inability to hydrolyze GTP, thus resulting in cellular transformation. Which of the following oncogenes is most likely implicated in this case?
    A. c-abl
    B. c-sis
    C. k-ras
    D. c-neu
    E. c-myc
17. It is known that some pre-existing conditions are found to precede development of malignant neoplasms. Which of the following conditions is most likely to be statistically unrelated to subsequent malignancy?
    A. Cervical squamous dysplasia
    B. Chronic alcoholism with hepatic cirrhosis
    C. Endometrial atypical hyperplasia
    D. Uterine leiomyoma
    E. Chronic ulcerative colitis
18. An unequivocal features of malignancy is
    A. Cellular pleomorphism
    B. Lymphocytic infiltration at the edge of the tumor
    C. Numerous mitotic figures
    D. Metastases
    E. Necrosis
19. A 62-year-old man with an 80 pack year history of smoking experiences an episode of hemoptysis. On physical examination his face shows puffiness, there is some pedal edema and bruises on the skin, and his blood pressure is 165/100mmHg. A chest radiograph reveals a 5cm right upper lobe lung mass. A fine needle aspiration of this mass shows small cell carcinoma. A bone scan shows no metastases. Immunohistochemical staining of the tumor cells is likely to be positive for which

of the following hormones?

A. Parathormone related peptide
B. Erythropoietin
C. ACTH
D. Insulin
E. Gastrin

20. A 25-year-old woman in excellent health found a 2 cm firm, rounded mass beneath the skin of her upper back. The overlying skin appears normal. The mass is palpable and soft. She has no symptom over many years. Which of the following neoplasms is she most likely to have?

A. Metastatic carcinoma
B. Melanoma
C. Leiomyoma
D. Lipoma
E. Fibrosarcoma

21. A 14-year-old boy has had lumps in the right neck for the past 5 months. On physical examination there is painless lymphadenopathy in the right cervical region. One of the lymph nodes is biopsied and on microscopic examination shows many lymphocytes that are large, with clumped chromatin and occasional mitoses. The characterization of this population of lymphocytes as a neoplasm is best accomplished by which of the following methods?

A. Immunohistochemical staining identifying leukocyte common antigen
B. Gene rearrangement studies that demonstrate clonality
C. A differential white blood count showing a lymphocytosis
D. Flow cytometry indicating a high S-phase
E. Demonstrating necrosis within the mass by light microscopy

22. A 58-year-old man dies after a year-long illness. At autopsy the liver contains multiple tumor masses from 2cm to 5cm that are mostly firm and tan, and these masses grossly exhibit umbilication with central necrosis. Which of the following statements would best characterize the significance of such findings?

A. There is multicentric origin of a benign neoplasm
B. The primary neoplasm is in the stomach
C. A carcinogen was the underlying cause for the neoplasm
D. The neoplasm is at an advanced stage
E. The neoplasm is high grade

23. A 17-year-old healthy girl has a 0.5 cm reddish, slightly raised nodule on the skin of the upper part of her chest found on a routine physical examination. She states that this lesion has been present for years and has not appreciably changed in size or color. Which of the following neoplasms is this nodule most likely to be?

A. Hemangioma
B. Melanoma
C. Rhabdosarcoma
D. Osteoma
E. Lymphoma

24. A 35-year-old woman has had a feeling of abdominal discomfort for the past 7 months. On pelvic examination, there is a right adnexal mass. An abdominal CT scan demonstrates a 6 cm cystic mass involving the right ovary with small areas of calcification. The uterus is normal in size. The right fallopian tube and ovary are removed surgically. Grossly, on sectioning the mass is filled with abundant hair and sebum. Microscopically, the mass has glandular spaces lined by columnar epithelium, squamous epithelium with hair follicles, cartilage, and dense connective tissue. Which of the following statements regarding this type of neoplasm is most appropriate?
A. A human papillomavirus infection preceded development of this mass
B. Primitive immature tissues resembling those from an embryo can usually be seen
C. Metastases are unlikely to be present.
D. A sarcomatous element is usually present
E. Such a mass is often seen to arise in the testis

25. Which of the following neoplasms is most likely to develop from viral oncogenesis?
A. Transtional cell carcinoma
B. Retinoblastoma
C. T-cell leukemia
D. Seminoma
E. Hepatic angiosarcoma

26. A 51-year-old man has experienced vague abdominal discomfort for the past 4 months. On physical examination he has no lymphadenopathy, and no abdominal masses or organomegaly can be palpated. Bowel sounds are present. An abdominal CT scan shows a 20cm retroperitoneal soft tissue mass obscuring the left psoas muscle. A stool specimen tested for occult blood is negative. Which of the following neoplasms is this man most likely to have?
A. Melanoma
B. Hamartoma
C. Adenocarcinoma
D. Lymphoma
E. Liposarcoma

27. Which of the following statement(s) is/are correct?
A. The incidence of malignancy tends to increase with age
B. Certain tumors tend to occur in certain age groups
C. Certain tumors tend to occur in certain locations

D. Hormones may inhibit or promote tumor growth
E. All of the above statements are correct

28. Which one of the following statements is not true about paraneoplastic syndromes?
A. They may represent the earliest manifestation of an occult neoplasm
B. They may be serious and even be lethal
C. Cushing syndrome is the most common endocrinopathy
D. They can only be seen in cancer patients, not in sarcoma patients
E. They can mimic metastatic disease

29. Which one of the following statements is not correct about tumor antigens?
A. Tumor associated antigens are present on tumor cells and also on some normal cells
B. Tumor antigens may be normal cellular proteins
C. The expression of oncofetal antigens in adults is only limited to tumors
D. CD20 is a cell type-specific differentiation antigen
E. All of above

30. Among the following viruses, which one is an oncogenic RNA virus?
A. HIV
B. HCV
C. HBV
D. HTLV
E. HPV

31. Which of the following is true in identifying the steps involved in chemical carcinogenesis?
A. Promoters are tumorigenic by themselves
B. The effect of promoters is reversible
C. Initiation alone is sufficient for tumor formation
D. Initiators are direct-acting compounds
E. None of above

32. Which of the following statements about tumor angiogenesis is correct?
A. Tumor blood vessels are similar to normal vasculature
B. Tumor angiogenesis is related to the decreased production of VEGF
C. VEGF can only be produced by tumor cells
D. Angiogenesis is induced from a very early stage of tumor growth
E. During angiogenesis anti-angiogenesis molecules can also be induced by tumor cells

33. Which of the following statements is true about tumor suppressor genes?
A. The function of these genes is to prevent tumor formation
B. These genes were discovered by studying common diseases
C. WT1 and VHL genes act in a manner similar to the RB gene

D. Loss of heterozygosity involving the P16 gene is the mechanism of retinoblastoma
E. None of above

34. Which of the following does not belonged to a defective DNA repair syndrome?
A. HNPCC
B. Xeroderma pigmentosum
C. Ataxiatelangectasia
D. Li-Fraumeni syndorme
E. Bloom syndrome

35. Which of the following features is true about metastasis?
A. All cancers can metastasize, In general the more rapidly growing
B. In general，the more rapidly growing，the greater the likelihood that cancer will metastasize
C. The probability of metastasis can be determined by pathologic examination of the primary tumor
D. Nodal enlargement in proximity to a cancer definitely means lymphatic spread
E. Sarcoma can only metastasis by hematogenous spread

36. Which of the following statements is true about cancer stem cells?
A. Cancer stem cells have been identified in almost all kinds of cancers
B. Cancer stem cells have a low rate of replication
C. Cancer stem cells constitute 20% of the cells in breast tumors
D. Cancer stem cells can be killed easily by chemotherapy
E. None of above

37. A benign neoplasm characterized by an abnormal amount of tissue indigenous to the area is best referred to as
A. Teratoma
B. Mixed tumor
C. Hamartoma
D. Embryonal tumor
E. Keloid

38. The characteristic appearance of a neoplasm depends on which of the following?
A. parenchyma
B. mesenchyma
C. metastasis
D. atypia
E. mitosis

39. A 53-year-old man has a palpable mass lesion on digital rectal examination. Biopsy was performed. Of the following microscopic findings, which is most likely to indicate that the neoplasm is malignant?
A. Pleomorphism
B. Increased nuclear/cytoplasmic ratio
C. Necrosis
D. Invasion
E. Atypia

40. A 55-year-old woman goes to her physician for a routine physical examination. A

4 cm diameter non-tender mass is palpated in her right breast. The mass appears fixed to the chest wall. Another 2 cm non-tender mass is palpable in the left axilla. A chest radiograph reveals multiple 0.5 to 2 cm nodules in both lungs. Which of the following classifications best rerpresents the stage of her disease?

A. T_1 N_1 M_0
B. T_1 N_0 M_1
C. T_2 N_1 M_0
D. T_3 N_0 M_1
E. T_4 N_1 M_1

41. Which of the following features is not true in describing carcinoma in situ?
A. Dysplastic changes involve the entire thickness of the epithelium
B. It is preinvasive neoplasm
C. It is intramucosal carcinoma
D. It is early stage carcinoma
E. It does not metastastasize

42. Which of the following features is true in describing malignant neoplasm?
A. The presence of giant cells indicate that a tumor is malignant
B. Lack of differentiation is a hallmark of malignant transformation
C. The presence of mitoses indicate that a tumor is malignant
D. The bigger the neoplasm the higher the tumor grade
E. Anaplastic tumors do not differ cytologically from normal counterparts

43. Which of the following conditions is a not preneoplastic disorder?
A. Chronic atrophic gastritis
B. Chronic ulcerative colitis
C. Leukoplakia of the oral cavity
D. Thyroid adenoma
E. Papilloma of the penis

44. Which of the following tumors is not malignant?
A. Mesothelioma
B. Seminoma
C. Osteoma
D. Wilm's tumor
E. Leukemia

45. The term border line tumor means:
A. Tumor consisting of both carcinoma and sarcoma
B. Tumor consisting of squamous carcinoma and adenocarcinoma
C. Tumor consisting of epithelial tissue and meschymal tissue
D. Tumor bordering between benign and malignant
E. Tumor bordering between hyperplasia and carcinoma

46. The patient survival rate with stage $T_4N_1M_1$ neoplam 5 years from the time of diagnosis is less than 50%, regardless of therapy. Which of the following clinical findings is most likely to be characteristic of this group of patients?

A. Cachexia
B. Cardiac murmur
C. Icterus
D. Tympany
E. Splenomegaly

47. Which of the following factors presents the greatest risk for development of malignant melanoma in the skin of hand?
A. Smoking tobacco
B. Asbestos exposure
C. Chemotherapy
D. Ultraviolet radiation
E. Allergy to latex

48. Angiogenesis in a neoplasm is best characterized as:
A. blood-borne spread of tumor cells
B. invasion of blood vessels by tumor cells
C. the ingrowth of new blood vessels into a growing tumor
D. a tumor arising from primitive blood forming tissue
E. central necrosis seen in rapidly growing tumors resulting from the tumor outgrowing its blood supply

49. Which of the following techniques used during partial colectomy for colon carcinoma can best aid the surgeon in determining whether the resection is adequate to reduce the probability of a recurrence?
A. Fine-needle aspiration
B. Serum carcinoembryonic antigen assay
C. Frozen section
D. Electron microscopy
E. Flow cytometry

50. Which of the following is a risk factor for hepatic angiosarcoma?
A. Arsenic
B. Asbestos
C. Benzene
D. Beryllium
E. Vinyl chloride

51. Which gene is most likely to have mutation in the familial breast carcinoma?
A. BCL2 (antiapoptosis gene)
B. BRCAl (DNA repair gene)
C. EGF (epidermal growth factor gene)
D. ERBB2 (growth factor receptor gene)
E. K-RAS (GTP-binding protein gene)

52. Which of the following best describes the mechanism of sporadic colonic adenomas evolving into an invasive carcinoma?
A. Protooncogenes can be activated by chromosomal translocation
B. Extensive regeneration of tissues increases the risk of cancer-causing mutations
C. Malignant transformation involves accumulation of mutations in protooncogenes and tumor suppressor genes in a step-wise fashion
D. Inherited defects in DNA repair increase the susceptibility to develop

cancer

E. Overexpression of growth factor receptor genes is associated with poor prognosis

53. Which of the following features of malignancies is related to its sensitivity to chemotherapeutic agents?

A. High growth fraction

B. Monoclonality

C. Poor vascularity

D. Prone to metastasis

E. Strong expression of tumor antigens

54. A 14-year-old girl in a family of three children has had skin nodules over the past 5 years. On physical examination, she is of appropriate height and weight. The skin lesions are 1- to 3-cm maculopapular nodules that are erythematous to brown-colored and have areas of ulceration. Biopsy of the skin lesions shows squamous cell carcinoma. The child has no history of recurrent infections, and her parents and other relatives are not affected. Which of the following mechanisms is most likely to produce neoplasia in her condition?

A. Infection with human papillomavirus

B. Inactivation of p53

C. Ingestion of food contaminated with Aspergillus flavus

D. Failure of nucleotide excision repair of DNA

E. Chromosomal translocation

55. A needle biopsy of the liver shows a mucin secreting adenocarcinoma. The most likely site of origin is?

A. Bladder

B. Cerebellum

C. Prostate

D. Colon

E. Lung

56. A 55-year-old woman was diagnosed with well-differentiated ductal carcinoma of right breast and there were three firm lymph nodes palpable in the right axilla. There was no family history of cancer. 6 months later, she died of multiple metastases in the lung, liver and brain. Which of the following molecular abnormalities is most likely to be found in this setting?

A. Inactivation of one copy of the BRCA1 gene in breast cancer cells

B. Amplification of the ERBB2 (HER2) gene in breast cancer cells

C. Deletion of one copy of the p53 gene in all of the patient's cells

D. Deletion of the RB locus in the patient's normal somatic cells

E. Fusion of BCR and C-ABL genes in the cancer cells

57. Which of the following investigational research procedures can most clearly distinguish whether an isolated 1cm pedunculated polyp represents hyperplasia of the

colonic mucosa or a tubular adenoma?

A. Immunoperoxidase staining for keratin
B. Immunoperoxidase staining for Ki-67
C. Flow cytometry to determine the frequency of cells in the S phase
D. Histochemical staining for mucin
E. Determination of clonality by pattern of X chromosome inactivation

58. Which of the following molecular abnormalities in cervical carcinoma is most likely related to infection with HPV-16?

A. Trapping of the RAS protein in a GTP-bound state
B. activation of the c-myc gene
C. Inability to repair DNA damage
D. Functional inactivation of the RB1 protein
E. Increased expression of epidermal growth factor receptor

59. Which of the following factors is most likely responsible for the lymph node metastases in breast carcinoma?

A. Increased laminin receptors on tumor cells
B. Presence of keratin in tumor cells
C. Diminished apoptosis of tumor cells
D. Tumor cell monoclonality
E. Lymphadenitis

60. A 45-year-old man has had hepatitis for 14 years, and HBsAg and anti-HCV are positive. He develops hepatocellular carcinoma. Which of the following best explains why this patient developed hepatocellular carcinoma?

A. The consistent integration of these viruses in the vicinity of protooncogenes
B. The ability of viral genes to inactivate RB and p53 expression
C. Viral induced injury to liver cells followed by extensive regeneration
D. The ability of these viruses to capture protooncogenes from the host DNA
E. The ability of these viruses to cause immunosuppression in the host

61. Which of the following maligancies does not increase in incidence in the setting of alcohol abuse?

A. Oropharyngeal
B. Laryngeal
C. Esophageal
D. Gastric
E. Hepatocellular

62. Which of the following carcinomas tends to spread via vascular invasion as opposed to lymphatic invasion?

A. lung carcinoma
B. Renal cell carcinoma
C. gastric carcinoma
D. colon carcinoma
E. nasopharyngeal carcinoma

63. In what organ has cytologic screening produced a marked reduction in mortality?

A. Breast
B. Lung
C. Pancreas
D. Uterine cervix
E. Stomach

64. The increase in size of a neoplasm depends on the doubling time which
A. is related to the growth fraction and rate of cell loss
B. is related to its clonality
C. is the same as the length of the cell cycle
D. is related to the length of the cell cycle
E. is related to the age of the patient

65. Which of the following is a neoplasm of mesenchymal origin?
A. leukemia
B. seminal cell tumor
C. fibrosarcoma
D. liposarcoma
E. all of the above

66. Patients with hereditary nonpolyposis colon cancer typically develop multiple lesions of the colon during middle age. Molecular analysis of the cells from these lesions shows changes in hPMSl, hPMS2, and hMLHl genes. Which of the following principles of carcinogenesis is best illustrated by this study?
A. Tumor initiators are mutagenic
B. Tumor promoters induce proliferation
C. Many oncogenes are activated by translocations
D. Inability to repair DNA is carcinogenic
E. Carcinogenesis is a multistep process

67. A healthy 23-year-old woman undergoes a routine physical examination. A discrete, firm, rubbery, movable mass is found in the left breast. She has no axillary lymphadenopathy. The skin overlying the breast and the nipple appear normal. Which of the following neoplasms is most likely to be present?
A. Fibroadenoma
B. Lipoma
C. Intraductal carcinoma
D. Malignant lymphoma
E. Leiomyoma

68. A 66-year-old man has developed a chronic cough over the past 2 months. There are no abnormal physical examination findings. A chest x-ray shows a 3 cm mass in the right upper lobe. After fine needle aspiration biopsy, the mass is resected. Gross and microscopic examination of the mass is performed and it is determined that a neoplasm is present. Which of the following findings regarding this mass is most likely to be irrelevant for determining further therapy and prognosis?
A. The size of the neoplasm
B. The differentiation of the neoplasm

C. Is the neoplasm invading the margin of resection
D. How much inflammation is present in the neoplasm
E. Is the neoplasm primary or metastatic

69. A 75-year-old healthy man is found to have a palpable firm nodule in the prostate gland by rectal examination. His serum prostate specific antigen (PSA) is 19 ng/mL. The microscopic appearance seen on biopsy of the prostate will probably be most consistent with which of the following conditions?
A. Chronic prostatitis
B. Adenocarcinoma
C. Infarction
D. Leiomyoma
E. Rhabdomyosarcoma

70. A 47-year-old man has noted a lump on his left shoulder that has enlarged over the past 4 months. On physical examination there is an enlarged, non-tender supraclavicular lymph node. A biopsy of the node is done and on microscopic examination there is a metastatic neoplasm. Which of the following is the most likely primary neoplasm?
A. Cerebral glioma
B. Adenocarcinoma of colon
C. Leiomyosarcoma of shoulder soft tissue
D. Fibrosarcoma of shoulder soft tissue
E. Liposarcoma of the retroperitoneum

71. A child is born with a single functional allele of a tumor suppressor gene. At the age of five the remaining normal allele is lost through a point mutation. As a result, the ability to inhibit cell cycle progression until the cell is ready to divide is lost. Which of the following neoplasms is most likely to arise via this mechanism?
A. Infiltrating ductal carcinoma of breast
B. Small cell anaplastic carcinoma of the lung
C. Cerebral astrocytoma metastasis
D. Retinoblastoma of eye
E. Chronic myeloid leukemia

72. A 35-year-old man with a family history of colon carcinoma undergoes a surveillance colonoscopy. It reveals hundreds of polyps in the colon, and two focal 0.5 cm ulcerated areas. A biopsy from one ulcer reveals irregularly shaped glands that have penetrated into the muscular layer. Which of the following molecular events is believed to occur very early in the evolution of his colonic disease process?
A. Activation of the WNT signaling pathway
B. Inability to hydrolyze GTP-bound RAS
C. Mutations in mismatch repair genes
D. Loss of heterozygosity affecting the p53 gene

E. Translocation of BCL2 from mitochondria to cytoplasm

73. A 63-year-old man sees his physician because of cough and hemoptysis. He has a 65 pack-year history of smoking. A chest CT scan shows a 5-cm right hilar mass. Bronchoscopy is performed, and lung biopsy specimens show small cell anaplastic lung carcinoma. His family history shows three first-degree maternal relatives who developed leukemia, sarcoma, and carcinoma before the age of 40 years. Which of the following genes is most likely to have undergone mutation to produce these findings?

A. APC
B. BCL2
C. ERBB2
D. K-RAS
E. p53

74. A 40-year-old man has generalized nontender enlargement of lymph node in the groin and armpit and hepatosplenomegaly. An inguinal lymph node biopsy specimen shows a malignant tumor of lymphoid cells. Immunoperoxidase staining of the tumor cells with antibody to BCL2 is positive in the lymphocytic cell nuclei. Which of the following mechanisms has most likely produced this lymphoma?

A. Increased tyrosine kinase activity
B. Reduced DNA repair
C. Gene amplifications
D. Lack of apoptosis
E. Loss of cell cycle inhibition

答案

中文题

1.B	2.A	3.A	4.D	5.E	6.B	7.C	8.C	9.B	10.C
11.C	12.D	13.C	14.C	15.A	16.E	17.A	18.E	19.E	20.C
21.B	22.E	23.D	24.E	25.E	26.C	27.E	28.D	29.E	30.D
31.C	32.E	33.B	34.B	35.C	36.B	37.B	38.E	39.A	40.D
41.C	42.A	43.C	44.D	45.B	46.E	47.B	48.E	49.C	50.C
51.B	52.E	53.A	54.D	55.B	56.E	57.E	58.C	59.B	60.D
61.C	62.E	63.E	64.A	65.E	66.E	67.D	68.A	69.B	70.A
71.C	72.D	73.C	74.D	75.D	76.D	77.A	78.D	79.C	80.E
81.B	82.B	83.A	84.E	85.E	86.A	87.C	88.E	89.A	90.A
91.E	92.B	93.B	94.C	95.E	96.E	97.E	98.A	99.B	100.C
101.D	102.A	103.E	104.A	105.C	106.E	107.B	108.D	109.E	110.A
111.C	112.E	113.D	114.A	115.C	116.C	117.C	118.A	119.E	120.C
121.D	122.C	123.D	124.C	125.D	126.D	127.D	128.D	129.E	130.B
131.E	132.B	133.E	134.B	135.D	136.D	137.B	138.C	139.D	140.E
141.D	142.B	143.C	144.D	145.E	146.E	147.A	148.C	149.E	150.E
151.D	152.C	153.C	154.D	155.E	156.C	157.D	158.B	159.B	160.C

英文题

1.E	2.B	3.B	4.C	5.C	6.B	7.E	8.B	9.C	10.E
11.B	12.C	13.A	14.D	15.B	16.C	17.D	18.D	19.C	20.D
21.B	22.D	23.A	24.C	25.C	26.E	27.E	28.D	29.C	30.D
31.B	32.E	33.C	34.D	35.B	36.B	37.C	38.A	39.D	40.E
41.C	42.B	43.D	44.C	45.D	46.A	47.D	48.C	49.C	50.E
51.B	52.C	53.A	54.D	55.D	56.B	57.E	58.D	59.A	60.C
61.D	62.B	63.D	64.A	65.E	66.D	67.A	68.D	69.B	70.B
71.D	72.A	73.E	74.D						

5 心血管疾病

中文题

1. 关于风湿病的叙述，不正确的是
   A. 是累及全身结缔组织的变态反应性疾病
   B. 以心脏病变对患者危害最大
   C. 风湿性心内膜炎引起的慢性心瓣膜病
   D. 风湿性关节炎常可导致关节畸形
   E. 皮下结节和环形红斑对临床诊断风湿病有帮助
2. 关于风湿病的叙述，错误的是
   A. 多见于寒冷地区
   B. 早期咽拭子培养，溶血性链球菌阳性率高达 70%～90%
   C. 抗生素的广泛应用，降低了风湿病的发病率
   D. 抗 O 滴度升高提示本病是由于溶血性链球菌直接作用所致
   E. 是一种结缔组织病
3. 对风湿病最具病理诊断意义的病变是
   A. 纤维素样变性
   B. 风湿小结
   C. 心肌变性
   D. 纤维素渗出
   E. 纤维组织增生
4. 风湿病常引起
   A. 细动脉玻璃样变
   B. 大、中动脉粥瘤
   C. 多发性栓塞性小脓肿
   D. 疣状心内膜炎
   E. 瓣膜穿孔
5. 不符合风湿病病变的描述是
   A. 胶原纤维的纤维素样变性
   B. Aschoff 小体形成
   C. 心肌细胞变性坏死
   D. 心外膜纤维素渗出
   E. Aschoff 细胞来源于心肌细胞
6. 风湿性病变中，对机体危害最大的是
   A. 反复发作的风湿性关节炎
   B. 反复发作的环形红斑
   C. 反复发作的风湿性心内膜炎
   D. 风湿性皮下结节
   E. 风湿性动脉炎
7. 不符合风湿病的病因和发病机制的描述是
   A. 抗生素的广泛应用，降低了风湿病的发病率
   B. 患者血清抗链球菌抗体滴度明显升高
   C. 多见于热带地区
   D. 发病前多先有急性扁桃体炎和咽峡炎
   E. 本病多始发于儿童
8. 不符合 Aschoff 小体的描述是
   A. 可见纤维素样变性
   B. 可见淋巴细胞
   C. Aschoff 细胞具有异型性
   D. 可发生纤维化
   E. Aschoff 细胞和 Anitschkow 细胞是

主要成分

9. 马氏斑（Mac Callum's Patch）位于
   A. 右心房
   B. 左心房
   C. 右心室
   D. 左心室
   E. 心外膜
10. 关于风湿性心内膜炎的描述，正确的是
   A. 瓣膜赘生物由血小板纤维素组成
   B. 瓣膜赘生物易脱落
   C. 受累瓣膜容易发生穿孔
   D. 受累瓣膜以主动脉瓣最多见
   E. 瓣膜赘生物内有细菌
11. 风湿病常见的联合瓣膜损害是
   A. 二尖瓣和肺动脉瓣
   B. 二尖瓣和三尖瓣
   C. 二尖瓣和主动脉瓣
   D. 主动脉瓣和肺动脉瓣
   E. 主动脉瓣和三尖瓣
12. 风湿病引起急性心功能不全的最常见原因是
   A. 疣状心内膜炎
   B. 风湿性心肌炎
   C. 绒毛心
   D. 缩窄性心包炎
   E. 马氏斑
13. 有关风湿性心脏病的描述，错误的是
   A. 常引起二尖瓣增厚、粘连
   B. 常引起肺动脉瓣增厚、粘连
   C. 常引起主动脉瓣增厚、粘连
   D. 可引起心包粘连
   E. 可引起心肌间质内小瘢痕形成
14. 关于慢性风湿性瓣膜病的描述，错误的是
   A. 瓣膜硬化
   B. 瓣膜粘连
   C. 腱索增粗
   D. 瓣膜卷曲
   E. 瓣膜断裂
15. 风湿性心内膜炎疣状赘生物的成分是
   A. 血小板和纤维素
   B. 纤维组织
   C. 细菌团和炎细胞
   D. 肉芽组织
   E. 弹力纤维
16. 风湿性关节炎不引起
   A. 关节积液
   B. 关节红肿
   C. 关节周围风湿小结形成
   D. 关节软骨破坏
   E. 关节疼痛
17. 风湿性心外膜炎引起
   A. 球形心
   B. 靴形心
   C. 梨形心
   D. 绒毛心
   E. 虎斑心
18. 风湿性动脉炎表现为
   A. 细动脉玻璃样变
   B. 大动脉粥瘤形成
   C. 中等动脉纤维素样坏死
   D. 小动脉纤维素样坏死
   E. 小动脉肌层肥厚
19. 女性，13 岁，一月前患扁桃腺炎，经治

疗痊愈。1 周来患者发热，关节肿胀。心尖区闻及舒张期 II～III 级杂音、心包摩擦音。血白细胞升高，血沉加快。抗 O 滴度升高。符合患者心脏疾病的描述是

A. 病毒性心肌炎
B. 细菌性心肌炎
C. 寄生虫性心肌炎
D. 风湿性心肌炎
E. 心肌病

20. 女性，13 岁，一月前患扁桃腺炎，经治疗痊愈。1 周来患者发热，关节肿胀。心尖区闻及舒张期 II～III 级杂音、心包摩擦音。血白细胞升高，血沉加快。抗 O 滴度升高。符合患者疾病的描述是

A. 关节病变属纤维素性炎
B. 心外膜病变为浆液性纤维素性炎
C. 心肌病
D. 本病与细菌直接感染有关
E. 心内膜形成较大赘生物

21. 女性，13 岁，一月前患扁桃腺炎，经治疗痊愈。1 周来患者发热，关节肿胀。心尖区闻及舒张期 II～III 级杂音、心包摩擦音。血白细胞升高，血沉加快。抗 O 滴度升高。符合患者心脏疾病的描述是

A. 亚急性细菌性心内膜炎
B. 二尖瓣狭窄
C. 二尖瓣关闭不全
D. 急性细菌性心内膜炎
E. 赘疣性血栓性心内膜炎

22. 慢性风湿性瓣膜病常见的联合瓣膜损害是

A. 二尖瓣、肺动脉瓣
B. 二尖瓣和三尖瓣
C. 二尖瓣和主动脉瓣
D. 主动脉瓣和肺动脉瓣
E. 主动脉瓣和三尖瓣

23. 急性风湿病最常见的致死原因是

A. 风湿性关节炎
B. 风湿性心脏病
C. 风湿性脑病变
D. 风湿性动脉炎
E. 风湿性肺炎

24. 不符合风湿病病变的描述是

A. 皮肤环形红斑有助于风湿病的诊断
B. 皮下结节有助于风湿病的诊断
C. 小舞蹈症由椎体外系统受累引起
D. Aschoff 小体仅出现于心肌内
E. 皮下结节内可见纤维素样坏死

25. 亚急性细菌性心内膜炎的最常见的病原体是

A. 金黄色葡萄球菌
B. 草绿色链球菌
C. β 溶血性链球菌
D. 大肠杆菌
E. 绿脓杆菌

26. 不符合亚急性细菌性心内膜炎的是

A. 由致病力强的细菌引起
B. 发生于已有病变的心瓣膜
C. 瓣膜赘生物较大，污秽，易脱落
D. 多引起非感染性梗死
E. 皮肤可见 Osler 结节

27. 亚急性细菌性心内膜炎不引起
A. 慢性瓣膜病
B. 局灶性肾小球肾炎
C. 多发性栓塞性小脓肿
D. 紫癜
E. 脾肿大

28. 女性，14 岁，室间隔缺损患者。拔牙后 1 月来，出现发热（37℃～38℃），食欲下降，消瘦、贫血。查体：心前区闻及杂音，心脏轻度增大，皮肤有散在出血点，甲下线状出血。蛋白尿（+）。符合该患者疾病的描述是
A. 急性细菌性心内膜炎
B. 亚急性细菌性心内膜炎
C. 赘疣性血栓性心内膜炎
D. 疣状心内膜炎
E. Libman-Sacks 血栓性心内膜炎

29. 女性，14 岁，室间隔缺损患者。拔牙后 1 月来，出现发热（37℃～38℃），食欲下降，消瘦、贫血。查体：心前区闻及杂音，心脏轻度增大，皮肤有散在出血点，甲下线状出血。蛋白尿（+）。符合该患者肾脏疾病的描述是
A. 弥漫性毛细血管增生性肾小球肾炎
B. 新月体性肾小球肾炎
C. 膜性肾小球肾炎
D. 局灶性肾小球肾炎
E. 轻微病变性肾小球肾炎

30. 感染性心内膜炎最常见的病原体是
A. 细菌
B. 霉菌
C. 病毒
D. 衣原体
E. 立克次体

31. 不符合亚急性细菌性心内膜炎的的描述是
A. 赘生物呈息肉样
B. 灰黄、污秽
C. 质脆、易脱落
D. 内含细菌菌落
E. 大量纤维组织增生

32. 不符合亚急性细菌性心内膜炎的的描述是
A. 引起脑梗死
B. 引起肾梗死
C. 引起脾梗死
D. 引起肺梗死
E. 引起肝梗死

33. 急性细菌性心内膜炎不引起
A. 高热
B. 心脏杂音
C. 疣状心内膜炎
D. 心内膜溃疡、穿孔
E. 多发性栓塞

34. 急性细菌性心内膜炎的瓣膜赘生物中，不具有的成分是
A. 大量中性白细胞
B. 坏死组织
C. 细菌菌团
D. 肉芽组织
E. 肉芽肿

35. 急性细菌性心内膜炎不引起
A. 瓣膜破裂
B. 慢性瓣膜病

C. 败血症
D. 细动脉壁纤维素样坏死
E. 血管化脓菌栓塞

36. 不符合急性细菌性心内膜炎的是
A. 瓣膜赘生物体积大、污秽、易脱落
B. 常伴有败血症
C. 发生于已有病变的心瓣膜
D. 可引起乳头肌断裂
E. 可引起心瓣膜穿孔，溃疡形成

37. 女性，26 岁，既往体健。正常产后 1 周来，阴道分泌物增多，伴有臭味，近 3 天来高热、寒战、神志昏迷。白细胞 2 万/mm^3，肝、脾肿大，心脏左心增大，心前区可闻及收缩期杂音。皮肤有多形淤斑，有的可见脓点。符合该患者疾病的描述是
A. 毒血症
B. 细菌血症
C. 败血症
D. 脓毒败血症
E. 病毒血症

38. 女性，26 岁，既往体健。正常产后 1 周来，阴道分泌物增多，伴有臭味，近 3 天来高热、寒战、神志昏迷。白细胞 2 万/mm^3，肝、脾肿大，心脏左心增大，心前区可闻及收缩期杂音。皮肤有多形淤斑，有的可见脓点。不符合该患者心内膜病变的描述是
A. 心内膜赘生物体积大、污秽、松脆
B. 心内膜赘生物内大量细菌
C. 心内膜赘生物由血小板和纤维素构成
D. 可引起瓣膜穿孔
E. 原本健康的瓣膜受累

39. 不符合非感染性内膜炎的描述是
A. 可见于慢性消耗性疾病
B. 可见于系统性红斑狼疮
C. 赘生物内可见血小板和纤维素
D. 赘生物内可有纤维素样坏死
E. 赘生物内可有细菌团

40. 不符合真菌性心内膜炎的描述是
A. 长期应用抗生素
B. 长期应用激素
C. 应用免疫抑制剂
D. 以皮炎芽生真菌多见
E. 以念珠菌、曲菌和组织胞浆菌多见

41. 二尖瓣关闭不全不引起
A. 左心室肥大，扩张
B. 右心室肥大，扩张
C. 左心房肥大，扩张
D. 右心房肥大，扩张
E. 心尖区舒张期隆隆样杂音

42. 联合瓣膜病可引起
A. 心前区收缩期杂音
B. 心前区舒张期杂音
C. 主动脉瓣区收缩期杂音
D. 主动脉瓣区舒张期杂音
E. 心前区收缩期杂音和主动脉瓣区收缩期杂音

43. 二尖瓣狭窄不引起
A. 右心房扩张
B. 左心房扩张
C. 主动脉瘤形成
D. 左心室肥大

E. 右心室肥大

44. 男性，34 岁，中学时期风湿病史。近年来，常感觉乏力。查体：血压 100/60mmHg，心前区闻及收缩期杂音 II～III 级，双肺底少许湿　音，肝肿大（肋下 3cm），下肢轻度可凹性水肿。符合患者疾病的描述是

A. 房间隔缺损

B. 二尖瓣关闭不全

C. 二尖瓣狭窄

D. 主动脉瓣关闭不全

E. 主动脉瓣狭窄

45. 女性，40 岁，既往有风湿病史。2 年来，经常感觉乏力、心悸。查体：血压 100/60mmHg，心前区可闻及舒张期杂音 II～III 级，双肺底少许湿　音，肝肿大（肋下 3cm），下肢轻度可凹性水肿。符合患者疾病的描述是

A. 房间隔缺损

B. 二尖瓣关闭不全

C. 二尖瓣狭窄

D. 主动脉瓣关闭不全

E. 主动脉瓣狭窄

46. 不符合慢性心瓣膜病的描述是

A. 多由风湿性心内膜炎引起

B. 表现为心瓣膜狭窄或关闭不全

C. 二尖瓣最常受累，其次是主动脉瓣

D. 狭窄和关闭不全常同时存在

E. 病变不会同时累及两个瓣膜

47. 二尖瓣关闭不全可引起

A. 球形心

B. 靴形心

C. 梨形心

D. 绒毛心

E. 虎斑心

48. 二尖瓣狭窄可引起

A. 球形心

B. 靴形心

C. 梨形心

D. 绒毛心

E. 虎斑心

49. 不符合主动脉瓣狭窄的描述是

A. 左心室肥大

B. 右心肥大、扩张

C. 主动脉瓣区吹风样收缩期杂音

D. 水冲脉

E. 心绞痛

50. 不符合主动脉瓣关闭不全的描述是

A. 左心室肥大、扩张

B. 右心肥大、扩张

C. 主动脉瓣区舒张期杂音

D. 脉压差增大，毛细血管搏动

E. 水冲脉

51. 引起主动脉瓣关闭不全的最常见原因

A. 风湿性心内膜炎

B. 主动脉粥样硬化

C. 梅毒性主动脉炎

D. Marfan 综合征

E. 感染性心内膜炎

52. 主动脉瓣狭窄不引起

A. 左心室肥大，扩张

B. 右心室肥大，扩张

C. 主动脉瓣区收缩期吹风样杂音

D. 脉压差增大

E. 心绞痛

53. 引起脉压差增大的慢性瓣膜病是

A. 主动脉瓣口狭窄

B. 主动脉瓣关闭不全

C. 二尖瓣狭窄

D. 二尖瓣关闭不全

E. 二尖瓣狭窄及主动脉瓣口狭窄

54. 男性，60 岁，既往无风湿病史。近年来，经常出现心绞痛。查体：血压 90/50mmHg，主动脉瓣闻及收缩期吹风样杂音 II～III 级，左心向左向下扩大。符合患者疾病的诊断是

A. 房间隔缺损

B. 二尖瓣关闭不全

C. 二尖瓣狭窄

D. 主动脉瓣关闭不全

E. 主动脉瓣狭窄

55. 女性，62 岁，既往无风湿病史。近年来，经常出现心绞痛。查体：血压 120/50mmHg，主动脉瓣闻及舒张期隆隆样杂音 II～III 级，左心向左向下扩大。出现水冲脉和毛细血管搏动。符合患者疾病的诊断是

A. 房间隔缺损

B. 二尖瓣关闭不全

C. 二尖瓣狭窄

D. 主动脉瓣关闭不全

E. 主动脉瓣狭窄

56. 女性，55 岁，近年来，经常出现心绞痛。查体：血压 120/50mmHg，主动脉瓣闻及舒张期隆隆样杂音 II～III 级，左心向左向下扩大。出现水冲脉和毛细血管搏动。不符合患者疾病的描述是

A. 可由风湿病引起

B. 可由 Marfan 综合征引起

C. 可由动脉粥样硬化引起

D. 可由梅毒引起

E. 可由结节性多动脉炎引起

57. 有关高血压病的描述，错误的是

A. 常引起左心肥大

B. 脑出血是主要死因

C. 引起颗粒性肾固缩

D. 引起视乳头水肿

E. 常引起下肢坏疽

58. 恶性高血压患者死于尿毒症主要是因为

A. 肾小管坏死

B. 肾小球纤维化

C. 小动脉纤维素样坏死

D. 肾间质出血

E. 肾动脉内血栓形成

59. 不伴有高血压的疾病是

A. 肾动脉狭窄

B. 慢性肾小球肾炎

C. 成人型主动脉缩窄

D. 特发性扩张性心肌病

E. 原发性醛固酮增多症

60. 良性高血压血压升高的主要因素是

A. 小动脉壁纤维素样坏死

B. 细小动脉壁玻璃样变

C. 肌型动脉中膜

D. 肌型动脉内膜增厚

E. 中型动脉粥样硬化

61. 高血压病的最主要的致病原因主要是

A. 摄盐量过多
B. 吸烟
C. 交感神经兴奋性增强
D. 肾原发性颗粒性萎缩
E. 遗传素质

62. 恶性高血压患者死于尿毒症的主要原因是
A. 肾小管坏死
B. 肾小球纤维化
C. 肾细动脉纤维素样坏死
D. 肾间质出血
E. 肾小动脉增生性动脉内膜炎

63. 良性高血压的基本病变是
A. 细动脉玻璃样变
B. 大、中动脉粥瘤
C. 小动脉纤维素样坏死
D. 动脉中层粘液变性
E. 动脉中层钙化

64. 不伴有高血压的疾病是
A. 肥厚性心肌病
B. 慢性肾小球肾炎
C. 成人性主动脉缩窄
D. 嗜铬细胞瘤
E. 原发性醛固酮增多症

65. 男性，50 岁，高血压病史 20 年。近 1 月来，血压升高至 180/110mmHg，出现头痛、头晕、眼花，有时恶心、呕吐，甚至意识模糊。符合本患者疾病的描述是
A. 脑出血
B. 脑梗死
C. 高血压危象
D. 恶性高血压
E. 心肌梗死

66. 女性，45 岁，高血压病史 15 年，并有高血压家族史。近半年来经常出现气喘、心悸，劳动后加重。查体：肥胖体型，血压 170/100mmHg，X 线检查，左心肥大。符合患者心脏病变的是
A. 心肌肥大
B. 心肌萎缩
C. 心肌梗死
D. 心肌纤维化
E. 心肌炎细胞浸润

67. 男性，50 岁，高血压病史近 20 年。近 2 年来经常性头痛、头晕，休息后减轻。查体：肥胖体形，血压 170/105mmHg，左心肥大，眼底动脉普遍狭窄，蛋白尿（+）。符合患者肾脏病变的描述是
A. 部分肾小球萎缩纤维化
B. 肾小动脉纤维素样坏死
C. 肾小管坏死
D. 部分肾小球新月体形成
E. 部分肾小球纤维素样坏死

68. 男性，40 岁，高血压病史 11 年，未经治疗。近半年来血压显著升高（180 ～ 170/130 ～120mmHg），近 1 个月来出现尿毒症。符合患者肾脏病变的是
A. 肾动脉粥样硬化
B. 肾小动脉纤维素样坏死
C. 肾小动脉淀粉样变
D. 肾小动脉炎细胞浸润
E. 肾弓形动脉粥样硬化

69. 女性，65 岁，高血压病史近 30 年。1 小

时前突然意识丧失，送急诊。查体，血压 180/100mmHg，患者神志昏迷，口角向右上方歪斜，右侧上、下肢瘫痪。符合患者疾病的描述是

A. 脑梗死
B. 脑出血
C. 脑肿瘤
D. 脑脓肿
E. 高血压脑病

70. 高血压性心脏病代偿期的特征是

A. 左心室扩张
B. 左心室向心性肥大
C. 心壁肉柱扁平
D. 弥漫性心肌纤维化
E. 左心室远心性肥大

71. 不符合原发性颗粒性固缩肾的描述是

A. 肾动脉纤维素样坏死
B. 部分肾小球纤维化
C. 入球小动脉玻璃样变
D. 部分肾小球代偿性肥大
E. 肾小管内可见蛋白管形

72. 关于高血压病脑出血的描述，错误的是

A. 以基底节、内囊部多见
B. 脑小动脉痉挛通透性增加，引起漏出性出血
C. 微小动脉瘤形成、破裂
D. 小动脉壁坏死，发生破裂性出血
E. 出血灶小，症状不明显

73. 动脉粥样硬化的危险因素不包括

A. 吸烟
B. 糖尿病
C. 高血压病
D. 动脉内膜的损伤
E. 雌激素

74. 动脉粥样硬化的危险因素不包括

A. 血浆低密度脂蛋白（LDL）水平持续升高
B. 血浆极低密度脂蛋白（VLDL）水平持续升高
C. 甲状腺素
D. 家族性高胆固醇血症
E. 家族性高甘油三酯血症

75. 抑制动脉粥样硬化症发病的因素是

A. 胆固醇
B. 高密度脂蛋白
C. 低密度脂蛋白
D. 极低密度脂蛋白
E. β 脂蛋白

76. 不参与动脉粥样硬化斑块形成的细胞是

A. 单核细胞
B. 血小板
C. 内皮细胞
D. T 淋巴细胞
E. B 淋巴细胞

77. 动脉粥样硬化症引起的非致死性病变是

A. 冠心病
B. 脑出血
C. 夹层动脉瘤
D. 肾梗死
E. 脑梗死

78. 动脉粥瘤内产生胶原的细胞是

A. 纤维母细胞
B. 巨噬细胞
C. 内皮细胞
D. 平滑肌细胞
E. 淋巴细胞

79. 动脉粥样硬化症中，不产生刺激平滑肌增生的生长因子的细胞是
A. 血小板
B. 内皮细胞
C. 平滑肌细胞
D. 巨噬细胞
E. 淋巴细胞

80. 动脉粥瘤的粥样物质的主要成分是
A. 胆固醇和胆固醇酯
B. 高密度脂蛋白
C. 低密度脂蛋白
D. 极低密度脂蛋白
E. β 脂蛋白

81. 动脉粥样硬化最早期的病变是
A. 纤维斑块
B. 粥瘤
C. 脂纹
D. 钙化
E. 溃疡形成

82. 动脉粥样硬化早期病变特点是
A. 血栓形成
B. 纤维帽形成
C. 泡沫细胞形成
D. 坏死灶形成
E. 溃疡形成

83. 粥瘤所不具备的成分是
A. 纤维瘢痕
B. 中性粒细胞
C. 无定形的坏死物质
D. 泡沫状细胞
E. 胆固醇结晶

84. 动脉粥样硬化的早期病变中，最早迁入内膜的细胞是
A. 平滑肌细胞
B. 淋巴细胞
C. 中性粒细胞
D. 单核细胞
E. 纤维母细胞

85. 动脉粥样硬化症不会出现
A. 玻璃样变
B. 淀粉样物质
C. 出血
D. 血栓形成
E. 钙化

86. 动脉粥样硬化的起始病变是
A. 平滑肌细胞迁入内膜
B. 内皮细胞慢性损伤
C. 内皮下单核细胞浸润
D. 血小板聚集
E. 纤维组织增生

87. 不符合动脉粥样硬化症的描述是
A. 泡沫细胞可来源于平滑肌
B. 脂纹可见于儿童的主动脉
C. 粥样硬化是不可逆的过程
D. 可引起动脉管腔狭窄
E. 可引起动脉瘤形成

88. 动脉粥样硬化症的复合病变不包括
A. 溃疡
B. 动脉瘤形成

C. 出血
D. 血栓形成
E. 肉芽肿形成

89. 不符合肾动脉粥样硬化的描述是
A. 可引起肾动脉狭窄
B. 可引起肾动脉血栓形成
C. 可引起肾性高血压
D. 可引起肾脏大块的凹陷瘢痕
E. 常引起肾功能不全

90. 关于主动脉粥样硬化的描述，错误的是
A. 好发于主动脉后壁和各分支开口处
B. 可引起动脉夹层
C. 可引起主动脉瓣钙化
D. 可引起管腔狭窄
E. 可继发溃疡、钙化、出血、血栓形成

91. 关于脑动脉粥样硬化的描述，错误的是
A. 引起动脉瘤形成
B. 继发脑血栓形成
C. 大脑前动脉最易受累
D. 引起脑萎缩
E. 引起脑出血

92. 不符合动脉粥样硬化性肾萎缩的叙述是
A. 可引起纤维组织增生
B. 可引起肾小动脉纤维素样坏死
C. 可引起肾梗死
D. 单侧或双侧肾受累
E. 出现大的凹陷瘢痕

93. 主动脉粥样硬化不会引起
A. 管腔狭窄
B. 血栓形成
C. 溃疡形成
D. 动脉瘤形成
E. 主动脉瓣硬化

94. 男性，73 岁，3 天前突然感觉右下肢沉重，活动受限，并伴右上肢活动受限。1 天前右侧上下肢活动能力完全丧失。符合患者疾病的描述是
A. 脑出血
B. 脑脓肿
C. 脑动脉粥样硬化伴血栓形成
D. 脑肿瘤
E. 高血压脑病

95. 主动脉粥样硬化可引起的致死性病变是
A. 溃疡形成
B. 夹层动脉瘤形成
C. 血栓形成
D. 钙化灶形成
E. 出血灶形成

96. 女性，50 岁，急性腹痛入院。既往有高血压病史。查体：血压 90/50mmHg，四肢温度低，肢端色苍白、腹部无明显包块，无腹肌紧张及反跳痛。超声诊断为主动脉夹层。可引起主动脉夹层的是
A. 巨细胞性动脉炎
B. 高安动脉炎
C. 风湿性主动脉炎
D. 动脉粥样硬化症
E. 动脉中层钙化

97. 冠状动脉粥样硬化不会引起
A. 心绞痛

B. 心肌梗死
C. 冠状动脉性猝死
D. 原发性心肌病
E. 心功能不全

98. 冠状动脉粥样硬化不会引起
A. 心肌代偿肥大
B. 心肌糖原沉积
C. 心肌间出血
D. 心肌纤维化
E. 心肌变性坏死

99. 冠状动脉粥样硬化最常累及
A. 左冠状动脉前降支
B. 左冠状动脉主干
C. 左冠状动脉回旋支
D. 右冠状动脉主干
E. 右冠状动脉回旋支

100. 引起冠状动脉性心脏病的最常见的原因是
A. 高安动脉炎
B. 冠状动脉粥样硬化
C. 巨细胞性动脉炎
D. 结节性多动脉炎
E. Wegener 肉芽肿

101. 男性，70 岁，既往体健，数小时前猝死。尸检结果：主动脉广泛粥瘤形成，两冠状动脉主干均可见粥样硬化斑块，并有管腔狭窄，未见血栓。心肌暗红色，有光泽。符合患者疾病的描述是
A. 心肌病
B. 冠状动脉性猝死
C. 克山病
D. 心肌梗塞
E. 心肌炎

102. 冠心病
A. 细动脉玻璃样变性
B. 夹层动脉瘤形成
C. 冠状动脉粥样硬化
D. 细动脉壁纤维素样坏死
E. 小血管化脓性菌栓塞

103. 心肌梗死的好发部位的顺序是
A. 左室侧壁＞左室后壁及室间隔后 1/3 ＞左室前壁及室间隔前 2/3
B. 左室前壁及室间隔前 2/3 ＞左室侧壁＞左室后壁及室间隔后 1/3
C. 左室前壁及室间隔前 2/3 ＞左室后壁及室间隔前 1/3 ＞左室侧壁
D. 左室后壁及室间隔后 1/3 ＞左室侧壁＞左室前壁及室间隔后 2/3
E. 左室后壁及室间隔后 1/3 ＞左室前壁及室间隔前 2/3 ＞左室侧壁

104. 最常引起区域性心肌梗死的原因是
A. 因体力活动造成心脏过度负荷
B. 冠状动脉痉挛
C. 冠状动脉粥样硬化并发血栓形成
D. 冠状动脉栓塞
E. 梅毒性主动脉炎所致的冠状动脉口狭窄或闭塞

105. 符合心内膜下心肌梗死的描述是
A. 由冠状动脉粥样硬化并发血栓形成引起
B. 梗死灶累及心壁各层
C. 可伴发心脏破裂
D. 由梅毒性主动脉炎引起
E. 各支冠状动脉均有狭窄

106. 有关心肌梗死的叙述，错误的是
A. 多由冠状动脉粥样硬化引起
B. 多发生于冠状动脉前降支供血的区域
C. 多为出血性梗死
D. 可引起心脏破裂
E. 可继发附壁血栓

107. 急性心肌梗死大量中性粒细胞浸润出现在
A. 梗死形成后 6 小时
B. 梗死形成后 24 小时
C. 梗死形成后第 2 天
D. 梗死形成后第 4 天
E. 梗死形成后第 7 天

108. 心肌梗死的好发部位
A. 左室后壁
B. 右心室
C. 左室侧壁
D. 室间隔
E. 左室前壁及心尖部

109. 最常引起区域性心肌梗死的原因是
A. 冠状动脉纤维素样坏死
B. 冠状动脉痉挛
C. 冠状动脉粥样硬化并发血栓形成
D. 冠状动脉栓塞
E. 冠状动脉破裂

110. 心肌梗死的最常见并发症是
A. 心脏破裂
B. 室壁瘤
C. 附壁血栓形成
D. 纤维素性心包炎
E. 心功能不全

111. 心内膜下心肌梗死的特点是
A. 引起心脏破裂
B. 引起纤维素性心包炎
C. 不波及肉柱和乳头肌
D. 多发性小灶状坏死
E. 引起室壁瘤

112. 不符合心绞痛的描述是
A. 由心肌急性暂时性缺血引起
B. 表现为心前区疼痛
C. 稳定性心绞痛病情重
D. 变异性心绞痛患者冠状动脉明显狭窄
E. 不稳定性心绞痛为心肌梗死前兆

113. 扩张性心肌病的病变特点是
A. 室间隔不匀称肥厚
B. 心肌细胞排列紊乱
C. 心室扩张、心肌间纤维化
D. 不伴有附壁血栓形成
E. 以上都不是

114. 肥厚性心肌病的病变特点是
A. 四个心腔扩张
B. 心肌细胞异常肥大，排列紊乱
C. 常伴有附壁血栓形成
D. 弥漫性心肌间质纤维化
E. 心肌细胞变性坏死

115. 克山病时心脏的病变特点是
A. 心肌细胞变性、坏死
B. 心肌间质纤维化
C. 心肌坏死灶与冠状动脉供血区无关
D. 附壁血栓形成
E. 以上都是

116. 原发性心肌病是指

A. 高血压引起的心肌损害
B. 动脉粥样硬化引起的心肌损害
C. 风湿病引起的心肌损害
D. 原因不明的心肌损害
E. 白喉引起的心肌损害

117. 限制性心肌病的病变特点是
A. 心肌细胞变性、坏死
B. 心肌细胞肥大、核深染
C. 心室充盈受限
D. 附壁血栓形成
E. 心外膜纤维化

118. 心内膜心肌纤维化的病变特点是
A. 心肌细胞变性、坏死
B. 心肌间质纤维化
C. 室间隔不规则增厚
D. 附壁血栓形成
E. 心内膜和内膜下纤维组织增生

119. 不符合心内膜弹力纤维增生症的病变特点是
A. 婴幼儿多见
B. 可合并先天性心脏病
C. 心内膜弹力纤维增生
D. 心肌间纤维组织增生
E. 心内膜胶原纤维增生

120. 不符合心脏淀粉样物质沉积的病变特点是
A. 老年人多见
B. 淀粉样物质沉积可累及全身
C. 淀粉样物质沉积可累及心脏
D. 淀粉样物质沉积在心内膜下和心肌间
E. 心肌细胞广泛变性坏死

121. 符合先天性心脏病的描述是
A. 血流自左向右分流组不发生紫绀
B. 妊娠早期预防病毒感染可降低其发病率
C. 室间隔缺损、房间隔缺损、动脉导管未闭少见
D. 无紫绀的先天性心脏病对胎儿健康影响不大
E. 亚急性感染性心内膜炎和肺炎是少见的并发症

122. Fallot 四联症所不具备的是
A. 肺动脉流出道狭窄
B. 右心室肥大
C. 房间隔缺损
D. 高位室间隔缺损，伴室间隔膜部缺损
E. 主动脉右移

123. 不符合室间隔缺损的描述是
A. 血流自右向左分流
B. 室间隔膜部缺损最多见
C. 左第三肋间闻响亮的收缩期杂音
D. 可出现传导阻滞
E. 晚期可发生肺小动脉硬化

124. 不符合房间隔缺损的描述是
A. 多为第二房间隔缺损
B. 血流自左心房向右心房分流
C. 缺损较大者、导致右心室肥大
D. 可导致肺动脉高压
E. 可合并二尖瓣裂缺

125. 不符合幼年型主动脉缩窄的描述是
A. 动脉导管前的主动脉峡部狭窄
B. 常合并动脉导管开放

C. 右心室肥大
D. 下肢不发生紫绀
E. 肺动脉血可进入降主动脉

126. 不符合成人型主动脉缩窄的描述是
A. 动脉导管后的主动脉峡部狭窄
B. 狭窄程度一般较轻
C. 胸主动脉与腹主动脉之间存在较高压差
D. 主动脉弓与降主动脉的分支之间可形成侧支循环
E. 下肢有紫绀

127. 自右向左分流（青紫型）的先天性心脏病是
A. 房间隔缺损
B. 动脉导管未闭
C. 肺动脉狭窄
D. 法洛四联症
E. 室间隔缺损

128. 无分流型先天性心脏病是
A. 房间隔缺损
B. 动脉导管未闭
C. 肺动脉狭窄
D. 法洛四联症
E. 室间隔缺损

129. 孤立性心肌炎时，心肌间质的肉芽肿不含有
A. 干酪样坏死
B. 淋巴细胞
C. 单核细胞
D. 多核巨细胞
E. 嗜酸性粒细胞

130. 属于特发性心肌炎的是
A. 弓形体性心肌炎
B. 风湿性心肌炎
C. 病毒性心肌炎
D. 细菌性心肌炎
E. 孤立性心肌炎

131. 细菌毒素引起的心肌炎是
A. 葡萄球菌性心肌炎
B. 链球菌性心肌炎
C. 肺炎双球菌性心肌炎
D. 脑膜炎双球菌性心肌炎
E. 白喉性心肌炎

132. 特发性心肌炎是
A. 弓形体性心肌炎
B. 原因不明的心肌炎
C. 病毒性心肌炎
D. 细菌性心肌炎
E. 风湿性心肌炎

133. 弓形虫性心肌炎的主要诊断依据是
A. 心肌细胞变性坏死
B. 淋巴细胞浸润
C. 中性粒细胞浸润
D. 弓形虫包囊
E. 纤维组织增生

134. 不属于免疫反应性心肌炎的是
A. 类风湿性心肌炎
B. 风湿性心肌炎
C. 特发性心肌炎
D. 红斑狼疮性心肌炎
E. 药物性心肌炎

135. 病毒性心肌炎的主要诊断依据是
A. 心肌细胞变性坏死
B. 淋巴细胞浸润

C. 中性粒细胞浸润
D. 心肌活检组织病毒基因检查阳性
E. 纤维组织增生

136. 病毒性心肌炎的常见病原体是
A. 腺病毒
B. 巨细胞病毒
C. 麻疹病毒和腮腺炎病毒
D. 乙型脑炎病毒和脊髓灰质炎病毒
E. 柯萨奇病毒和埃可病毒

137. 高血压可引起
A. 假性动脉瘤
B. 蜿蜒状动脉瘤
C. 夹层动脉瘤
D. 多发性微动脉瘤
E. 侵蚀性动脉瘤

138. 不符合梅毒性主动脉炎的描述是
A. 引起升主动脉内膜粗糙不平
B. 引起主动脉中膜树胶肿形成
C. 引起冠状动脉开口部狭窄
D. 引起主动脉瓣粘连狭窄
E. 引起主动脉瓣口扩大

139. 不引起纤维素样坏死的动脉炎是
A. 结节性多动脉炎
B. 风湿性动脉炎
C. 梅毒性主动脉炎
D. 狼疮动脉炎
E. Wegener 肉芽肿性动脉炎

140. 不符合血栓闭塞性脉管炎的描述是
A. 多累及下肢中等动脉和静脉
B. 绝大多数发生于女性
C. 动脉壁纤维素样坏死，伴有炎性细胞浸润
D. 伴有血栓形成及机化
E. 与吸烟关系密切

141. 颞动脉炎的病变特点是
A. 动脉壁内糖原沉积
B. 动脉壁粘液样变性
C. 动脉壁内脂类物质沉积
D. 动脉壁纤维素样坏死
E. 动脉壁形成多核巨细胞和类上皮细胞的肉芽肿

142. 结节性多动脉炎的病理特点是
A. 动脉壁内糖原沉积
B. 动脉壁粘液样变性
C. 动脉壁内脂类物质沉积
D. 动脉壁纤维素样坏死及炎细胞浸润
E. 动脉壁形成多核巨细胞和类上皮细胞的肉芽肿

143. Marfan 综合征的病变特点是
A. 动脉壁玻璃样变性
B. 主动脉中膜粘液样变性及坏死，弹性纤维断裂缺失
C. 动脉壁内脂类物质沉积
D. 动脉壁纤维素样坏死及炎细胞浸润
E. 动脉壁形成多核巨细胞和类上皮细胞的肉芽肿

144. 血管外伤可引起
A. 假性动脉瘤
B. 蜿蜒状动脉瘤
C. 夹层动脉瘤
D. 多发性微动脉瘤
E. 侵蚀性动脉瘤

145. Marfan 综合征可引起
A. 假性动脉瘤

B. 蜿蜒状动脉瘤
C. 夹层动脉瘤
D. 多发性微动脉瘤
E. 侵蚀性动脉瘤

146. 慢性空洞性肺结核可引起
A. 假性动脉瘤
B. 蜿蜒状动脉瘤
C. 夹层动脉瘤
D. 多发性微动脉瘤
E. 侵蚀性动脉瘤

147. 尿毒症引起的心包炎可表现为
A. 出血性心包炎
B. 化脓性心包炎
C. 浆液性心包炎
D. 纤维素性心包炎
E. 肉芽肿性心包炎

148. 结核性心包炎的特征性病理变化是
A. 浆液渗出
B. 伴有出血
C. 纤维蛋白渗出
D. 心包粘连
E. 肉芽肿形成

149. 不符合心脏肿瘤的特点是
A. 原发性肿瘤罕见
B. 转移性肿瘤多见
C. 粘液瘤最常见
D. 横纹肌瘤较常见
E. 可发生恶性间皮瘤

150. 引起缩窄性心包炎的最常见原因是
A. 风湿性心包炎
B. 狼疮性心包炎
C. 结核性心包炎
D. 病毒性心包炎
E. 尿毒症性心包炎

151. 非感染性心包炎是
A. 化脓性心包炎
B. 真菌性心包炎
C. 结核性心包炎
D. 病毒性心包炎
E. 尿毒症性心包炎

152. 肿瘤性心包炎的特征性病理变化是
A. 浆液渗出
B. 伴有出血
C. 纤维蛋白渗出
D. 可见肿瘤细胞
E. 肉芽肿形成

153. verrucous endocartis
A. 风湿性心肌炎
B. 风湿性心内膜炎
C. 疣状心内膜炎
D. 风湿性心外膜炎
E. 风湿性全心炎

154. mitral stenosis
A. 二尖瓣狭窄
B. 二尖瓣关闭不全
C. 主动脉瓣狭窄
D. 主动脉瓣关闭不全
E. 心瓣膜病

155. malignant hypertension
A. 良性高血压
B. 恶性高血压
C. 临界性高血压
D. 良性增生
E. 恶性增生

156. atheroma
A. 粥样硬化
B. 脂纹
C. 动脉瘤
D. 粥瘤
E. 硬化

157. stable angina pectoris
A. 心肌梗死
B. 内膜下心肌梗死
C. 心绞痛
D. 变异性心绞痛
E. 稳定性心绞痛

158. idiopathic myocarditisa
A. 寄生虫性心肌炎
B. 细菌性心包炎
C. 原发性心肌病
D. 特发性心肌炎
E. 扩张性心肌病

159. hypertrophic cardiomyopathy
A. 扩张性心肌病
B. 特发性心肌炎
C. 肥厚性心肌病
D. 缩窄性心包炎
E. 限制性心肌病

160. tetralogy of Fallot
A. 室间隔缺损
B. 房间隔缺损
C. 动脉导管开放
D. 法洛四联症
E. 主动脉狭窄

英文题

1. A 30-year-old woman had acute rheumatic fever as a child and when she was around 20 years of age she developed a new cardiac murmur. Now she presents with fever, intermittent severe pain in the left upper quadrant of her abdomen, and Roth spots in the retina. At the present time one culture submitted to the hospital laboratory grew out a specific type of bacteria. Which of the following is the most likely cause of her disease?
A. Candida species
B. Staphylococcus aureus
C. Tubercle bacillus
D. Alpha-hemolytic viridans streptococci
E. Group A streptococci

2. A 10-year-old boy has experienced increased fatigue and decreased exercise tolerance for the past 2 weeks. He presented with an acute onset of fever, and pain in several joints 2 weeks before. Physical examination finds an enlarged heart and systolic murmur. Laboratory tests find an elevated erythrocyte sedimentation rate and an elevated antistreptolysin O titer. Which of the following lesions is most likely to be found in the heart of this patient?
A. Foci of fibrinoid necrosis surrounded with Anitsehkow cells in the intersti-

tial tissue of the heart

B. Large fragile vegetations formed along the lines of closure of the mitral valves

C. Coagulative necrosis of the cardiac fibers with neutrophilic infiltration

D. Haphazardly arranged hypertrophied myocytes in the interventricular septum

E. Mural thrombosis in the left atrium

3. A 30-year-old man who has a history of rheumatic fever during childhood presents with shortness of breath and cough. Physical examination finds an early diastolic opening snap with a rumbling late diastolic murmur. Which of the following is the most likely diagnosis?

A. Pulmonic stenosis

B. Aortic regurgitation

C. Mitral regurgitation

D. Aortic stenosis

E. Mitral stenosis

4. A 65-year-old man presents with signs of angina, syncopal episodes with exertion and left heart failure. Physical examination reveals a midsystolic ejection murmur. Pressure studies reveal that the left ventricular pressure during systole is markedly greater than the aortic pressure. Which of the following is the most frequent cause of the cardiac valvular abnormality present in this patient?

A. Rheumatic fever

B. Congenital aplasia

C. Atherosclerosis of aortic root

D. Acute infective endocarditis

E. Aneurysm of aortic root

5. A 65-year-old man presents shortness of breath, cough, and nocturnal dyspnea. Physical examination reveals a hyperdynamic, bounding, "water-hammer" pulse and a diastolic murmur. Which of the following is the most likely diagnosis?

A. Mitral stenosis

B. Mitral insufficiency

C. Constrictive pericarditis

D. Aortic insufficiency

E. Aortic stenosis

6. A 70-year-old woman presents with increasing chest pain and occasional syncopal episodes, especially with physical exertion. She has nocturnal dyspnea and orthopnea. Physical examination reveals a midsystolic ejection murmur. An echocardiogram shows concentric hypertrophy of the left ventricle. Which of the following is the most likely diagnosis?

A. Mitral stenosis

B. Mitral regurgitation

C. Constrictive pericarditis

D. Aortic regurgitation

E. Aortic stenosis

7. A 60-year-old woman developed left heart failure and died suddenly. A section

from her heart at the time of autopsy reveals marked thickening of the wall of the left ventricle with decrease in cavity size, but the thickness of the right ventricle is within normal limits. Many of the nuclei of the myocytes in the wall of the left ventricle are hyperchromatic and rectangular. The endocardium and the cardiac valves are normal. What is the most likely cause of this cardiac pathology?

A. Eccentric hypertrophy
B. Cor pulmonale
C. Systemic hypertension
D. Volume overload
E. Ischemic heart disease

8. Several days after a myocardial infarction, a 60-year-old man develops a sudden onset of a new pansystolic murmur along with a diastolic flow murmur. Workup reveals increased left atrial pressure that develops late in systole and extends into diastole. Which of the following is the most likely cause of the abnormalities present in this individual?

A. Thrombosis of the left atrial cavity
B. Rupture of a papillary muscle
C. Rupture of the left ventricular wall
D. Aneurysmal dilation of the left ventricle
E. Obstruction of the aortic valve

9. Areas of myocardial infarction undergo a series of gross changes. Which of the following is the earliest histologic change in myocardial infarction?

A. Formation of wavy fibers
B. Pyknosis, karyorrhexis and karyolysis of cardiac muscle fibers
C. Liquefaction necrosis of cardiac muscle fibers
D. Hyperemia and hemorrhage in the border zone around the infarction
E. Interstitial hemorrhage and neutrophil infiltration

10. Which of the following complications will be eariest to arise following a transmural myocardial infarction?

A. Rupture of the heart
B. Ventricular aneurysm
C. Mural thrombosis
D. Acute serofibrinous pericarditis
E. Congestive heart failure

11. Arrange the following numbered statements in the correct order according to the expected sequence of events that normally occur during healing of a myocardial infarction. 1 = Collagen is deposited, forming a fibrous scar; 2 = Mitochondrial swelling and distortion of cristae; 3 = Granulation tissue begins to form; 4 = Macrophages begin to arrive at the area of coagulative necrosis; 5 = Neutrophils begin to arrive at the area of coagulative necrosis

A. 5, then 4, then 3, then 2, then 1

B. 2, then 5, then 4, then 3, then 1
C. 2, then 4, then 5, then 3, then 1
D. 4, then 5, then 3, then 2, then 1
E. 2, then 3, then 4, then 5, then 1

12. A 65-year-old man develops intense crushing substernal, chest pain that is not relieved by nitroglycerin. The ECG shows ST-segment elevation and abnormal Q waves. His condition rapidly worsens and he dies 2 days later. At autopsy a 2.5 cm^2 necrotic area is found that involves the anterior left ventricle. What is the most likely cause of this patient's myocardial infarction?
A. Fibrinoid necrosis of main trunk of left coronary artery
B. Severe stenosis of all three main branches of coronary artery without thrombosis
C. Thrombosis of the anterior descending branch of the left coronary artery
D. Complete occlusion of the left circumflex coronary artery
E. Vasospasm of the right coronary artery

13. A 65-year-old man presents with recurrent chest pain that comes on whenever he feels tired while working over the past 3 years. In the past 3 months he has found the pain can not be relieved by rest, and the pain has increased in duration and frequency. Which of the following is most likely present in this patient?
A. LDH and CPK level is increased in his serum
B. Newly formed thrombosis on the previous atheromatous plaque in the anterior descending branch of the left coronary artery
C. Fibrinous exudation on the epicardium
D. The ECG shows elevated ST segment
E. Pulmonary edema and hepatomegaly are present


14. A 65-year-old man presents with recurrent chest pain that comes on whenever he feels tired while he is working. He states that the pain goes away after a couple of minutes if he stops and rests or takes nitroglycerin. His ECG shows only nonspecific changes. Which of the following is the most likely diagnosis?
A. Rheumatic pericarditis
B. Atypical angina
C. Stable angina
D. Myocardial infarction
E. Unstable angina

15. A 65-year-old man has a long history of poorly controlled hypertension. On physical examination, he is afebrile. His neck veins are distended. Crepitant rales can be heard in both lower lungs. He has pitting edema of his feet bilaterally and mild hepatomegaly. A chest x-ray reveals his heart is dilated and enlarged. Which of the following is the basic defect that caused these

clinical signs?

A. Decreased viscosity of blood caused by decreased diastolic filling of his heart
B. Increased contractility of cardiac muscle caused by high-output cardiac failure
C. Decreased peripheral resistance caused by low-output cardiac failure
D. Increased afterload caused by systolic dysfunction of the heart
E. Increased preload caused by diastolic dysfunction of his heart

16. A 55-year-old woman was being evaluated for shortness of breath, ankle edema, and mild hepatomegaly in the hospital. She died 1 week later after admission. The gross appearance of the liver at autopsy is shown. Which of the following abnormalities is most likely present?



A. Right heart failure
B. Splenic infarction
C. Atherosclerosis of descending aorta
D. Portal vein thrombosis
E. A pulmonary saddle embolus

17. A 65-year-old man was found to have a 5-cm pulsatile mass in his abdomen during a routine physical examination. Angiography reveals a marked dilation in his aorta distal to the renal arteries. Which of the following is the most likely cause of this aneurysm?

A. A congenital defect
B. Hypertension
C. Trauma
D. Rheumatic fever
E. Atherosclerosis

18. A 60-year-old man has had hypertension for 20 years. His blood pressure has ranged from 150/95 mmHg to 220/110 mmHg. He has diabetes mellitus and has had signs of renal failure since last year. During the workup of her disease, a renal biopsy is performed. Which of the following vascular lesions is most likely to be found in the kidney of this patient?

A. Dystrophic calcification in the tunica media of muscular arteries
B. The lumens of the afferent glomerular arterioles are narrowed by uniform, homogenous, pink deposits within the vessel walls
C. Fibrinoid necrosis of the intima and tunica media of vascular wall with infiltration of monocytes and neutrophilic

granulocytes

D. A concentric obliteration of arterioles by an increase in smooth muscle cells and protein deposition in a laminar configuration

E. Chronic granulomatous inflammation with giant cells and breaks in the internal elastic lamina

19. A 50-year-old woman presents with a long history of hypertension, and signs of renal failure. A renal biopsy is performed and shows the lumens of the small blood vessels are narrowed by uniform, homogenous, pink deposits within the walls of these vessels. Which of the following is the most likely diagnosis?

A. Hyperplastic arteriolosclerosis

B. Thromboangiitis obliterans

C. Arteriosclerosis obliterans

D. Hyaline arteriolosclerosis

E. Medial calcific sclerosis

20. A 33-year-old man presents with signs and symptoms of hyperglycemia. Laboratory evaluation of his serum also finds hypertriglyceridemia, which is due to his diabetes. The most common type of secondary hyperlipidemia associated with diabetes mellitus is characterized by elevated serum levels of which one of the following substances?

A. Intermediate-density lipoproteins

B. High-density lipoproteins

C. Very-low-density lipoproteins

D. Low-density lipoproteins

E. Chylomicrons

21. A 30-year-old woman has angina pectoris of 6 months duration. On physical examination, her blood pressure is 135/85mm Hg. She is 162cm tall and weighs 80kg. Coronary angiography shows 75% narrowing of the anterior descending branch of the left coronary artery. Which of the following is the major risk factor associated with these findings?

A. Job occupation

B. Gender

C. Diabetes mellitus

D. Obesity

E. Type A personality


22. For more than a decade, a 40-year-old man has had poorly controlled hypertension ranging from 150/90mm Hg to 160/95mm Hg. Over the past 3 months, his blood pressure has increased to 250/125mm Hg. A chest radiograph shows a prominent border on the left side of the heart. Laboratory studies show that his serum creatinine level has increased during this time from 1.7mg/dL to 3.8mg/dL. Which of the following lesions is most likely to be found in the afferent glomerular arterioles of this patient?

A. Granuloma formation

B. Fibrinoid necrosis


C. Dystrophic calcification
D. Hyaline degeneration
E. Amyloid change

23. A 60-year-old woman has reported increasing fatigue over the past year. Laboratory studies show a serum creatinine level of 4.7 mg/dL（normal values: male: 0.6-1.2mg/dl, female: 0.5-1.1mg/dl）and urea nitrogen level of 44 mg/dL (normal value: adult: 9-20mg/dl, child: 5-18mg/dl). An abdominal ultrasound scan shows that her kidneys are symmetrically shrunken and granular. The high-magnification microscopic appearance of the kidneys is shown. These findings are most likely to indicate which of the following underlying conditions?




A. Benign hypertension
B. Hypercholesterolemia
C. Malignant hypertension
D. Advanced breast carcinoma with widespread metastasis
E. Rheumatic arteritis

24. A 60-year-old man has experienced chest pain at rest for the past year. The figure shows the microscopic appearance representative of the patient's left anterior descending artery. Which of the following sets of histologic features is most likely to be found in this patient?



A. A core of foam cells covered by a cap of smooth muscle cells
B. Collection of foam cells, necrotic areas, cholesterol clefts and calcification
C. A lipid core, granulation tissue
D. Lipid-filled foam cells and T lymphocytes
E. Cholesterol clefts surrounded by proliferating smooth muscle cells

25. The gross appearance of the aorta of a 10-year-old girl who died of acute leukemia at autopsy is shown in the preceding figure. Histologic examination of the linear pale marking is most likely to show which of the following features?



A. A lipid core, granulation tissue
B. Collection of foam cells, necrotic areas, and calcification
C. Cholesterol clefts surrounded by proliferating smooth muscle cells
D. A core of lipid debris covered by a cap of smooth muscle cells
E. Lipid-filled foam cells and T lymphocytes

26. Smoking, hypertension, and hypercholesterolemia are the main factors that are found to play a significant role in the causation of atherosclerosis. Which of the following events is the most important direct biologic consequence of these factors?
A. Alterations of hepatic lipoprotein receptors
B. Endothelial injury and its sequelae
C. Inhibition of LDL oxidation
D. Alterations of endogenous factors regulating vasomotor tone
E. Conversion of smooth muscle cells to foam cells

27. The formation of atheromatous plaque in coronary arteries is observed in an experiment. Atherosclerotic plaques are shown to change slowly but constantly in ways that can promote clinical events, including acute coronary syndromes. Which of the following types of cells is the initial target in the pathogenesis of this arterial lesion?
A. Neutrophils
B. Smooth muscle cells
C. Endothelial cells
D. Platelets
E. Monocytes

28. A 50-year-old man has had pain in the left shoulder and arm for the past 12 hours. Over the next 6 hours, he develops shortness of breath, which persists for 2 days. The patient is admitted to the hospital and continues to experience dyspnea for the next 3 days. Coronary angiography shows 50% proximal occlusion of the right coronary and circumflex arteries, and 75% occlusion of the left anterior descending artery. On day 7 after the onset of shoulder pain, he has a cardiac arrest. Which of the following atherosclerosis plaque alterations is not significantly associated with the patient's death?
A. Intermittent platelet aggregation
B. Thrombosis
C. Ulceration of the plaque surface
D. Thinning of the media
E. Hemorrhage into the plaque substance

29. A 57-year-old man has experienced angina on exertion for the past 7 years. A coronary angiogram performed 2 years ago showed 75% stenosis of the left anterior descending coronary artery and 50% stenosis of the right coronary artery. For the past 3 weeks, the frequency and sever-

ity of the anginal attacks have increased, and pain sometimes occurs even when he is lying in bed. Which of the following is most likely to explain these findings?

A. Increasing stenosis of right coronary artery
B. Hypertrophy of ischemic myocardium with increased oxygen demand
C. Reduction in oxygen-carrying capacity due to pulmonary congestion
D. Elevated level of blood pressure
E. Fissuring of plaque in left coronary artery with superimposed mural (partial) thrombosis

30. A 33-year-old woman has had increasingly severe dyspnea, orthopnea, and swelling of the legs for the past 2 weeks. On physical examination, her temperature is 37.8℃, pulse 83/min, respirations 20/min, and blood pressure 100/60mmHg. An ECG shows episodes of ventricular tachycardia. An echocardiogram shows right and left ventricular dilation but no valvular deformities. An endomyocardial biopsy shows focal myocyte necrosis and lymphocytic infiltrate. Which of the following organisms most likely caused the infection?

A. Haemolytic streptococcus
B. Mycobacterium tuberculosis
C. Staphylococcus aureus
D. Coxsackievirus A
E. Streptococcus viridans

31. A 55-year-old man has had increasing abdominal discomfort and swelling of his legs for the past 3 years. On physical examination, the liver is enlarged and tender and can be palpated 8 cm below the right costal margin. Pitting edema is observed on the lower extremities. A chest radiograph shows pleural effusions, and increased lucency of lung fields. Thoracentesis on the right side yields 500mL of clear fluid with few cells. Which of the following is most likely to produce these findings?

A. Chronic obstructive pulmonary disease
B. Pulmonary valve stenosis
C. Tricuspid valve Stenosis
D. Systemic hypertension
E. Acute myocardial infarction

32. A 11-year-old girl manifests choreiform movements and begins to complain of pain in her knees and hips, particularly with movement 3 weeks after a bout of acute pharyngitis. She develops subcutaneous nodules over the skin of her arms and torso at the same time. A friction rub is heard on auscultation of the chest. Which of the following serum laboratory findings is most characteristic of the disease affecting this patient?

A. Increased creatine kinase-MB fraction

(CK-MB)


B. Positive blood culture for Staphylococcus aureus

C. High double-stranded DNA autoantibody titer

D. Elevated anti-streptolysin O level

E. Elevated cholesterol level

33. A 37-year-old woman dies suddenly at home. External examination of the body shows no evidence of trauma. The microscopic appearance of the proximal left anterior descending artery at autopsy is shown. Which of the following conditions is most likely to be the underlying cause of death?



A. Marfan syndrome

B. Rheumatic arteritis

C. Diabetes mellitus

D. Aneurysm

E. Advanced breast carcinoma with widespread metastases

34. A 33-year-old man died suddenly after a week-long illness. He experienced chest pain and then became increasingly dyspneic and nauseated and lost consciousness multiple times. External examination shows no evidence of trauma. The body height is 166cm and he weighs 75kg. This figure which shows the gross appearance of the chest cavity at autopsy with the pericardial sac opened, is most consistent with which of the following underlying causes of death?



A. Dilated cardiomyopathy

B. Transmural myocardial infarction

C. Disseminated tuberculosis

D. Rheumatic heart disease

E. Marfan syndrome

35. A 16-year-old girl has fever and chest pain. Two weeks earlier, she had a mild upper respiratory tract infection. On physical examination, her temperature is 37 ℃, pulse 90/min, respirations 20/min, and blood pressure 85/45mm Hg. A friction rub is audible on auscultation of the chest. A chest radiograph shows pulmonary edema. An echocardiogram shows small vegetations at the closure line

of the mitral and aortic valves. An endocardial biopsy shows focal interstitial inflammation with Aschoff nodules and Anichkov cells. Her condition improves over the next month. Which of the following additional complications of this illness is most likely to be seen in the patient?

A. Dilated cardiomyopathy
B. Mitral valve prolapse
C. Constrictive pericarditis
D. Left ventricular aneurysm
E. Mitral stenosis and insufficiency

36. A 48-year-old man has experienced worsening exercise tolerance accompanied by marked shortness of breath for the past 6 months. He was healthy previously and he is now afebrile, but he has pitting edema to the knees. A chest radiograph shows cardiomegaly and pulmonary edema with pleural effusions. An echocardiogram shows four-chamber cardiac dilation and mitral and tricuspid valvular regurgitation, with an ejection fraction of 30%. A coronary angiogram shows no more than 10% narrowing of the major coronary arteries. Which of the following is the most likely diagnosis?

A. Constrictive pericarditis
B. Hypertrophic cardiomyopathy
C. Rheumatic heart disease
D. Idiopathic dilated cardiomyopathy
E. Diabetes mellitus

37. An 18-year-old man was send to emergency room for suddenly collapse. His vital signs include a temperature of 37.1℃, pulse 84/min, respirations 18/min, and blood pressure 80/40mm Hg. Laboratory findings include hemoglobin of 13.56g/dL, platelet count 252,000/mm^3, WBC count 7230/mm^3, serum glucose 73mg/dL, and creatinine 1.2mg/dL. The total creatine kinase (CK) level is elevated, with a CK-MB fraction of 10%. Which of the following underlying conditions is most likely to be present in this patient?

A. Congenital ventricular septal defect
B. Familial hypercholesterolemia
C. Marfan syndrome
D. Rheumatic fever
E. Down syndrome

38. A 55-year-old man experienced chest pain with exertion 3 years ago, and an angiogram showed 75% occlusion of his left anterior descending artery. He was prescribed long acting nitrates and beta blockers, and he remained asymptomatic until the past week. He experienced sudden cardiac arrest when he was admitted to hospital yesterday. Which of the following laboratory findings is most characteristic of the disease affecting this patient?

A. High double-stranded DNA autoanti-

body titer


B. Positive ANCA determination

C. Increased creatine kinase-MB fraction (CK-MB)

D. Elevated anti-streptolysin O titer

E. Elevated renin level

39. A 50-year-old man has had increasing dyspnea for the past 2 days. He experiences sudden cardiac arrest and cannot be resuscitated. The light microscopic appearance of the left ventricular free wall at autopsy is shown above. Which of the following is the most likely diagnosis?



A. Viral myocarditis

B. Acute rheumatic myocarditis

C. Myocardial infarction

D. Hypertrophic cardiomyopathy

E. Septic embolization

40. A 50-year-old man experiences crushing substernal chest pain one morning. Over the next 4 hours, the pain persists and begins to radiate to his left arm. He becomes diaphoretic and short of breath and finally goes to the hospital in the evening. An elevated serum value in which of the following laboratory tests would be most useful for diagnosis of this patient on admission to the hospital?

A. CK-MB fraction

B. Lipase

C. AST

D. C-reactive protein

E. ALT

41. A 11-year-old girl who is normally developed has chronic progressive exercise intolerance. Physical examination shows a temperature of 37.1℃, pulse 70/min, respirations 14/min, and blood pressure 100/60 mmHg. A chest radiograph shows cardiomegaly and mild pulmonary edema. An echocardiogram shows severe left ventricular hypertrophy and a prominent interventricular septum. The right ventricle is slightly thickened. During systole, the anterior leaflet of the mitral valve moves into the outflow tract of the left ventricle. The ejection fraction is abnormally high, and the ventricular volume and cardiac output are both low. Which of the following is the most likely cause of the cardiac abnormalities in this patient?

A. Endomyocardial fibrosis

B. Mutations in the P-myosin heavy chain

C. Autoimmunity against myocardial fibers

D. Elevated anti-streptolysin O titer

E. ECHO virus infection

42. A 28-year-old man was found dead at home. An external examination by the medical examiner showed splinter hemorrhages under the fingernails and no signs of trauma. The gross appearance of the heart at autopsy is shown. Which of the following conditions is most likely to have been the cause of his death?



A. Acute infective endocarditis
B. Acute rhematic heart disease
C. Acute myocardial infarction
D. Vasculitis
E. Viral myocarditis

43. A 65-year-old man with a history of diabetes mellitus had chest pain and an elevated ST level 1 year ago. He was treated in the hospital with antiarrhythmic agents for 1 week. An echocardiogram showed an ejection fraction of 28%. He now has markedly reduced exercise tolerance. Diffuse crackles are heard on auscultation of the lungs. The representative gross appearance of the heart is shown in the following figure. Which of the following complications of this disease is the patient most likely to develop?



A. Infective endocarditis
B. Chronic valvular disease of the heart
C. Hypertrophic cardiomyopathy
D. Constrictive pericarditis
E. Systemic thromboembolism

44. A 16-year-old girl died suddenly when she just finished a running. She had been healthy all her life and complained only of limited episodes of chest pain in games during the current schoo1 year. Which of the following pathologic findings of the heart is the medical examiner most likely to find?

A. Large, friable vegetations with destruction of aortic valve cusps
B. Thickening of the endocardium with white color change
C. Mitral valvular stenosis with left atrial enlargement
D. Haphazardly arranged hypertrophied septal myocytes
E. Extensive myocardial hemosiderin deposition

45. A 65-year-old man has had progressive dyspnea for the past 2 years. On physical examination, he is afebrile, his pulse is 77/min, respirations 17/min, and extensive rales are heard in all lung fields. An echocardiogram shows that the left ventricular wall is markedly hypertrophied. A chest radiograph shows pulmonary edema and a prominent left-sided heart shadow. Which of the following conditions has most likely produced these findings?
    A. Mitral stenosis
    B. Centrilobular emphysema
    C. Silicosis
    D. Systemic hypertension
    E. Tricuspid valve regurgitation
46. For the past 2 and half years, a 55-year-old man has experienced increased fatigue and decreased exercise tolerance. On auscultation, diffuse crackles are audible. The abdomen is distended with a fluid wave, and there is bilateral pitting edema to the knees. A chest radiograph shows pulmonary edema and marked cardiomegaly. An echocardiogram shows mild tricuspid and mitral regurgitation and reduced right and left ventricular wall motion, with an ejection fraction of 30%. Over the next year he suffers cerebral, renal, and splenic infarctions, and also has infarctions of two toes of his right foot. Which of the following is most likely to have produced these findings?
    A. Rheumatic heart disease
    B. Hypertrophic cardiomyopathy
    C. Dilated cardiomyopathy
    D. Coronary atherosclerosis
    E. Restrictive cardiomyopathy
47. A 57-year-old man has a sudden onset of severe substernal chest pain that radiates to the neck. He has tachycardia, hyperventilation, and hypotension on physical examination. No cardiac murmurs are heard on auscultation. Emergent coronary angiography shows a thrombotic occlusion of the left circumflex artery with areas of 60% to 75% narrowing in the proximal circumflex and anterior descending arteries. Which of the following complications of this disease is likely to occur in the future?
    A. Myocardial rupture
    B. Thromboembolism
    C. ventricular aneurysm
    D. Pericarditis
    E. All above
48. A 17-year-old man has had a low-grade fever for 3 weeks. On physical examination, his temperature is 38.5 ℃, pulse 104/min, respirations 28/min, and blood pressure of 97/60 mm Hg. His spleen tip is palpable and tender. There are also tender hemorrhagic nodules on the palms and soles. A heart murmur is heard on auscul-

tation. Which of the following infectious agents is most likely to be cultured from this patient's blood?

A. Coxsackievirus B
B. Streptococcus viridans group
C. Staphylococci
D. Mycobacterium tuberculosis
E. Candida albicans

49. A 10-year-old boy had a sore throat and fever 3 weeks ago, and a throat culture was positive for group A b-hemolytic Streptococcus. He is afebrile. On auscultation, a murmur due to mitral regurgitation is audible, and there are diffuse rales over both lungs. Over the next 2 days he has several episodes of atrial fibrillation accompanied by signs of acute left ventricular failure. Which of the following pathologic changes which occurred in this child's heart during hospitalization is most likely to be the cause of the left ventricular failure?

A. Fibrinous pericarditis
B. Fibrosis of mitral valve with fusion of commissures
C. Myocarditis
D. Verrucous endocarditis
E. Amyloidosis

50. A 62 year-old man experiences episodes of severe substernal chest pain every time he performs a task that requires moderate exercise. The episodes have become more frequent and severe over the past year, but they can be relieved by sublingual nitroglycerin. On physical examination, there are no murmurs or gallops. A chest radiograph shows the lungs are clear and the heart size is nearly normal. Laboratory studies show the total serum cholesterol is 223mg/dL. Which of the following cardiac lesions is most likely to be present?

A. Rheumatic mitral stenosis
B. Coronary atherosclerosis
C. Restrictive cardiomyopathy
D. Viral myocarditis
E. Pericardial tamponade

答案

中文题

1.D	2.B	3.B	4.D	5.E	6.C	7.C	8.C	9.B	10.A
11.C	12.B	13.B	14.E	15.A	16.D	17.D	18.C	19.D	20.B
21.B	22.C	23.B	24.D	25.B	26.A	27.C	28.B	29.D	30.A
31.E	32.E	33.C	34.E	35.D	36.C	37.D	38.C	39.E	40.D
41.E	42.E	43.D	44.B	45.C	46.E	47.A	48.C	49.D	50.C
51.A	52.D	53.B	54.E	55.D	56.E	57.E	58.C	59.D	60.B
61.D	62.C	63.A	64.A	65.C	66.A	67.A	68.B	69.B	70.B
71.A	72.D	73.E	74.C	75.B	76.E	77.D	78.D	79.E	80.A
81.C	82.C	83.B	84.D	85.B	86.B	87.C	88.E	89.E	90.D
91.C	92.B	93.A	94.C	95.B	96.D	97.D	98.B	99.A	100.B
101.B	102.C	103.C	104.C	105.E	106.C	107.A	108.E	109.C	110.E
111.D	112.C	113.C	114.B	115.E	116.D	117.C	118.E	119.D	120.E
121.B	122.C	123.A	124.B	125.D	126.E	127.D	128.C	129.A	130.E
131.E	132.B	133.D	134.C	135.D	136.E	137.D	138.D	139.C	140.B
141.E	142.D	143.B	144.A	145.C	146.E	147.D	148.E	149.B	150.C
151.E	152.E	153.C	154.A	155.B	156.D	157.E	158.D	159.C	160.D

英文题

1.D	2.A	3.E	4.A	5.D	6.E	7.C	8.B	9.A	10.D
11.B	12.C	13.B	14.C	15.D	16.A	17.E	18.B	19.D	20.C
21.C	22.B	23.A	24.B	25.E	26.B	27.C	28.D	29.E	30.D
31.A	32.D	33.C	34.B	35.E	36.D	37.B	38.C	39.C	40.A
41.B	42.A	43.E	44.D	45.D	46.C	47.E	48.B	49.C	50.B

6　呼吸系统疾病

中文题

1. 最常见的慢性阻塞性肺病是
   A. 支气管扩张
   B. 支气管哮喘
   C. 慢性支气管炎
   D. 矽肺
   E. 慢性空洞性肺结核
2. 不符合慢性支气管炎的描述是
   A. 可引起肺心病
   B. 可引起肺气肿
   C. 吸烟是常见的原因
   D. 可伴喘息症状
   E. 支气管呈囊性扩张
3. 与慢性支气管炎发病无关的是
   A. 呼吸道反复病毒感染
   B. 肺孢子虫感染
   C. 呼吸道细菌感染
   D. 大气污染
   E. 吸烟
4. 慢性支气管炎的临床表现是
   A. 慢性咳嗽
   B. 咳铁锈色痰
   C. 肺实变
   D. 最大通气量明显降低
   E. 大咯血
5. 慢性支气管炎的主要病变不包括
   A. 粘膜上皮纤毛脱落、倒伏
   B. 肉芽肿形成
   C. 粘液腺肥大
   D. 管壁淋巴细胞、浆细胞浸润
   E. 鳞状上皮化生
6. 慢性支气管炎患者通气功能障碍的病理基础是
   A. 粘液腺肥大、增生
   B. 上皮纤毛倒伏、脱落
   C. 软骨变性萎缩
   D. 小气道阻塞
   E. 管壁平滑肌萎缩
7. 女性，50 岁，慢性咳嗽，咳白色泡沫粘痰，每年秋冬发作，持续三个月左右。不符合该患者疾病的描述是
   A. 可发生肺气肿
   B. 可发生支气管肺炎
   C. 可发生肺源性心脏病
   D. 可发生支气管扩张
   E. 可发生鳞状上皮化生
8. 女性，50 岁，慢性咳嗽，咳白色泡沫粘痰，每年秋冬发作，持续三个月左右。不符合该患者疾病的描述是
   A. 支气管粘膜上皮变性坏死
   B. 支气管粘膜上皮杯状细胞增多
   C. 支气管粘液腺肥大
   D. 支气管软骨变性萎缩
   E. 支气管壁弹力纤维增生
9. 不符合急性细支气管炎的特点是
   A. 累及管径＜ 2mm 的细支气管
   B. 常见于婴幼儿
   C. 多由流感杆菌、葡萄球菌等细菌感染引起

D. 多引起喘憋性呼吸困难
E. 可引起呼吸衰竭

10. 急性细支气管炎出现呼吸困难的主要病变基础是
A. 细支气管上皮坏死脱落
B. 细支气管管腔内炎性渗出物阻塞
C. 肺间质炎细胞浸润
D. 肺萎陷
E. 急性阻塞性肺气肿

11. 不符合支气管哮喘的描述是
A. 属 I 型超敏反应性疾病
B. 有多种炎症介质参与
C. 支气管内大量粘液分泌
D. 支气管壁多量嗜碱粒细胞浸润
E. 支气管平滑肌痉挛

12. 引起支气管哮喘的主要原因是
A. 支气管粘膜内杯状细胞增多
B. 支气管粘膜水肿
C. 支气管壁平滑肌痉挛
D. 支气管腔内粘液增多
E. 支气管管壁炎细胞浸润

13. 符合支气管哮喘的描述是
A. 单基因遗传病
B. 常染色体显性遗传病
C. 常染色体隐性遗传病
D. 多因素遗传病
E. 性连锁遗传病

14. 不符合支气管哮喘的描述是
A. 多种炎细胞参与
B. 多种炎症介质参与
C. 多种细胞因子参与
D. 嗜酸粒细胞是最主要的效应细胞
E. 气道的高反应性是重要特征

15. 不符合支气管哮喘病变的描述是
A. 可见胆固醇结晶
B. 可见 Curschmann 螺旋
C. 支气管内粘液栓填塞
D. 支气管壁多量炎细胞浸润
E. 支气管平滑肌肥厚

16. 不符合支气管哮喘的合并症的描述是
A. 可引起肺气肿
B. 可引起肺炎
C. 可引起肺心病
D. 可引起支气管扩张
E. 不引起急性死亡

17. 引起肺气肿的因素有
A. 小气道的狭窄、阻塞
B. 肺泡壁破坏、弹性减弱
C. 弹性蛋白酶增多
D. 遗传性 α_1-抗胰蛋白酶缺乏
E. 以上都是

18. 腺泡中央型肺气肿病变主要累及
A. 肺泡管
B. 终末细支气管
C. 肺泡囊
D. 呼吸性细支气管
E. 细支气管

19. 累及肺腺泡各部分的肺气肿为
A. 腺泡中央型
B. 全腺泡型
C. 腺泡周围型
D. 间质性肺气肿
E. 不规则型

20. 在小叶间隔与肺膜连接处形成串珠状

小气泡，应诊断为

A. 小叶周围型

B. 不规则型

C. 肺大泡

D. 间质性肺气肿

E. 全小叶型

21. 肺气肿可导致

A. 肺源性心脏病

B. 自发性气胸

C. 肺性脑病

D. 筒状胸

E. 以上都是

22. 肺气肿的病变累及

A. 肺泡

B. 终末细支气管

C. 呼吸性细支气管

D. 肺泡管

E. 以上都是

23. 全腺泡型肺气肿的主要原因是

A. 小气道阻塞

B. 支气管壁炎细胞浸润

C. 肺组织炎细胞浸润

D. 先天性 α_1-抗胰蛋白酶缺乏

E. 支气管壁扩张

24. 女性，60 岁，慢性支气管炎近 10 年，近 3 年来感觉气短、胸闷、呼吸困难。查体：桶状胸，胸廓呼吸运动减弱；叩诊呈过清音，心浊音界缩小，肝浊音界下降；听诊呼吸音减弱。不符合该患者疾病的描述是

A. 肺泡扩张

B. 肺泡间隔变窄

C. 肺泡融合成囊泡状

D. 肺间质广泛纤维化

E. 肺组织膨胀

25. 下列疾病中，最常引起肺心病者是

A. 肺动脉栓塞

B. 胸廓畸形

C. 矽肺

D. 慢性肺结核

E. 肺气肿

26. 诊断肺心病的主要标准是

A. 心脏增大

B. 心尖钝圆

C. 右心室扩张

D. 肺动脉圆锥膨隆

E. 肺动脉下 2cm 处右心室壁厚超过 5mm

27. 肺心病的心肌病变不包括

A. 心肌细胞肥大，核大深染

B. 心肌细胞增生，数目增多

C. 心肌纤维萎缩、变细

D. 心肌细胞肌浆溶解

E. 心肌间质纤维化

28. 肺心病时，肺内小动脉的变化不包括

A. 小动脉炎

B. 小动脉玻璃样变

C. 无肌性细动脉肌化

D. 小动脉弹力纤维增生

E. 肌型小动脉中膜肥厚

29. 引起肺心病的关键环节是

A. 小气道阻塞

B. 肺部感染

C. 肺动脉高压

D. 肺间质纤维组织增生
E. 肺水肿

30. 肺心病可引起
A. 呼吸衰竭
B. 心力衰竭
C. 肺性脑病
D. DIC
E. 以上都是

31. 慢性肺源性心脏病最常见的原因是
A. 矽肺
B. 硬变性肺结核
C. 弥漫性肺间质纤维化
D. 慢性阻塞性肺气肿
E. 结节性多动脉炎

32. 引起急性肺源性心脏病的常见原因是
A. 肺脂肪栓塞
B. 肺空气栓塞
C. 肺总动脉主干血栓栓塞
D. 肺动脉小分支血栓栓塞
E. 肺动脉分支肿瘤栓塞

33. 不符合支气管扩张特点的是
A. 多发生于叶及段等大支气管
B. 支气管壁的炎症损伤是主要发病基础
C. 肺脓肿为其常见合并症
D. 可导致肺心病
E. 临床表现咳大量脓痰及反复咯血

34. 支气管扩张常合并
A. 肺肉质变
B. 弥漫性肺气肿
C. 肺结核
D. 肺脓肿
E. 早期肺门淋巴结病变

35. 支气管扩张的发病基础是
A. 支气管的炎症损伤和管腔阻塞
B. 支气管旁淋巴组织增生
C. 支气管发育异常
D. 肺免疫功能障碍
E. 肺内胰蛋白酶水平增高

36. 支气管扩张的病变不包括
A. 支气管呈节段性囊性扩张
B. 囊内为脓性渗出物
C. 支气管壁结构破坏
D. 支气管壁多量炎细胞浸润
E. 支气管上皮呈乳头状增生

37. 肺炎是指
A. 各种致炎因子引起的肺的炎症
B. 各种细菌引起的肺的炎症
C. 各种原因引起的肺泡的炎症
D. 各种原因引起肺的急性渗出性炎症
E. 以中性白细胞渗出为主的肺的炎症

38. 肺炎中最常见的病原体是
A. 病毒
B. 衣原体
C. 支原体
D. 细菌
E. 真菌

39. 在肺炎的诊治中，最重要的是
A. 确定炎症的部位
B. 确定炎症的类型
C. 确定炎症的分布特点
D. 确定炎症的严重程度和病原体
E. 确定感染的途径

40. 医院内肺炎常见的病原体是

A. 肺炎链球菌等多见
B. 绿脓杆菌及其他假单胞菌等多见
C. 厌氧菌多见
D. 病毒多见
E. 支原体多见

41. 不符合小叶性肺炎的是
A. 由多种致病菌引起
B. 常为其他疾病的合并症
C. 病变呈灶状、小叶分布
D. 病变为纤维素性炎
E. 可导致支气管扩张

42. 小叶性肺炎的典型病变是
A. 支气管急性卡他性炎
B. 以细支气管为中心的小叶性化脓性炎
C. 肺泡的纤维素性炎
D. 肺泡出血性炎
E. 肺间质非化脓性炎

43. 不符合小叶性肺炎的是
A. 主要由病毒引起
B. 小叶和小叶融合性病变
C. 病变以肺的背部和下部较重
D. 可导致心力衰竭
E. 可引起肺脓肿

44. 不符合小叶性肺炎的是
A. 由多种细菌引起
B. 病变从细支气管开始
C. 属化脓性炎症
D. 常咳铁锈色痰
E. 可破坏肺泡壁

45. 小叶性肺炎一般不引起
A. 肺肉质变
B. 脓胸
C. 脓毒败血症
D. 支气管扩张
E. 肺脓肿

46. 不属于小叶性肺炎的是
A. 麻疹后肺炎
B. 病毒性肺炎
C. 手术后肺炎
D. 吸入性肺炎
E. 坠积性肺炎

47. 男性，80 岁，直肠癌根治术后 2 周。7 天来发热（39 ℃）、咳嗽、咳粘液脓痰。查体：双肺下叶背侧可闻及湿罗音。X 线见双肺下叶散在不规则斑片状模糊阴影。符合该患者疾病的描述是
A. 肺泡纤维素性炎
B. 支气管肺泡化脓性炎
C. 间质性肺炎
D. 病毒性肺炎
E. 支原体性肺炎

48. 男性，80 岁，直肠癌根治术后 2 周。7 天来发热（39 ℃）、咳嗽、咳粘液脓痰。查体：双肺下叶背侧可闻及湿罗音。X 线见双肺下叶散在不规则斑片状模糊阴影。不符合该患者疾病的合并症的描述是
A. 中毒性休克
B. 肺脓肿
C. 脓胸
D. 肺栓塞
E. 心力衰竭

49. 大叶性肺炎的病变属于

A. 化脓性炎
B. 出血性炎
C. 纤维素性炎
D. 浆液性炎
E. 肉芽肿性炎

50. 大叶性肺炎发生肺肉质变是因为
A. 局部组织缺血
B. 纤维素渗出过多
C. 机体抵抗力降低
D. 吞噬细胞数量少或功能障碍
E. 合并其他细菌感染

51. 不符合大叶性肺炎的描述是
A. 多由肺炎链球菌感染引起
B. 累及整个或多个大叶
C. 属于纤维素性炎
D. 破坏小支气管壁和肺泡壁结构
E. 咳铁锈色痰

52. 大叶性肺炎的合并症有
A. 肺肉质变
B. 感染性休克
C. 纤维素性胸膜炎
D. 肺脓肿
E. 以上都是

53. 大叶性肺炎合并金黄色葡萄球菌感染可引起
A. 败血症
B. 肺脓肿
C. 脓胸
D. 感染性休克
E. 以上都是

54. 不符合大叶性肺炎病理变化的描述是
A. 肺泡腔内浆液渗出
B. 肺泡腔内纤维素渗出
C. 肺泡腔内可有红细胞
D. 肺泡腔内中性粒细胞渗出
E. 肺泡腔内多量心力衰竭细胞

55. 男性，30 岁，2 天前上呼吸道感染，病情迅速加重，发热（39.8℃），伴寒战、胸痛、咳嗽、咳铁锈色痰。X 线检查见左肺下叶大片致密阴影。符合该患者疾病的描述是
A. 大叶性肺炎
B. 小叶性肺炎
C. 间质性肺炎
D. 病毒性肺炎
E. 支原体性肺炎

56. 男性，30 岁，2 天前上呼吸道感染，病情迅速加重，发热（39.8℃），伴寒战、胸痛、咳嗽、咳铁锈色痰。X 线检查见左肺下叶大片致密阴影。不符合该患者疾病的描述是
A. 可合并中毒性休克
B. 可合并肺脓肿
C. 可合并肺肉质变
D. 可合并支气管扩张
E. 可合并胸膜炎

57. 病毒性肺炎的特点中不包括
A. 多由上呼吸道病毒感染蔓延引起
B. 炎症主要累及肺间质
C. 肺泡内可有浆液渗出
D. 肺组织可出现片状坏死
E. 细胞内可出现包涵体

58. 病毒性肺炎具有诊断价值的组织学特点是

A. 肺间质中单核细胞、淋巴细胞浸润
B. 肺泡腔内浆液渗出
C. 肺泡上皮细胞增生
D. 胞浆内和胞核内出现包涵体
E. 细支气管上皮增生

59. 麻疹病毒性肺炎的特点是
A. 间质性肺炎
B. 肺泡上皮细胞增生
C. 形成核内包涵体
D. 形成胞浆内包涵体
E. 形成核内包涵体，也形成胞浆内包涵体

60. 病毒性肺炎最常见的病原体是
A. 腺病毒
B. 巨细胞病毒
C. 流感病毒
D. 副流感病毒
E. 鼻病毒

61. 病毒性肺炎中，不形成病毒包涵体的是
A. 腺病毒
B. 麻疹病毒
C. 巨细胞病毒
D. 呼吸道合胞病毒
E. 流感病毒

62. 病毒性肺炎的肉眼特点是
A. 大叶性病变
B. 小叶性病变
C. 斑片状病变
D. 小叶融合性病变
E. 结节状病变

63. 巨细胞病毒、单纯疱疹病毒引起的肺炎多见于
A. 青壮年
B. 儿童
C. 老年人
D. 艾滋病患者
E. 妇女

64. 病毒性肺炎的确诊主要依据是
A. X 线检查
B. 血清学检查
C. 病毒和病毒抗原的检查
D. 病毒感染流行状况
E. 儿童和老年人

65. 确诊支原体肺炎的依据是
A. 患者多为儿童和青年
B. 起病急，多有发热、头痛、咽痛及剧烈干性呛咳
C. X 线检查，肺部呈段性分布的纹理增加及网状阴影
D. 病变呈间质性炎症，肺泡腔内可无渗出物
E. 以上都不是

66. 不符合原发性非典型性肺炎的描述是
A. 可由病毒引起
B. 可由支原体引起
C. 可由衣原体引起
D. 可由细菌引起
E. 可呈间质性肺炎

67. 原发性非典型性肺炎的病变特点是
A. 大叶性病变
B. 小叶性病变
C. 斑片状病变
D. 小叶融合性病变
E. 结节状病变

68. 原发性非典型性肺炎的病变特点是
A. 肺纤维素性炎
B. 肺化脓性炎
C. 肺浆液性炎
D. 肺出血性炎
E. 肺间质性炎

69. 确诊支原体肺炎的可靠依据是
A. X 线检查
B. 血清学检查
C. 病原学检查
D. 支原体感染流行状况
E. 排除细菌性肺炎的可能

70. 不符合支原体肺炎的描述是
A. 肺炎支原体不直接破坏肺组织
B. 多有发热、头痛、咽痛及剧烈干性呛咳
C. 病变呈斑片状阴影
D. 肺间质内淋巴单核细胞浸润
E. 肺泡不受累

71. 符合支原体肺炎的描述是
A. 可合并中毒性休克
B. 可合并肺脓肿
C. 可合并肺肉质变
D. 可合并急性支气管炎
E. 可合并胸膜炎

72. 肺炎支原体的主要致病作用是
A. 毒素
B. 对肺组织的直接破坏作用
C. 引起过敏反应
D. 损伤 T 细胞
E. 损伤 B 细胞

73. 军团菌肺炎的特点是
A. 大叶性病变
B. 小叶性病变
C. 斑片状病变
D. 小叶融合性病变
E. 小叶性病变、小叶融合性病变或大叶性病变

74. 诊断军团菌肺炎的确切依据是
A. X 线检查
B. 尿液内抗军团菌抗体检查
C. 支气管、肺冲洗液病原学检查
D. 军团菌感染流行状况
E. 病情危重

75. 不符合军团菌肺炎特点的是
A. 可呈散发性
B. 可呈流行性
C. 病情急、死亡率高
D. 红霉素治疗效果良好
E. 呈大叶性病变

76. 不符合金黄色葡萄球菌肺炎特点的是
A. 多呈大叶性病变
B. 多呈小叶性病变
C. 多见于医院内肺炎
D. 对青霉素有抗药性
E. 多引起肺脓肿

77. 符合克雷白杆菌肺炎特点的是
A. 咳铁锈色痰
B. 咳黄色脓痰
C. 咳淡绿色痰
D. 咳臭味痰
E. 咳粘冻样痰

78. 不符合绿脓杆菌肺炎特点的是
A. 易引起败血症

B. 多呈小叶性病变
C. 多见于社区性肺炎
D. 多发生于严重烧伤、白血病、化学治疗患者
E. 多引起肺脓肿

79. 不符合流感嗜血杆菌肺炎特点的是
A. 起病隐匿
B. 多呈小叶性病变
C. 多发生于慢性支气管炎和支气管扩张患者
D. 病情较轻
E. 多引起肺脓肿

80. 不符合吸入性肺炎特点的是
A. 多呈大叶性病变
B. 多呈小叶性病变
C. 多见于医院内肺炎
D. 多发生于危重病人
E. 多引起肺脓肿

81. 引起矽肺最主要的是
A. 10μm 的矽尘颗粒
B. 8μm 的矽尘颗粒
C. 6μm 的矽尘颗粒
D. 4μm 的矽尘颗粒
E. 1～2μm 的矽尘颗粒

82. 不符合矽肺的病理变化的是
A. 矽结节形成
B. 肺纤维组织弥漫增生
C. 肺纤维化具有病理诊断意义
D. 晚期矽结节可融合成团
E. 早期矽结节由巨噬细胞构成

83. 不符合矽肺合并症的是
A. 肺癌
B. 肺心病
C. 肺气肿
D. 自发性气胸
E. 矽肺结核病

84. 不符合矽肺各期的病变特点的是
A. 早期病变累及周围肺组织
B. 肺门淋巴结常受累
C. I 期矽肺累及肺门部
D. II 期矽肺累及全肺 1/3 以下
E. III 期矽肺累及肺的大部分，矽结节密集融合成块

85. 符合矽结节的描述是
A. 早期是纤维结节
B. 晚期是细胞结节
C. 其发生与 T 淋巴细胞关系密切
D. 具有病理诊断意义
E. 不含免疫球蛋白

86. 不符合矽肺发病特点的是
A. 免疫反应参与矽肺发病
B. 游离的二氧化硅具有细胞毒性
C. 纤维化与巨噬细胞分泌的细胞因子有关
D. 较大的矽尘颗粒致病作用更强
E. 病变首先累及淋巴系统

87. 不符合矽肺结核病的描述是
A. 矽肺结核性空洞数目多，直径大
B. 易引起大咯血
C. 易发生于矽肺晚期
D. 病变发展缓慢
E. 与肺组织的防御能力下降有关

88. 男性，60 岁，煤矿工人，近 10 年来出现憋气、心悸、口唇青紫。X 片检查，

肺门部可见较密集的结节，多数直径小于 1cm。符合该患者疾病描述的是

A. 煤矽肺
B. 肺结核
C. 肺癌
D. 结节病
E. 肺炎性假瘤

89. 男性，60 岁，煤矿工人，近 10 年来出现憋气、心悸、口唇青紫。X 片检查，肺门部可见较密集的结节，多数直径小于 1cm。不符合该患者疾病描述的是

A. 可引起呼吸衰竭
B. 可合并肺结核
C. 可合并肺源性心脏病
D. 可合并肺气肿
E. 可合并间皮瘤

90. 男性，60 岁，煤矿工人，近 10 年来出现憋气、心悸、口唇青紫。X 片检查，肺门部可见较密集的结节，多数直径小于 1cm。不符合该患者疾病描述的是

A. 矽结节对本病有诊断意义
B. 大于 5μm 的矽粉尘最有致病性
C. 肺间质弥漫性纤维化
D. II 期矽肺矽结节直径小于 1cm
E. III 期矽肺矽结节直径大于 2cm

91. 晚期矽肺脏层胸膜下可出现

A. 腺泡中央型肺气肿
B. 全腺泡型肺气肿
C. 腺泡周围型肺气肿
D. 不规则型肺气肿
E. 间质性肺气肿

92. 石棉肺的病理诊断依据是

A. 弥漫性纤维化的肺组织内可见石棉小体
B. 壁层胸膜斑
C. 肺间质纤维化
D. 肺脱屑性间质性肺炎
E. 肺气肿

93. 不符合石棉肺合并症的描述是

A. 可引起肺脓肿
B. 可引起间皮瘤
C. 可引起肺心病
D. 可引起肺气肿
E. 可引起肺结核

94. 石棉小体是

A. 肉芽肿性病变
B. 呈同心圆排列的纤维结节
C. 胸膜纤维瘢痕
D. 有铁蛋白沉积的石棉纤维
E. 巨噬细胞构成的小体

95. 不符合农民肺的描述是

A. 由于吸入高温放线菌引起的过敏性肺泡炎
B. 肺内出现肉芽肿病变
C. 类上皮细胞内可见放线菌孢子
D. 肺间质纤维组织增生
E. 属 I 型超敏反应性疾病

96. 石棉粉尘能引起

A. 脱屑性间质性肺炎
B. 巨细胞肺炎
C. 大叶性肺炎
D. 过敏性肺泡炎
E. 小叶性肺炎

97. 不符合鼻咽癌特征的是

A. 多起源于鼻咽部柱状上皮
B. 多发生于鼻咽顶部
C. 癌组织浸润能力强
D. 可破坏颅底骨向颅内浸润
E. 晚期方经淋巴道及血道转移

98. 最常见的鼻咽癌的组织学类型是
A. 未分化癌
B. 泡状核细胞癌
C. 腺泡状腺癌
D. 低分化鳞状细胞癌
E. 高分化鳞状细胞癌

99. 在我国，头颈部恶性肿瘤中发病率最高的是
A. 恶性淋巴瘤
B. 喉癌
C. 扁桃体癌
D. 基底细胞癌
E. 鼻咽癌

100. 鼻咽癌的病因中，最重要的发病因素是
A. 遗传因素
B. 亚硝胺类
C. 多环芳烃类
D. 激素
E. EB 病毒

101. 鼻咽癌的辅助诊断中，常用的检验指标是
A. CEA
B. AFP
C. CA19-9
D. 酸性磷酸酶
E. 抗 EB 病毒壳抗原的癌基因抗体

102. 泡状核细胞癌的特点是
A. 出现角化珠
B. 形成腺体
C. 形成癌巢
D. 核大呈空泡状
E. 对放射治疗敏感

103. 鼻咽癌的显著特点是
A. 早期症状明显
B. 可引起脑神经破坏
C. 早期发生淋巴道转移
D. 可引起耳鸣、鼻堵
E. 可有家族聚集倾向

104. 鼻咽癌最常见的肉眼类型是
A. 溃疡型
B. 菜花型
C. 粘膜下型
D. 浸润型
E. 结节型

105. 不易发生肺真菌病的患者是
A. 慢性消耗性疾病患者
B. 长期使用广谱抗生素患者
C. 长期使用肾上腺皮质激素患者
D. 接受放射治疗、化学治疗、免疫抑制剂的患者
E. 静脉输液患者

106. 肺真菌病的确诊主要依据
A. 患者免疫功能低下
B. 血清学检查
C. X 线检查
D. 真菌检查或在病变中查到真菌
E. 炎症的病理学类型

107. 不符合肺孢子虫性肺炎的描述是
A. 仅发生于艾滋病患者

B. 可引起肺实变
C. 可引起间质性肺炎
D. 肺泡腔内可见泡沫状蛋白
E. 肺活检组织内可查到滋养体

108. 不符合巨细胞病毒性肺炎的描述是
A. 成人多见于艾滋病患者和免疫功能低下患者
B. 可合并胃肠道、中枢神经和视网膜感染
C. 可引起呼吸窘迫综合征
D. 可见巨细胞形成和巨大的核内包涵体
E. 正常人群仅有少数人抗巨细胞病毒抗体阳性

109. 不符合念珠菌病的描述是
A. 是常见的深部真菌病
B. 可累及消化道、呼吸道、泌尿道、心脏和脑
C. 可形成小脓肿和肉芽肿
D. 芽生孢子和假菌丝是主要诊断依据
E. 多为外源性感染

110. 不符合隐球病的描述是
A. 主要通过呼吸道感染
B. 病原体为芽生酵母菌
C. 以肺部病变多见
D. 可引起脑膜炎
E. 可形成肉芽肿

111. 不符合曲菌病的描述是
A. 主要通过呼吸道传染
B. 以肺部病变多见
C. 可形成脓肿
D. 可引起过敏性支气管炎
E. 菌丝有隔，呈直角分支

112. 不符合放线菌病的描述是
A. 主要通过外源性感染
B. 常引起慢性化脓性炎
C. 脓液中可见“硫磺颗粒”
D. 蓝紫色分支状菌丝放射状排列
E. 病变可累及颌面部、胸部和腹部

113. 女性，60 岁，40 年吸烟史。慢性咳嗽 30 年，近 1 月来痰中带血丝。X 线检查，显示左肺门处不规则巨大阴影。支气管镜检查，左主支气管粘膜不平，管腔狭窄。活检病理诊断最可能是
A. 肺结核病
B. 肺结节病
C. 细支气管肺泡细胞癌
D. 肺鳞状细胞癌
E. 肺炎性假瘤

114. 女性，50 岁，本人不吸烟，但家中多个成员吸烟。查体偶然发现右肺近胸膜处有一直径 3cm 的边界较清的阴影，伴肺门淋巴结肿大。该结节最可能是
A. 鳞状细胞癌
B. 细支气管肺泡细胞癌
C. 腺癌
D. 大细胞癌
E. 小细胞癌

115. 男性，60 岁，3 个月来咳嗽、胸痛。X 线检查，左肺下叶大片云雾状、不均匀密度增高阴影，经用抗生素治疗无效。痰脱落细胞检查，可见腺癌细胞。对该患者疾病的诊断是
A. 鳞状细胞癌

B. 类癌
C. 细支气管肺泡细胞癌
D. 小细胞癌
E. 巨细胞癌

116. 肺癌的最常见肉眼类型是
A. 中央型
B. 周围型
C. 弥漫型
D. 管内型
E. 管壁浸润型

117. 弥漫型肺癌多为
A. 大细胞癌
B. 鳞状细胞癌
C. 小细胞癌
D. 细支气管肺泡细胞癌
E. 类癌

118. 不符合肺腺癌的描述是
A. 多为周围型
B. 常有吸烟史
C. 可伴有粘液分泌
D. 常有肺门淋巴结转移
E. 常有乳头形成

119. 不符合肺癌的特点的是
A. 发病隐匿
B. 生长迅速，转移较早
C. 易发生脑、骨及肾上腺转移
D. 可发生种植转移
E. 不引起内分泌紊乱

120. 肺癌不会引起
A. Horner 综合征
B. Cushing 综合征
C. 类癌综合征
D. 甲状腺机能亢进
E. 骨关节病

121. 肺癌的肉眼形态不包括
A. 肺门部巨大肿块
B. 肺膜下的球形结节
C. 散布于两肺的多发性结节
D. 大叶性实性病变
E. 小叶融合性实性病变

122. 肺癌的主要组织来源是
A. 支气管上皮细胞
B. Clara 细胞
C. Kultschitzky 细胞
D. I 型肺泡上皮细胞
E. II 性肺泡上皮细胞

123. 最容易引起的皮肤潮红、水泻、心动过速、支气管哮喘等症状的肺癌类型是
A. 低分化鳞状细胞癌
B. 未分化腺癌
C. 大细胞癌
D. 类癌
E. 粘液癌

124. 累及整个大叶呈肺炎样外观的肺癌类型是
A. 低分化鳞状细胞癌
B. 大细胞癌
C. 小细胞癌
D. 粘液癌
E. 细支气管肺泡细胞癌

125. 可能来源与 II 型肺泡上皮细胞的肺癌是
A. 腺癌
B. 瘢痕癌

C. 小细胞癌
D. 细支气管肺泡细胞癌
E. 腺样囊性癌

126. 不符合隐性肺癌的描述是
A. 临床检查无明显肺癌体征
B. X 线检查未见阴影
C. 病理证实为原位癌或早期侵润性癌
D. 无淋巴结转移
E. 痰细胞学检查癌细胞阴性

127. 不符合小细胞肺癌特点的描述是
A. 多发生于肺周边部
B. 细胞小、密集排列，由结缔组织加以分隔
C. 分化程度低、恶性度高
D. 肿瘤细胞胞浆内出现神经内分泌颗粒
E. 起源于嗜银细胞

128. 腺癌起源于
A. 主支气管粘膜上皮细胞
B. 支气管腺体
C. 小支气管粘膜上皮细胞
D. 嗜银细胞
E. II 型肺泡上皮细胞

129. 不符合急性鼻炎的描述是
A. 由各种呼吸道病毒引起
B. 可呈肉芽肿性炎
C. 可由各种变应原引起
D. 可呈浆液性炎
E. 可呈化脓性炎

130. “臭鼻症”指
A. 急性过敏性鼻炎
B. 慢性肥厚性鼻炎
C. 慢性萎缩性鼻炎
D. 急性化脓性鼻炎
E. 急性浆液性鼻炎

131. 鼻腔恶性肉芽肿实质上是指
A. 以巨噬细胞增生为主的炎性肉芽肿
B. Wegener 肉芽肿
C. 血管中心型非霍奇金淋巴瘤
D. 致死性中线肉芽肿
E. 中线恶性网状细胞增生症

132. 符合鼻腔内翻性乳头状瘤的特点的是
A. 慢性炎症性增生
B. 不易恶变
C. 鳞状上皮团向间质内生长
D. 与 HPV 感染无关
E. 术后不复发

133. 不符合扁桃体炎的描述是
A. 多由乙型溶血性链球菌引起
B. 可呈急性化脓性扁桃体炎
C. 可呈慢性肥大性扁桃体炎
D. 可呈慢性纤维性扁桃体炎
E. 中老年患者多见

134. 不符合鼻息肉的特点的是
A. 良性肿瘤
B. 肉眼呈灰白色
C. 间质水肿
D. 腺体增生，并可呈囊状扩张
E. 间质炎性细胞侵润

135. 不符合鼻窦炎的特点的是
A. 以上颌窦炎多见
B. 可呈浆液性炎
C. 可呈化脓性炎
D. 可呈慢性炎

E. 多由血源性细菌引起

136. 不符合咽炎的特点的是

A. 多由葡萄球菌引起

B. 可呈纤维素性炎

C. 可呈急性化脓性炎

D. 可呈慢性肥厚性炎

E. 可呈慢性萎缩性炎

137. 不符合特发性肺间质纤维化的描述是

A. 可能是一种自身免疫性疾病

B. 早期表现为脱屑性间质性肺炎

C. 晚期可形成蜂窝肺

D. 蜂窝肺小囊腔由扩张的细支气管和融合的肺泡构成

E. 蜂窝肺的囊壁由肺泡壁构成

138. 不符合特发性肺间质纤维化合并症的是

A. 肺气肿

B. 肺心病

C. 肺炎

D. 肺癌

E. 间皮瘤

139. 符合特发性肺间质纤维化病因的描述是

A. 是一种自身免疫性疾病

B. 是一种遗传病

C. 是一种病毒感染性疾病

D. 是一种原因不明的疾病

E. 是一种职业病

140. 蜂窝肺是

A. 大叶性肺炎引起的肺肉质变

B. 阻塞性肺气肿引起的肺病变

C. 特发性肺间质纤维化引起的肺病变

D. 矽肺引起的肺病变

E. 农民肺引起的肺病变

141. 不符合成人呼吸窘迫症的描述是

A. 是一种肺的原发性疾病

B. 表现为突发性进行性呼吸困难

C. 肺泡毛细血管膜弥漫性损伤

D. 病死率高达 36%～68%

E. X 线检查两肺弥漫性浸润性阴影

142. 不符合成人呼吸窘迫症的病因是

A. 严重创伤

B. 严重感染

C. 休克

D. 中毒

E. 晚期癌症

143. 成人呼吸窘迫症的发病机制的中心环节是

A. 肺内淋巴细胞渗出

B. 肺内单核细胞渗出

C. 肺泡毛细血管膜弥漫性损伤

D. 肺毛细血管内血小板聚集

E. 肺表面活性物质减少

144. 成人呼吸窘迫症的特征性病理变化是

A. 肺水肿

B. 透明膜形成

C. 肺出血

D. 肺纤维化

E. 肺萎陷

145. 不符合喉癌的描述是

A. 腺癌少见

B. 声门上癌转移率高

C. 声带癌转移率高

D. 声带癌最多见

E. 鳞癌最多见

146. 不符合喉炎特点的描述是

A. 儿童急性喉炎可引起哮吼，甚至窒息

B. 声带小结与长期用声过度或用声不当有关

C. 慢性喉炎与吸烟关系密切

D. 急性喉炎与病毒、细菌感染关系密切

E. 声带小结易恶变，是一种癌前病变

147. 肺脓肿最常见的病原体是

A. 厌氧菌

B. 需氧菌

C. 革兰氏阴性菌

D. 革兰氏阳性菌

E. 兼性厌氧菌

148. 吸入性肺脓肿易发生于

A. 左上肺

B. 左下肺

C. 右上肺

D. 右上肺腋亚段和右下肺上部

E. 右下肺

149. 不符合肺脓肿的描述是

A. 吸入性肺脓肿常为多发

B. 肺炎和支气管扩张引起的肺脓肿常为多发

C. 可引起脓胸、脓气胸

D. 可引起败血症

E. 可引起支气管胸膜瘘

150. 符合癌性胸水的描述是

A. 比重大于 1.020

B. 可见多量淋巴细胞

C. 可见多量中性粒细胞

D. 可见肿瘤细胞

E. 可见多量间皮细胞

151. 引起胸膜漏出的疾病是

A. 肺癌

B. 肺梗死

C. 肺结核

D. 肺脓肿

E. 肺淤血

152. 男性，50 岁，近一月来感到胸部胀闷，近 3 天伴有气促、心慌。X 线检查，左胸腔大量积液，肺门部可见 5×3cm 大小的阴影。查体：患者无发热，结核菌实验阴性。该患者胸水最可能为

A. 炎性渗出液

B. 癌性胸水

C. 胸腔积血

D. 胸腔积脓

E. 乳糜液

153. emphysema

A. 肺气肿

B. 肺梗死

C. 肺结核

D. 肺脓肿

E. 肺淤血

154. bronchopneumonia

A. 大叶性肺炎

B. 支气管肺炎

C. 间质性肺炎

D. 腺泡中央型肺气肿

E. 全腺泡型肺气肿

155. nasopharyngeal carcinoma

A. 肺癌
B. 鼻咽癌
C. 喉癌
D. 支气管癌
E. 肺泡细胞癌

156. asbestosis
A. 农民肺
B. 矽肺
C. 煤矽肺
D. 尘肺
E. 石棉肺

157. chronic cor pulmonale
A. 慢性阻塞性肺气肿
B. 慢性支气管炎
C. 支气管扩张症
D. 慢性肺纤维化
E. 慢性肺源性心脏病

158. respiratory distress syndrome
A. 特发性肺间质纤维化
B. 呼吸窘迫综合征
C. 肺副肿瘤综合征
D. 上腔静脉综合征
E. 交感神经麻痹综合征

159. actinomycosis
A. 念珠菌病
B. 组织胞浆菌病
C. 曲菌病
D. 放线菌病
E. 球孢子菌病

160. mesothelioma
A. 胸膜炎
B. 气胸
C. 血胸
D. 间皮瘤
E. 错构瘤

英文题

1. A 70-year-old woman is referred to an ophthalmologist because of difficulty with her right eye. She also has pain in the right upper chest. The findings on physical examination include enophthalmos, meiosis, anhidrosis, and ptosis. A chest radiograph shows right upper lobe opacification and bony destruction of the right first rib. Which of the following conditions is most likely to be present?
A. Bronchopneumonia
B. Bronchiectasis
C. Bronchogenic carcinoma
D. Sarcoidosis
E. Tuberculosis

2. Which of the following is the most frequent site of intestinal tuberculosis
A. Ileocecum
B. Jejunum
C. Descending colon
D. Duodenum
E. Ascending colon

3. Which of the following is not present in secondary pulmonary tuberculosis?

A. The apex of the lung is the most common site of lesions
B. Fibro-cavities may form
C. The course of the disease is long and the state of the illness is sometimes good and sometimes bad
D. Old and new lesions are mixed
E. It mainly spreads through the lymphatics

4. Which of the following descriptions of secondary tuberculosis is appropriate?
A. Old and new lesions are mixed and the lesions of the upper lung are more severe
B. There are obvious lesion of tuberculosis in the lymph nodes of the hilus of lung
C. It does not readily cause cavitation
D. The vast majority of patients recover naturally
E. It spreads easily through lymph vessels and blood vessels

5. Which of the following is characteristic of chronic fibro-cavitative pulmonary tuberculosis?
A. The cavities are the same size and regular shape
B. There is formation of one or a few acute cavities in the lung
C. The beam-like tissue frequently seen in the cavity is the remaining branch of the bronchus
D. There is caseous necrosis in the inner wall of the cavity containing a large number of mycobacterium tuberculosis
E. The cavity does not communicate with the bronchus

6. Which of the following is not characteristic of infiltrative pulmonary tuberculosis?
A. The lesion may be extensive
B. There is a small amount of caseous necrosis in the center of the lesion
C. The alveolar cavity is full of serous fluid, mononuclear cells and lymphocytes microscopically
D. The alveolar cavity is filled with a large number of neutrophils
E. Acute cavity formation may be seen

7. Which of the following is the definition of tuberculoma?
A. It is a benign tumor caused by mycobacterium tuberculosis
B. It is a malignant tumor caused by mycobacterium tuberculosis
C. It is a tumor-like invasive lesion caused by mycobacterium tuberculosis
D. It is a solitary focus of caseous necrosis caused by mycobacterium tuberculosis
E. It is a cold abscess caused by mycobacterium tuberculosis

8. Which of the following is the most com-

mon type of adult pulmonary tuberculosis?

A. Focal pulmonary tuberculosis
B. Infiltrative pulmonary tuberculosis
C. Tuberculoma
D. Caseous pneumonia
E. Chronic fibro-cavitative pulmonary tuberculosis

9. Which of the following is not correct regarding secondary tuberculosis?

A. It is mainly caused by endogenic reinfection
B. It mainly disseminates through lymphatics and blood vessels
C. It mainly disseminates through air passages
D. It mostly starts from the apex of the lung
E. Old and new lesions are mixed together

10. Which of the following descriptions of tuberculosis is not appropriate?

A. Tuberculosis is a chronic infectious disease
B. Tuberculosis is attributed to type III allergic reaction
C. Extrapulmonary tuberculosis is mostly caused by blood born dissemination
D. Mycobacterium tuberculosis is still present in calcified necrotic foci
E. Cavitative pulmonary tuberculosis is an exoteric tuberculosis

11. Which of the following is caused by primary pulmonary tuberculosis?

A. Acute miliary tuberculosis
B. Focal pulmonary tuberculosis
C. Infiltrative pulmonary tuberculosis
D. Caseous pneumonia
E. Cavitative tuberculosis

12. Which of the following is the most frequent site of primary tuberculosis in the lung?

A. Hilum of lung
B. Apex of lung
C. Lower portion of upper lobe or upper portion of lower lobe of lung
D. Diaphragmatic surface of the lung
E. Pleura

13. Which of the following is the most frequent site of tuberculosis in the male reproductive system?

A. Seminal vesicle
B. Prostate
C. Spermatic ducts
D. Epididymis
E. Testicle

14. Which of the following is true of ulcerative-type intestinal tuberculosis?

A. The ulcers are round or oval-shaped and the edges are well demarcated
B. It mostly occurs in the hepatic flexure of the colon
C. The major axis of the ulcer is parallel

too the intestinal axis

D. Only a little serous exudation is seen at the bottom of ulcers

E. It mostly occurs in the splenic flexure of the colon

15. Which of the following is not true of tuberculosis of bone?

A. It is frequently seen in the upper thoracic vertebrae

B. It may form a sequestrum

C. It may form a cold abscess

D. It may form caseous necrosis

E. It often involves vertebrae, phalanges and long bones

16. Which of the following is not true of tuberculous meningitis?

A. There is a large amount of fibrinopurulent exudate at the base of the brain

B. It is caused mainly by blood born dissemination of mycobacterium tuberculosis

C. It is often a part of systemic miliary tuberculosis

D. The lesions at the base of the brain are the most obvious

E. It is mostly found in children

17. Which of the following is true of intestinal tuberculosis?

A. It may be present as primary tuberculosis

B. It mainly involves the jejunum

C. It may cause cancer

D. It can lead to the formation of longitudinal ulcers

E. It mainly involves ileum

18. Which of the following is the most frequent site of tuberculosis of the female reproductive system?

A. Cervix

B. Vagina

C. Endometrium

D. Fallopian tube

E. Ovary

19. Which of the following is true of small cell carcinoma of the lung?

A. It does not have a close relationship with smoking

B. It may be associated with secretion of ectopic hormones

C. It originates in metaplasia of epithelial cells

D. The 5-year survival rate of it is high

E. Among lung cancers its incidence is highest

20. Which of the following is not true of small cell carcinoma of the lung?

A. It occurs mostly in the periphery of the lung

B. The tumor cells are small with intensive arrays

C. Neuroendocrine granules can be seen in tumor cells

D. It is high-grade malignant tumor

E. It is not suitable for surgical resection,

but is sensitive to chemotherapy and radiotherapy

21. Which of the following is not true of squamous cell carcinoma of the lung?
    A. It often occurs in smokers
    B. The vast majority arises in bronchial squamous metaplasia
    C. It is mostly central type
    D. The tumor often produces ectopic hormones
    E. It is easily found by fiberoptic bronchoscopy
22. Which of the following types of lung cancer has the lowest degree of malignancy?
    A. Carcinoid tumor
    B. Adenocarcinoma
    C. Squamous cell carcinoma
    D. Large cell carcinoma
    E. Small cell carcinoma
23. 50-year-old woman is not a smoker, but many members of her family are smokers. A shadow of 3 cm in diameter with a clear border and hilar lymphadenopathy near the pleura of the right lung was discovered incidentally during a routine checkup. Which of the following is likely to be the correct diagnosis?
    A. Bronchioloalveolar carcinoma
    B. Squamous cell carcinoma
    C. Adenocarcinoma
    D. Carcinoid tumor
    E. Small cell carcinoma
24. Which of the following types of lung cancer can cause symptoms such as skin flushing, watery diarrhea, tachycardia and bronchial asthma?
    A. Differentiated adenocarcinoma
    B. Poorly differentiated squamous cell carcinoma
    C. Large cell carcinoma
    D. Carcinoid tumor
    E. Bronchioloalveolar carcinoma
25. Which of the following descriptions of lung cancer is correct?
    A. Adenocarcinoma is the most frequent
    B. Most small cell carcinoma is diffuse type
    C. Most patients with squamous cell carcinoma have a history of smoking
    D. Peripheral type lung cancer is mostly squamous cell carcinoma
    E. Bronchioloalveolar carcinoma is mostly central type
26. Which of the following is not caused by lung cancer?
    A. Cushing syndrome
    B. Horner's syndrome
    C. Carcinoid syndrome
    D. Hyperthyroidism
    E. Superior vena cava syndrome
27. 60-year-old man has suffered from a cough and chest pain for 3 months. A chest x-ray shows large shadows with cloud-like and irregular increasing density

in the left lower lobe. Treatment with antibiotics has no effect in this patient. Exfoliative cytology of the sputum reveals adenocarcinoma cells. Which of the following diagnoses is likely to be correct?

A. Carcinoid
B. Squamous cell carcinoma
C. Bronchioloalveolar carcinoma
D. Small cell carcinoma
E. Large cell carcinoma

28. 70-year-old man has a 40-year history of smoking. He has suffered from a chronic cough for 30 years, and has been coughing bloody sputum for nearly 1 month. A chest x-ray shows irregular huge shadow in left hilum of the lung. Bronchoscopy shows irregular mucosa in the left main stem bronchus and narrowing of the lumen. Which of the following diagnoses is likely to be correct?

A. Pulmonary sarcoidosis
B. Tuberculosis
C. Bronchioloalveolar carcinoma
D. Squamous cell carcinoma of the lung
E. Pulmonary adenocarcinoma

29. Which of the following types of lung cancer is likely to involve the whole lobe and show a pneumonia-like appearance?

A. Bronchioloalveolar carcinoma
B. Large cell carcinoma
C. Small cell carcinoma
D. Mucinous adenocarcinoma
E. Signet-ring cell carcinoma

30. Which of the following is not true of silicosis?

A. It can easily be complicated by tuberculosis
B. It can cause pulmonary fibrosis
C. Early silicon nodules are present in the alveolus
D. Necrosis can be seen in the center of silicon nodules
E. Silicon nodules are composed of concentric ring-like collagen fibers

31. Which of the following is true regarding silicosis?

A. Silicon granules greater than 5μm in diameter are strongly pathogenic
B. Silicic acid can result in the injury of macrophages
C. There is no immunoglobulin in silicon nodules
D. The fibrous nodule is an early lesion
E. There are no lesions of the pleura

32. Which of the following description is not consistent with silicotuberculosis?

A. It can easily cause massive hemoptysis
B. The tubercular cavities of silicosis are typically greater in number and of larger diameter
C. It is likely to occur in late stages of silicosis
D. The disease is slowly progressive

E. The frequency of complication by tuberculosis parallels the severity of the silicotic lesions

33. Which of the following is true of pulmonary disease?
    A. Silicosis is caused mainly by silicon granules less than 5μm in diameter
    B. Lobular pneumonia is caused mainly by mycoplasma
    C. α_1-antitrypsin deficiency is the most common cause of emphysema
    D. Bronchiectasis means expansion of the bronchial cavity
    E. Bronchopneumonia often occurs as an independent disease process
34. 60-year-old man who works in a coal mine presents with pulmonary distress, palpitation and cyanosis of his lips. A chest x-ray reveals extensive nodules in the pulmonary hilus, the majority of which are less than 1 cm in diameter. Which of the following is not true of this disease?
    A. Mesothelioma is a possible complication
    B. Pulmonary tuberculosis is a possible complication
    C. Pulmonary heart disease is a possible complication
    D. Emphysema is a possible complication
    E. Chronic bronchitis is a possible complication
35. 40-year-old man who works in a coal mine presents with respiratory distress, palpitation and cyanosis of his lips. A chest x-ray reveals extensive nodules in the hilus of the lung, the majority of which are less than 1 cm in diameter. Which of the following is not consistent with this patient's disease?
    A. Silicon nodules are the characteristic lesion of this disease
    B. Silicon granules greater than 5μm are the most pathogenic
    C. This is silicosis, phase II
    D. Diffuse interstitial fibrosis of lung
    E. May lead to pulmonary heart disease
36. Which of the following results from silicosis and leads to pulmonary heart disease?
    A. Pulmonary cavity formation due to silicosis
    B. Secondary infection of the lung
    C. Diffuse interstitial fibrosis of the lung
    D. Silicotuberculosis
    E. Chronic bronchitis
37. Which of the following lesions is likely to be caused by silicon particles?
    A. Suppurative inflammation of the lung
    B. Fibrinous inflammation of the lung
    C. Pulmonary fibrosis
    D. Interstitial pneumonia
    E. Serous inflammation of the lung
38. Which of the following cells are present in

early silicotic nodules?

A. large number of macrophages
B. A large number of eosinophils
C. A large number of plasma cells
D. A large number of fibroblasts
E. A large number of neutrophils

39. Which of the following is the main pathogenesis of bronchiectasis?

A. Chronic bronchial infection
B. Pulmonary ventilation barrier due to bronchial obstruction
C. Obstruction by bronchial tumo
D. Destruction of bronchial smooth muscle and elastic tissue
E. Squamous metaplasia of bronchial mucosa

40. Which of the following is characteristic of viral pneumonia?

A. Fibrinous pneumonia
B. Suppurative pneumonia
C. Hemorrhagic pneumonia
D. Pulmonary fibrosis
E. Interstitial pneumonia

41. Which of the following groups of findings is strongly characteristic of interstitial pneumonia?

A. Interstitial congestion and edema, infiltration of mononuclear cells and lymphocytes
B. Interstitial congestion and edema, lymphocytic infiltration
C. Interstitial congestion, edema and bleeding
D. Interstitial congestion and edema, neutrophil infiltration
E. Interstitial congestion and edema, infiltration of plasma cells

42. Which of the following types of pneumonia has inclusion bodies in alveolar epithelial cells and/or macrophages?

A. Streptococcus pneumonia
B. Viral pneumonia
C. Mycoplasma pneumonia
D. Bacillary pneumonia
E. Fungal pneumonia

43. Which of the following types of pneumonia is likely to be caused by influenza virus?

A. Lobular pneumonia
B. Lobar pneumonia
C. Interstitial pneumonia
D. Hypersensity pneumonia
E. Hemorrhagic pneumonia

44. Which of the following is a diagnostic indicator of viral pneumonia?

A. Interstitial pneumonia
B. Infiltration of lymphocytes and monocytes
C. Virus inclusion bodies within epithelial cells
D. Hyperplasia of alveolar cell
E. Hyaline membrane formation in alveoli

45. Which of the following lesions is charac-

teristic of viral pneumonia?

A. Lobular lesions
B. Lobar lesions
C. Reticulate or irregular patchy lesions
D. Confluent lobular lesions
E. Segmental lesions

46. Which of the following is not present in lobar pneumonia?

A. Fibrinous inflammation of alveoli
B. Bronchial suppuration
C. Toxic shock
D. Lung abscess
E. Pulmonary carnification

47. Which of the following diseases is not an interstitial type pneumonia?

A. Mycoplasmal pneumonia
B. Chlamydia pneumonia
C. Legionella pneumonia
D. Viral pneumonia
E. SARS

48. Which of the following is not true in legionella pneumonia?

A. It can manifest as lobar pneumonia
B. It can manifest as confluent bronchopneumonia
C. It can manifest as interstitial inflammation
D. It can manifest as lobular pneumonia
E. It can manifest as purulent pleuritis

49. Which of the following lesions is characteristic of lobar pneumonia?

A. Lymphocytic exudation
B. Fibrin exudation
C. Serous exudation
D. Monocyte and macrophage exudation
E. Neutrophil exudation

50. Which of the following lesions is characteristic of lobular pneumonia?

A. Lymphocytic exudation
B. Fibrin exudation
C. Serous exudation
D. Monocyte and macrophage exudation
E. Neutrophil exudation

51. Which of the following lesions is characteristic of lobular pneumonia?

A. Lobular suppurative inflammation surrounding bronchioles
B. Fibrinous inflammation of alveoli
C. Hemorrhagic inflammation of alveoli
D. Interstitial non-suppurative inflammation
E. Serous inflammation of alveoli

52. Which of the following is the typical lesion of lobular pneumonia?

A. Purulent inflammation of the trachea
B. Necrotizing inflammation of alveoli
C. Hyaline membrane formation of the alveolar wall
D. Fibrinous inflammation of alveoli
E. Suppurative inflammation surrounding bronchioles

53. Which of the following is not characteristic of lobular pneumonia?

A. The lesions begin from bronchioles

B. It is caused by a variety of bacteria
C. Suppurative inflammation
D. The patient often coughs up rust-colored sputum
E. The patient often coughs up mucopurulent sputum

54. Which of the following is not a complication of lobular pneumonia?
A. Pulmonary carnification
B. Lung abscess
C. Pyemia
D. Respiratory failure
E. Empyema

55. Which of the following lesions is characteristic of lobular pneumonia?
A. Fibrinous inflammation
B. Serous inflammation
C. Suppurative inflammation
D. Catarrhal inflammation
E. Necrotizing inflammation

56. Which of the following is characteristic of lobular pneumonia?
A. It often develops from acute bronchiolitis
B. Fibrinous inflammation of bronchi and alveoli
C. It can develop into a lobar pneumonia
D. It is often complicated by obstructive pulmonary emphysema
E. It will not destroy the alveolar structure

57. Which of the following is not characteristic of lobar pneumonia?
A. Many heart failure cells in the alveolar spaces
B. Serous exudation in the alveolar spaces
C. Red blood cell in the alveolar spaces alveolar cavity
D. Fibrinous inflammation in the alveolar space exudate
E. Neutrophils infiltration in the alveolar spaces

58. Which of the following is not characteristic of lobar pneumonia?
A. Involves a whole lobe or more than one lobes
B. Usually infected by Streptococcus pneumoniae
C. Fibrinous inflammation
D. The structure of the small bronchial wall and alveolar wall is damaged
E. Pulmonary carnification

59. Which of the following pathogens is found most frequently in pneumonia?
A. Bacteria
B. Chlamydia
C. Mycoplasma
D. Virus
E. Fungi

60. Which of the following is the most common cause of acute infection of the upper respiratory tract?
A. Cold

B. Bacteria
C. Mycoplasma
D. Virus
E. Fungi

61. Which of the following is characteristic of the red liver-like phage of lobar pneumonia?
A. Fibrinous inflammation
B. Hemorrhagic inflammation
C. Hemorrhagic serous inflammation
D. hemorrhagic fibrinous inflammation
E. Fibrinous purulent inflammation

62. Which of the following is characteristic of lobar pneumonia?
A. Hemorrhagic inflammation
B. Suppurative inflammation
C. Fibrinous inflammation
D. Serous inflammation
E. Necrotizing inflammation

63. Which of the following is characteristic of pulmonary carnification?
A. Sarcoma cells diffusely infiltrate the lung
B. Chronic pulmonary congestion and fibrosis
C. Cancer cells infiltrate the lung resulting consolidation
D. Organization of lung exudation
E. Formation of pulmonary inflammatory pseudotumor

64. Which of the following is the pathologic phases is present when a patient with lobar pneumonia coughs up rust colored sputum?
A. Congestion and edema phase
B. Red liver-like phase
C. Resolution phase
D. Gray liver-like phase
E. Recovery stage

65. Which of the following pathologic phases is present when rust colored sputum changes into mucoid purulent sputum in a patient with lobar pneumonia?
A. Gray liver-like phase
B. Red liver-like phase
C. Resolution phase
D. Recovery stage
E. Congestion and edema phase

66. There are macrophages in the sputum whose cytoplasm contains hemosiderin when there is lung congestion. Which of the following is the best name?
A. Large mononuclear cells
B. Dust cells
C. Macrophages in tissue
D. Heart failure cells
E. Foreign-body giant cells

67. A 35-year-old man has had an infection of the upper respiratory tract for two days. His condition rapidly deteriorated. He developed fever, chills, chest pain, cough, and rust-colored sputum. His temperature is 39.8℃. A chest radiograph shows a large dense shadow in the left lower lobe.

Which of the following conditions is not usually associated with this patients illness?

A. Lung abscess
B. Septic shock
C. Pulmonary carnification
D. Bronchiectasis
E. Fibrinous pleuritis

68. Which of the following is the cause of pulmonary carnification in patients with lobar pulmonary?

A. Fibrin exudation
B. Ischemia of local tissue
C. Lower resistance of the body
D. Smaller number and dysfunction of phagocytic cells
E. Highly pathogenic bacteria

69. Which of the following is likely to be present when lobar pneumonia is combined with staphylococcus infection?

A. Inflammatory pseudotumor
B. Lung abscess
C. Obstructive pulmonary emphysema
D. Pulmonary carnification
E. Fibrinous pleuritis

70. Which of the following changes in pulmonary vessels is not associated with chronic pulmonary heart disease?

A. The number of capillaries is reduced
B. Fibrinoid necrosis of medium-sized arteries
C. Muscularization of arterioles
D. Thickening of intima in small muscular arteries
E. Pulmonary arteritis

71. Which of the following pathological changes is not present in chronic pulmonary heart disease?

A. Fibrinoid necrosis of pulmonary muscular arterioles
B. Myocardial dissolution and fibrosis
C. Diffuse obstructive pulmonary emphysema
D. Pulmonary fibrosis
E. Muscularization of arterioles

72. Which of the following factors is not associated with chronic obstructive pulmonary emphysema?

A. Dysfunction of ventilation in small airways
B. Increase in endogenous elastase
C. Smoking
D. Hereditary α_1-antitrypsin deficiency
E. Pulmonary cystic fibrosis

73. Which of the following lesions is not present in chronic bronchitis?

A. The cilia of columnar epithelium show lodging and shedding
B. Hyperplasia of bronchial glands and metaplasia of mucous glands
C. There are a large numbers of lymphocytes in the bronchial wall
D. Atypical hyperplasia of cartilage cells in bronchial wall

E. Squamous metaplasia of bronchial mucosa

74. Which of the following diseases leads most frequently to pulmonary heart disease?
    A. Chronic obstructive pulmonary emphysema
    B. Thoracic malformation
    C. Silicosis
    D. Chronic tuberculosis
    E. Asthma
75. Which of the following is not appropriate in describing pulmonary heart disease?
    A. Persistent pulmonary hypertension is the basis of pathogenesis
    B. It may be caused by chronic obstructive pulmonary emphysema
    C. It can cause systemic congestion
    D. It can cause lung congestion
    E. It can cause right heart failure
76. Which of the following is not associated with chronic pulmonary heart disease?
    A. Chronic obstructive pulmonary disease
    B. Pulmonary hypertension
    C. Atherosclerosis of the pulmonary artery
    D. Myocardial hypertrophy of the right ventricle
    E. Thromboembolism of the main branch of the pulmonary artery
77. Which of the following types of arterial lesions is not associated wtih pulmonary heart disease?
    A. Small vessel arteritis
    B. Hyalinization of small arteries
    C. Fibroelastosis of arterioles
    D. Muscularization of arterioles
    E. Intimal hyperplasia of pulmonary arterioles
78. Which of the following is not an outcome of pulmonary heart disease?
    A. Respiratory failure
    B. Myocardial infarction
    C. Arrhythmia
    D. Pulmonary encephalopathy
    E. Expansion of the right ventricle
79. Which of the following is the common cause of acute pulmonary heart disease?
    A. Gas embolism
    B. Fat embolism
    C. Thromboembolism of the main branch of the pulmonary artery
    D. Thromboembolism of a small branch of the pulmonary artery
    E. Amniotic fluid embolism
80. Which of the following is the key mechanism of pulmonary heart disease?
    A. Infection of the lung
    B. Obstruction of small airways
    C. Pulmonary hypertension
    D. Pulmonary interstitial fibrosis
    E. Pulmonary thrombosis
81. Which is the following is not a cause of

pulmonary heart disease?

A. Pulmonary embolism
B. Interstitial pneumonia
C. Chronic fibro-cavitative pulmonary tuberculosis
D. Silicosis
E. Emphysema

82. Which of the following is the most common cause of pulmonary heart disease?

A. Chronic obstructive pulmonary emphysema
B. Tuberculoma
C. Lobar pneumonia
D. Lung cancer, peripheral type
E. Silicosis

83. Which of the following findings is the main standard for pathologic diagnosis of pulmonary heart disease?

A. The thickness of the right ventricular wall is more than 5 mm in the area two centimeters below the pulmonic valve
B. The apex of heart is blunt and round
C. There is expansion of the right ventricle
D. There is bulging of the pulmonary cone
E. There is endothelial thickening of the pulmonary artery

84. Which of the following chronic obstructive pulmonary diseases is the most common?

A. Asthma
B. Bronchiectasis
C. Silicosis
D. Chronic bronchitis
E. Chronic fibro-cavitative pulmonary tuberculosis

85. Which of the following is involved in lesions of centriacinar emphysema?

A. Terminal bronchiole
B. Alveolar ducts
C. Alveolar sac
D. Respiratory bronchiole
E. Alveoli

86. Which of the following is not true of centriacinar emphysema?

A. The lesion involves the central part of the acinus, and lesions of the respiratory bronchiole are most obvious, showing cystic dilatation
B. Chronic bronchiolitis is the main cause
C. The lesion is associated with hereditary α_1-antitrypsin deficiency
D. The lesion is frequent in smokers
E. Infection of small airways may play a role

87. Which of the following types of emphysema is closely associated with hereditary α_1-antitrypsin deficiency?

A. Centriacinar emphysema
B. Interstitial emphysema
C. Panacinar emphysema
D. Bullous emphysema

E. Paracicatrical emphysema

88. Which of the following chronic lung diseases is associated with hereditary α_1-antitrypsin deficiency?
    A. Goodpasture syndrome
    B. Panacinar emphysema
    C. Bronchiectasis
    D. Chronic bronchitis
    E. Pulmonary fibrosis
89. Which type of emphysema involves the entire acinus?
    A. Centriacinar emphysema
    B. Panacinar emphysema
    C. Interstitial emphysema
    D. Periacinar emphysema
    E. Irregular emphysema
90. Which of the following is not associated with the pathogenesis of emphysema?
    A. Destruction of alveolar wall and reduced elasticity
    B. Airway narrowing and obstruction
    C. Squamous metaplasia
    D. Increase in elastase
    E. Smoking
91. Which is the main factor leading to emphysema?
    A. Air pollution
    B. Smoking
    C. Infection of small airways
    D. Chronic obstructive bronchiolitis
    E. Cold
92. Which of the following is not included in emphysema lesions?
    A. Terminal bronchioles
    B. Alveoli
    C. Alveolar tube
    D. Respiratory bronchiole
    E. Alveolar sacs
93. A 70-year-old woman has a nearly 10 year history of chronic bronchitis She has had shortness of breath, chest distress and dyspnea for nearly 3 years. She has barrel chest which is hyperresonant on percussion, with weakened respiratory movement, and weakened breath sounds. Which of the following pathologic conditions is NOT typically associated with these clinical findings?
    A. Narrowing of alveolar septa
    B. Alveolar expansion
    C. Alveoli fusing into vesicles
    D. Extensive pulmonary fibrosis
    E. Fracture of the alveolar septum
94. There is formation of small beaded bubbles at the junction of the lobular interval and pleura. Which of the following is the most likely diagnosis?
    A. Emphysema, irregular type
    B. Periacinar emphysema
    C. Bullae of the lung
    D. Interstitial emphysema
    E. Centriacinar emphysema
95. Which of these is the most common complication of chronic bronchitis?

A. Emphysema
B. lobular pneumonia
C. Lung Abscess
D. Bronchiectasis
E. Asthma

96. Which type of epithelial metaplasia is prone to occur in patients with chronic bronchitis?
A. Transitional cell metaplasia
B. Mucinous metaplasia
C. Squamous metaplasia
D. Goblet metaplasia
E. Columnar cell metaplasia

97. Which of these is the pathological basis of ventilation dysfunction in patients with chronic bronchitis?
A. Lodging and shedding of epithelial cilia
B. Hypertrophy and hyperplasia of mucous glands
C. Degeneration and atrophy of cartilage
D. Obstruction of small airways
E. Squamous metaplasia of bronchial mucosa

98. A 60-year-old woman develops a chronic cough with white foamy mucous sputum in autumn and winter each year, continuing each time for about three months. Which of the following conditions is NOT true of the disease associated with these clinical findings?
A. It can cause lobar pneumonia
B. Smoking is a common cause
C. It can cause pulmonary heart disease
D. It may be associated with asthma
E. It can cause emphysema

99. Which factor is not closely related to the occurrence of chronic bronchitis?
A. Viral infection
B. Bacterial infection
C. Mycoplasma infection
D. Harmful gases
E. Smoking

100. A 60-year-old woman develops a chronic cough with white foamy mucous sputum in autumn and winter each year, which continues for about three months each time. Which of the following pathologic conditions is not associated with these clinical findings?
A. Fibroelastosis of the bronchial wall
B. Hyperplasia of goblet cells in mucosal epithelium
C. Hypertrophy of mucous glands
D. Degeneration and atrophy of bronchial cartilage
E. Squamous metaplasia of bronchial mucosa

101. Which of these lesions is the basis for sputum production in patients with chronic bronchitis?
A. Degeneration, necrosis and detachment of bronchial epithelial cells
B. Hypertrophy of glands, hyperplasia of

goblet cells in mucosal epithelium

C. Atrophy, fibrosis, calcification and ossification of cartilage

D. Lymphocytic infiltration in the bronchial wall

E. Squamous metaplasia of the bronchial mucosa

答案

中文题

1.C	2.E	3.B	4.A	5.B	6.D	7.D	8.E	9.C	10.B
11.D	12.C	13.D	14.D	15.A	16.E	17.E	18.D	19.B	20.D
21.E	22.E	23.D	24.D	25.E	26.E	27.B	28.B	29.C	30.E
31.D	32.C	33.A	34.D	35.A	36.E	37.A	38.D	39.D	40.B
41.D	42.B	43.A	44.D	45.A	46.B	47.B	48.D	49.C	50.D
51.D	52.E	53.E	54.E	55.A	56.D	57.D	58.D	59.E	60.C
61.E	62.C	63.B	64.C	65.E	66.D	67.C	68.E	69.C	70.E
71.D	72.C	73.E	74.C	75.E	76.A	77.E	78.C	79.D	80.A
81.E	82.C	83.A	84.A	85.D	86.D	87.D	88.A	89.E	90.B
91.D	92.A	93.A	94.D	95.E	96.A	97.E	98.D	99.E	100.E
101.E	102.D	103.C	104.E	105.E	106.D	107.A	108.E	109.E	110.C
111.E	112.A	113.D	114.C	115.C	116.A	117.D	118.B	119.E	120.D
121.D	122.A	123.D	124.E	125.D	126.E	127.A	128.C	129.B	130.C
131.C	132.C	133.E	134.A	135.E	136.A	137.E	138.E	139.D	140.C
141.A	142.E	143.C	144.B	145.C	146.E	147.A	148.D	149.A	150.D
151.E	152.B	153.A	154.B	155.B	156.E	157.E	158.B	159.D	160.D

英文题

1.C	2.A	3.E	4.A	5.D	6.D	7.D	8.B	9.B	10.B
11.A	12.C	13.D	14.C	15.A	16.A	17.A	18.D	19.B	20.A
21.D	22.A	23.C	24.D	25.C	26.D	27.C	28.D	29.A	30.C
31.B	32.D	33.A	34.A	35.B	36.C	37.C	38.A	39.D	40.E
41.A	42.B	43.C	44.C	45.C	46.B	47.C	48.C	49.B	50.E
51.A	52.E	53.D	54.A	55.C	56.A	57.A	58.D	59.A	60.D
61.D	62.C	63.D	64.B	65.A	66.D	67.D	68.D	69.B	70.B
71.A	72.E	73.D	74.A	75.D	76.E	77.B	78.B	79.C	80.C
81.B	82.A	83.A	84.D	85.D	86.C	87.C	88.B	89.B	90.C
91.D	92.A	93.D	94.D	95.A	96.C	97.D	98.A	99.C	100.A
101.B									

7　消化系统疾病

中文题

1. 慢性浅表性胃炎的主要病变特点是
   A. 胃粘膜变薄
   B. 胃粘膜水肿
   C. 可见粘膜下血管
   D. 淋巴细胞、浆细胞浸润胃粘膜浅层
   E. 胃粘膜糜烂
2. 符合慢性萎缩性胃炎 B 型表现为
   A. 胃体部弥漫性病变
   B. 不发生癌变
   C. 血清自身抗体阳性
   D. 常伴有恶性贫血
   E. 胃粘膜变薄
3. 慢性萎缩性胃炎的特征病变是
   A. 粘膜腺体萎缩
   B. 粘膜炎症
   C. 血清自身抗体阳性
   D. 肠上皮化生
   E. 假幽门腺化生
4. 慢性胃炎的主要病因是
   A. 吸烟
   B. 酗酒
   C. 幽门螺杆菌感染
   D. 免疫因素
   E. 胃大部切除
5. 不符合慢性萎缩性胃炎的描述是
   A. 可发生肠上皮化生
   B. 可引起胃酸增高
   C. 可合并胃溃疡
   D. 可引起胃泌素增高
   E. 可引起恶性贫血
6. 慢性胃炎与胃癌发生关系密切的病变是
   A. 肠上皮化生
   B. 上皮异型增生
   C. 淋巴滤泡形成
   D. 粘膜上皮再生
   E. 假幽门腺化生
7. 不符合 A 型慢性萎缩性胃炎的描述是
   A. 可发生肠上皮化生
   B. 可引起胃酸增高
   C. 可引起壁细胞减少
   D. 可引起 G 细胞增多
   E. 可引起恶性贫血
8. 不符合 B 型慢性萎缩性胃炎的描述是
   A. 可发生肠上皮化生
   B. 可引起胃酸减少
   C. 可引起假幽门腺化生
   D. 常由幽门螺杆菌引起
   E. 可引起恶性贫血
9. 慢性胃溃疡病变位置最常见于
   A. 胃前壁
   B. 胃后壁
   C. 胃大弯及胃底部
   D. 胃大弯近幽门处
   E. 胃小弯近幽门窦部
10. 慢性消化性溃疡最常见的好发部位是
   A. 胃小弯近幽门部
   B. 十二指肠球部

C. 胃与十二指肠复合
D. 胃体部
E. 十二指肠下段

11. 男性，40 岁，10 多年来，反酸、嗳气、上腹部饱胀，伴有规律性上腹痛，空腹发作，夜间更重，进食可缓解。胃镜检查，十二指肠球部可见直径约 1cm 的圆形溃疡。不符合该患者疾病的病因发病的描述是
A. 胃酸分泌过多
B. 幽门螺杆菌感染
C. 粘膜屏障受破坏
D. 胆汁反流
E. 女性易发生

12. 幽门螺杆菌感染引起慢性胃溃疡的机制是
A. 引起胆汁反流
B. 引起胃酸分泌增多
C. 引起胃排空延缓
D. 降低胃粘膜的防御能力
E. 引起免疫功能紊乱

13. 不符合慢性十二指肠溃疡的描述是
A. 多位于十二指肠球部
B. 直径多在 1cm 以内
C. 易发生癌变
D. 易合并穿孔
E. 易发生出血

14. 与慢性消化性溃疡的发生无关的因素是
A. 胃酸分泌增多
B. 胃泌素瘤
C. 胃窦部 G 细胞增多
D. 迷走神经兴奋性降低
E. 胆汁反流

15. 符合十二指肠溃疡的描述是
A. 多位于十二指肠降部
B. 与胃酸分泌过高有关
C. 不易发生癌变
D. 不易发生出血
E. 不易发生穿孔

16. 与消化性溃疡发生无关的因素是
A. 幽门螺杆菌感染
B. 胆汁反流入胃
C. 胃窦部 C 细胞减少
D. 正常粘膜屏障被破坏
E. 胃酸分泌过多

17. 与消化性溃疡的发生无关的因素是
A. 胃酸分泌增多
B. 胃粘膜萎缩
C. 胰岛细胞瘤
D. 迷走神经过度兴奋
E. 遗传因素

18. 关于胃溃疡的描述，错误的是
A. 可发生癌变
B. 直径多在 2.5cm 以内
C. 呈圆形或椭圆形
D. 多局限于粘膜下层
E. 多位于小弯侧

19. 慢性消化性溃疡不具备的病理变化是
A. 炎性渗出物
B. 肉芽组织
C. 瘢痕组织
D. 增殖性动脉内膜炎
E. 神经纤维肿瘤性增生

20. 不符合慢性胃溃疡合并症的是
    A. 穿孔
    B. 幽门狭窄
    C. 癌变
    D. 出血
    E. 粘膜萎缩
21. 女性，48 岁，10 多年来，反酸、嗳气、上腹部饱胀，伴有规律性上腹痛，空腹发作，夜间更重，进食可缓解。符合该患者疾病的诊断是
    A. 慢性胃炎
    B. 食管炎
    C. 十二指肠溃疡
    D. 胃癌
    E. 慢性胆囊炎
22. 男性，38 岁，10 多年来，反酸、嗳气、上腹部饱胀，伴有规律性上腹痛，空腹发作，夜间更重，进食可缓解。胃镜检查，十二指肠球部可见直径约 1cm 的圆形溃疡。不符合该患者疾病的描述是
    A. 易发生穿孔
    B. 易发生出血
    C. 易引起幽门梗阻
    D. 易发生癌变
    E. 可经不完全再生修复
23. 慢性胃溃疡最常见的合并症是
    A. 幽门狭窄
    B. 穿孔
    C. 出血
    D. 癌变
    E. 粘连
24. 女性，30 岁，10 年来，反酸、嗳气、上腹部饱胀，伴有规律性上腹痛，进食后加重。胃镜检查，胃窦小弯侧可见溃疡。不支持消化溃疡诊断的是
    A. 溃疡直径小于 2cm
    B. 溃疡边缘整齐
    C. 溃疡底部平坦
    D. 溃疡周围粘膜皱襞中断、结节状肥厚
    E. 溃疡呈椭圆形
25. 与阑尾炎发病无关的因素是
    A. 细菌感染
    B. 阑尾粘膜损伤
    C. 阑尾腔阻塞
    D. 纤维多的饮食
    E. 阑尾平滑肌痉挛
26. 最易引起阑尾穿孔的阑尾炎是
    A. 急性蜂窝织炎性阑尾炎
    B. 急性单纯性阑尾炎
    C. 急性坏疽性阑尾炎
    D. 慢性阑尾炎
    E. 慢性阑尾炎急性发作
27. 阑尾系膜化脓性血栓性静脉炎易引起
    A. 肝脓肿
    B. 肺脓肿
    C. 脑脓肿
    D. 心室壁脓肿
    E. 肾盂肾炎
28. 急性阑尾炎最常见的并发症是
    A. 肝脓肿
    B. 化脓性腹膜炎
    C. 肠瘘
    D. 肠梗阻

E. 阑尾粘液囊肿

29. 急性单纯性阑尾炎的病变特点是

A. 阑尾全层弥漫性中性粒细胞浸润

B. 阑尾壁局部坏疽

C. 阑尾粘膜坏死伴中性粒细胞浸润

D. 阑尾壁纤维组织增生伴慢性炎细胞浸润

E. 阑尾腔闭塞，远端粘液潴留

30. 急性蜂窝织炎性阑尾炎的病变特点是

A. 阑尾全层弥漫性中性粒细胞浸润

B. 阑尾壁局部坏疽

C. 阑尾粘膜坏死伴中性粒细胞浸润

D. 阑尾壁纤维组织增生伴慢性炎细胞浸润

E. 阑尾腔闭塞，远端粘液潴留

31. 慢性阑尾炎的病变特点是

A. 阑尾全层弥漫性中性粒细胞浸润

B. 阑尾壁局部坏疽

C. 阑尾粘膜坏死伴中性粒细胞浸润

D. 阑尾壁纤维组织增生伴慢性炎细胞浸润

E. 阑尾腔闭塞，远端粘液潴留

32. 女性，18 岁，转移性右下腹痛 1 天，伴恶心、呕吐。急诊入院，查体：体温 38.5℃，血液检查：中性粒细胞 18×10^9/L，右下腹压痛反跳痛明显。患者的疾病诊断最可能是

A. Meckel 憩室炎

B. 宫外孕

C. 急性阑尾炎

D. 肠系膜淋巴结炎

E. 卵巢滤泡破裂

33. 缺陷性肝炎病毒是

A. HAV

B. HBV

C. HCV

D. HDV

E. HEV

34. 我国病毒性肝炎最常见的病原体是

A. HAV

B. HBV

C. HCV

D. HDV

E. HEV

35. 经血传播的 RNA 肝炎病毒是

A. HAV

B. HBV

C. HCV

D. HDV

E. HEV

36. 丙型肝炎病毒（HCV）的主要传播途径是

A. 经血传播

B. 经消化道传播

C. 经呼吸道传播

D. 性传播

E. 母婴传播

37. 乙型肝炎病毒（HBV）的主要传播途径是

A. 经血传播

B. 经消化道传播

C. 经呼吸道传播

D. 性传播

E. 母婴传播

38. 甲型肝炎病毒（HAV）的主要传播途径是
A. 经血传播
B. 经消化道传播
C. 经呼吸道传播
D. 性传播
E. 母婴传播

39. 丁型肝炎病毒（HDV）主要感染
A. HAV 携带者
B. HBV 携带者
C. HCV 携带者
D. HEV 携带者
E. HGV 携带者

40. 可引起慢性肝炎的 RNA 病毒是
A. HAV
B. HBV
C. HCV
D. HDV
E. HEV

41. 病毒性肝炎的基本病变不包括
A. 肝细胞水肿
B. 肝细胞嗜酸性变
C. 肝细胞坏死
D. 肝细胞异型增生
E. 小胆管增生

42. 肝细胞碎片状坏死多见于
A. 急性普通型肝炎
B. 急性重型肝炎
C. 慢性持续性肝炎
D. 慢性活动性肝炎
E. 亚急性重型肝炎

43. 急性普通型肝炎的病变特点是
A. 肝细胞广泛变性，点灶状坏死
B. 肝细胞广泛变性，碎片状坏死
C. 肝细胞广泛变性，桥接坏死
D. 肝细胞大片坏死
E. 肝细胞广泛变性，亚大片坏死

44. 不符合病毒性肝炎基本病变的是
A. 肝细胞变性坏死
B. 小胆管增生
C. 纤维组织增生
D. 慢性炎细胞浸润
E. Kupffer 细胞异型增生

45. 不符合慢性活动性肝炎的描述是
A. 肝细胞大片坏死
B. 破坏肝小叶结构
C. 炎症反应明显
D. 容易转为肝硬化
E. 肝细胞弥漫变性

46. 毛玻璃样肝细胞出现于
A. 急性普通型肝炎
B. 急性重型肝炎
C. 慢性持续性肝炎
D. 慢性活动性肝炎
E. HBsAg 携带者

47. 病毒性肝炎时，肝细胞凋亡的病变是
A. 肝细胞灶状坏死
B. 肝细胞桥接坏死
C. 肝细胞大片坏死
D. 肝细胞嗜酸性坏死
E. 肝细胞碎片状坏死

48. 不符合肝细胞气球样变的描述是
A. 细胞内水分增多
B. 细胞内糖原增多

C. 内质网扩张，囊泡变
D. 线粒体肿胀嵴消失
E. 粗面内质网核蛋白体颗粒消失

49. 肝细胞碎片状坏死或桥接坏死常见于
A. 急性重型肝炎
B. 亚急性重型肝炎
C. 慢性持续性肝炎
D. 慢性活动性肝炎
E. 急性普通型肝炎

50. 健康携带者多见于
A. HBV 感染
B. HCV 感染
C. HAV 感染
D. HEV 感染
E. HDV 感染

51. 肝细胞亚大片坏死，伴肝细胞结节状再生常见于
A. 慢性持续性肝炎
B. 慢性活动性肝炎
C. 急性普通型肝炎
D. 急性重型肝炎
E. 亚急性重型肝炎

52. 急性重型肝炎最常发生于
A. HBV 感染
B. HCV 感染
C. HAV 感染
D. HEV 感染
E. 药物中毒

53. 病毒性肝炎患者血清 ALT 水平升高的原因是
A. 淋巴细胞浸润
B. 胆管上皮增生
C. 肝细胞增生
D. 肝细胞变性坏死
E. Kupffer 细胞增生

54. 女性，25 岁，近 2 周来，食欲不振，恶心、乏力，皮肤、巩膜黄染。肝区疼痛，肝功检查：ALT 100U，血总胆红素 18mg%，HBsAg（+）。肝穿刺检查，组织学改变有：弥漫性肝细胞水肿，Kupffer 细胞增生，局灶性肝细胞脂肪变性，肝小叶内淋巴细胞浸润，肝细胞点灶状坏死，病变符合
A. 急性重症肝炎
B. 亚急性重症肝炎
C. 急性普通型肝炎
D. 慢性活动性肝炎
E. 慢性持续性肝炎

55. 男性，27 岁，5 年前，食欲不振，恶心、乏力，皮肤、巩膜黄染。肝区疼痛，肝功检查：ALT 100U，血总胆红素 18mg%，HBsAg（+）。肝穿刺诊断为急性普通型肝炎，经治疗好转。近半年来，又出现乏力、肝区隐痛，ALT 60U～80U。符合该患者疾病的描述是
A. 肝细胞大片坏死
B. 肝细胞亚大片坏死
C. 肝细胞碎片坏死
D. 肝细胞点状坏死
E. 肝细胞灶状坏死

56. 女性，18 岁，1 周来高热、恶心、呕吐，嗜睡，皮肤、巩膜黄染，肝区疼痛。ALT 150U，胆红素 35mg%，HBsAg（+）。住院 2 天，出现肝臭，昏迷。因

胃肠道广泛出血而死亡。符合患者疾病的诊断是

A. 急性重症肝炎
B. 亚急性重症肝炎
C. 急性普通型肝炎
D. 慢性活动性肝炎
E. 慢性持续性肝炎

57. 肝硬化假小叶形成的最主要因素是

A. 肝细胞变性坏死
B. 肝细胞再生
C. 纤维结缔组织增生
D. 小胆管增生
E. 单核淋巴细胞浸润

58. 门脉性肝硬化的特点是

A. 结节大小不等
B. 结节大小相仿，纤维间隔薄而均匀
C. 肝脏呈深绿色
D. 纤维组织沿门静脉分支大量增生
E. 纤维间隔厚薄不均

59. 坏死后肝硬化的特点是

A. 结节大小不等，纤维间隔厚薄不均
B. 结节大小相仿
C. 肝脏呈深绿色
D. 纤维组织沿门静脉分支大量增生
E. 纤维间隔薄而均匀

60. 血吸虫性肝硬化病变特点是

A. 结节大小较一致，纤维间隔薄而均匀
B. 肝脏变小、变硬、深绿色
C. 结节大小不一，纤维间隔厚薄不均
D. 纤维结缔组织沿门静脉分支增生，将肝脏分隔为结节状
E. 肝脏淤血，纤维组织增生，分隔肝小叶

61. 胆汁性肝硬化病变特点是

A. 结节大小较一致，纤维间隔薄而均匀
B. 肝脏变小、变硬、深绿色
C. 结节大小不一，纤维间隔厚薄不均
D. 纤维结缔组织沿门静脉分支增生，将肝脏分隔为结节状
E. 肝脏淤血，纤维组织增生，分隔肝小叶

62. 假小叶的基本病变不包括

A. 肝细胞核增大、深染
B. 肝细胞团内中央静脉偏位或多个
C. 肝细胞团内出现汇管区
D. 肝细胞团被增生的纤维组织分隔包绕
E. 肝细胞索排列紊乱

63. 我国肝硬化最常见的原因是

A. 慢性酒精中毒
B. 营养缺乏
C. 毒物中毒
D. 病毒性肝炎
E. 药物中毒

64. 不符合假小叶病变特点的描述是

A. 肝细胞索围绕中央静脉呈放射状排列
B. 缺少中央静脉
C. 中央静脉偏位或有 2 个以上
D. 肝细胞排列紊乱
E. 出现汇管区

65. 肝硬化时，门脉高压症的主要原因是

A. 肝细胞索排列紊乱
B. 小叶下静脉、中央静脉及肝血窦受压
C. 肝动脉与肝静脉之间形成异常吻合支
D. 小胆管增生
E. 慢性炎细胞浸润

66. 肝硬化时，肝功能不全的表现有
A. 出血倾向
B. 食道下段静脉曲张及破裂出血
C. 胃肠道淤血
D. 脾功能亢进
E. 腹水

67. 不符合肝硬化腹水形成原因的描述是
A. 小叶下静脉受压
B. 窦内压升高
C. 肝细胞合成蛋白功能降低
D. 血中醛固酮、抗利尿素水平升高
E. 肝动脉与肝静脉异常吻合形成

68. 不符合门脉高压的临床表现是
A. 脾肿大
B. 腹水
C. 食管静脉曲张
D. 蜘蛛痣
E. 胃肠道淤血

69. 不符合肝功能不全的临床表现是
A. 皮肤粘膜出血
B. 男性乳房发育
C. 食管静脉曲张
D. 蜘蛛痣
E. 肝性脑病

70. 不符合肝硬化腹水的描述是
A. 混浊，可自行凝固
B. 与门脉高压有关
C. 与白蛋白减少有关
D. 与醛固酮升高有关
E. 与抗利尿激素升高有关

71. 男性，59 岁。慢性肝炎病史 20 年。近 3 年来双下肢浮肿，因上消化道出血急诊入院。查体：巩膜黄染，皮肤蜘蛛痣，腹水（+），脾肋缘下 3cm。符合该患者疾病的描述是
A. 酒精性肝硬化
B. 慢性肝淤血
C. 胆汁性肝硬化
D. 血吸虫性肝硬化
E. 门脉性肝硬化

72. 男性，59 岁。慢性肝炎病史 20 年。近 3 年来双下肢浮肿，因上消化道出血急诊入院。查体：巩膜黄染，皮肤蜘蛛痣，腹水（+），脾肋缘下 3cm。对该患者疾病最有诊断意义的是
A. 假小叶形成
B. 肝细胞增生
C. 小胆管增生
D. 肝细胞脂肪变性
E. 肝细胞水肿

73. 最常见的食管良性肿瘤是
A. 纤维瘤
B. 脂肪瘤
C. 血管瘤
D. 平滑肌瘤
E. 神经纤维瘤

74. 最常见的食管恶性肿瘤是

A. 腺癌
B. 鳞癌
C. 未分化癌
D. 类癌
E. 恶性黑色素瘤

75. 不符合早期食管癌的描述是
A. 癌的直径小于 2.0cm
B. 浸润深度不超过粘膜下层
C. 钡餐检查可有轻度局限性僵硬
D. 无明显临床症状
E. 无淋巴结转移

76. 早期食管癌是指
A. 粘膜内癌
B. 原位癌
C. 粘膜下癌
D. 无淋巴结转移
E. 以上都是

77. 中晚期食管癌最常见的肉眼形态是
A. 形成明显的环形狭窄
B. 形成较深的溃疡缺损
C. 如蘑菇状突入管腔
D. 灰白色、质地较软
E. 灰红色、质地较软

78. 食管癌的最常见的好发部位是
A. 颈段
B. 上段
C. 中段
D. 下段
E. 贲门

79. 食管的癌前病变是
A. 食管炎
B. 食管憩室
C. 食管狭窄
D. Barrett 食管
E. 贲门失驰

80. 不符合 Barrett 食管的描述是
A. 与食管反流有关
B. 内镜下食管下段可见色泽红润的斑块
C. 镜下腺上皮替代鳞状上皮
D. 可发生腺上皮异型增生
E. 与鳞癌发生关系密切

81. 早期胃癌最多见的类型是
A. 隆起型
B. 表浅型
C. 表浅凹陷型
D. 表浅平坦型
E. 凹陷型

82. 早期胃癌是指
A. 粘膜内癌
B. 未侵及肌层的癌
C. 粘膜下癌
D. 无淋巴结转移的癌
E. 侵及肌层的癌

83. 恶性度最低的胃癌组织学类型是
A. 管状腺癌
B. 印戒细胞癌
C. 单纯癌
D. 硬癌
E. 未分化癌

84. 胃癌最常见的好发部位是
A. 贲门
B. 大弯
C. 胃体

D. 胃窦小弯侧
E. 幽门

85. 革囊胃是指
A. 息肉型胃癌
B. 浸润溃疡型胃癌
C. 胶样型胃癌
D. 弥漫浸润型胃癌
E. 局限溃疡型胃癌

86. 分化较好的胃癌是
A. 腺癌
B. 硬癌
C. 髓样癌
D. 胶样癌
E. 印戒细胞癌

87. 早期胃癌是指
A. 原位癌
B. 粘膜内癌
C. 粘膜下癌
D. 无淋巴结转移的癌
E. 粘膜内癌及粘膜下癌

88. 不符合溃疡型胃癌的描述是
A. 呈火山口状
B. 多大于 2cm
C. 粘膜皱襞向溃疡集中
D. 边缘隆起不整齐
E. 底部凹凸不平

89. 不符合胃癌转移的描述是
A. 可转移至左锁骨上淋巴结
B. 可转移至肝门淋巴结
C. 可血行转移至肝脏
D. 血行转移引起克鲁根博（Krukenberg）瘤
E. 可种植转移至大网膜

90. 胃癌来源于
A. 腺颈部的干细胞
B. 表面粘液细胞
C. 壁细胞
D. 主细胞
E. 吸收细胞

91. 与胃癌预后的决定因素是
A. 胃癌的组织学类型
B. 胃癌的肉眼类型
C. 胃癌的分期
D. 患者的年龄
E. 胃癌的来源

92. 与胃癌发生关系最密切的病变是
A. 慢性表浅性胃炎
B. 慢性萎缩性胃炎
C. 肠上皮化生
D. 假幽门腺化生
E. 胃粘膜异型增生

93. 与胃癌关系密切的息肉是
A. 腺瘤性息肉
B. 增生性息肉
C. 炎症性纤维性息肉
D. Peutz-Jeghers 息肉
E. 幼年性息肉

94. 胃最常见的恶性肿瘤是
A. 平滑肌肉瘤
B. 类癌
C. 腺癌
D. 未分化癌
E. 恶性淋巴瘤

95. 胃癌最主要的转移途径是

A. 淋巴道转移
B. 消化道内转移
C. 血道转移
D. 胃内播散
E. 种植转移

96. 女性，50 岁，慢性萎缩性胃炎 20 余年。胃镜检查，胃窦部大弯侧可见一直径约 1cm 的息肉状隆起。活体检查，病理诊断为中分化腺癌。手术切除癌组织局限于粘膜内。符合该患者早期胃癌的肉眼类型是
A. I 型
B. IIa 型
C. IIb 型
D. IIc 型
E. III 型

97. 大肠最常见的恶性肿瘤是
A. 鳞癌
B. 腺癌
C. 平滑肌肉瘤
D. 黑色素瘤
E. 恶性淋巴瘤

98. 大肠癌最常见的好发部位是
A. 直肠和乙状结肠
B. 盲肠
C. 升结肠
D. 横结肠
E. 降结肠

99. 早期大肠癌最常见的肉眼类型是
A. I 型
B. IIa 型
C. IIb 型
D. IIc 型
E. III 型

100. 早期大肠癌是指
A. 粘膜内癌
B. 未侵及肌层的癌
C. 粘膜下癌
D. 无淋巴结转移的癌
E. 侵及肌层的癌

101. 大肠癌最常见的组织学类型是
A. 中分化腺癌
B. 低分化腺癌
C. 粘液腺癌
D. 印戒细胞癌
E. 未分化癌

102. 右侧大肠癌的常见肉眼类型是
A. 隆起型
B. 溃疡型
C. 浸润型
D. 胶样型
E. 早期癌

103. 左侧大肠癌的常见肉眼类型是
A. 隆起型
B. 溃疡型
C. 浸润型
D. 胶样型
E. 早期癌

104. 大肠鳞癌的好发部位是
A. 肛门直肠
B. 乙状结肠
C. 升结肠
D. 横结肠
E. 降结肠

105. 与大肠癌发生关系不密切的疾病是
A. 腺瘤
B. 家族性腺瘤性息肉病
C. 遗传性非息肉病性大肠癌综合征
D. 慢性溃疡性结肠炎
E. 增生性息肉

106. 与大肠癌发生关系密切的息肉是
A. 腺瘤性息肉
B. 增生性息肉
C. 炎症性息肉
D. Peutz-Jeghers 息肉
E. 幼年性息肉

107. 不符合大肠癌发生发展的描述是
A. 大多数经腺瘤癌变而来
B. 癌变是一个多步骤的演化过程
C. 累及多个基因
D. p53 基因的改变是早期事件
E. 遗传因素起重要作用

108. 不符合消化道类癌的描述是
A. 起源于神经内分泌系统
B. 属于低度恶性肿瘤
C. 阑尾和直肠类癌较少转移
D. 胃和回肠较多转移
E. 转移与肿瘤体积无关

109. 不符合胃肠道淋巴瘤的描述是
A. 与幽门螺杆菌感染有关
B. 多为 T 细胞源性
C. 可引起粘膜增厚，溃疡形成
D. 早期治疗预后良好
E. 肿瘤体积、浸润深度及组织学类型决定预后

110. 不符合大肠癌转移的描述是
A. 可直接蔓延至邻近器官
B. 可经淋巴道转移到结肠上、旁、中间和末端淋巴结
C. 晚期可经血道转移至肝、肺等器官
D. Dukes C 期是指伴淋巴结转移的大肠癌
E. 粘膜内癌常伴淋巴结转移

111. 男性，60 岁，半年来大便带血。结肠镜检查，距肛门 10cm 处可见一溃疡形肿物，约 3cm×5cm 大小，呈火山口状，底部凹凸不平。符合该肿物的病理学改变是
A. 肠粘膜腺体正常
B. 肠粘膜慢性炎
C. 腺体密集、腺上皮细胞异型性明显
D. 肉芽肿形成
E. 乳头形成、乳头被覆上皮轻度异型性

112. 男性，62 岁，直肠癌切除标本，病理检查为低分化腺癌，浸及肠壁深肌层，肠系膜淋巴结无转移。其分期为
A. Dukes A_1 期
B. Dukes A_2 期
C. Dukes B 期
D. Dukes C 期
E. Dukes D 期

113. 肝细胞癌的组织学特点是
A. 形成腺腔
B. 肿瘤细胞弥漫散在
C. 与肝细胞相似的癌细胞形成团块，其间为血窦
D. 形成巨细胞

E. 分泌胆汁

114. 肝脏最常见的恶性肿瘤是

A. 胆管上皮癌

B. 肝细胞癌

C. 血管肉瘤

D. 肝母细胞瘤

E. 恶性淋巴瘤

115. 肝细胞癌的肿瘤标记物是

A. CEA

B. AFP

C. PSA

D. CA19-9

E. 酸性磷酸酶

116. 早期肝癌（小肝癌）的诊断标准是

A. 直径在 1cm 以下，单个瘤结节

B. 直径在 5cm 以下，不超过 2 个瘤结节

C. 直径在 3cm 以下，单个瘤结节

D. 直径在 3cm 以下，不超过 2 个瘤结节

E. 直径在 5cm 以下，单个瘤结节

117. 与肝细胞癌的发生无关的是

A. HBV 感染

B. HCV 感染

C. 中华睾吸虫感染

D. 黄曲霉毒素

E. 亚硝胺类化合物

118. 不符合肝癌的描述是

A. 常合并肝硬化

B. 易发生肝内播散

C. 可引起腹腔大出血

D. 中晚期肝癌预后差

E. 不发生淋巴道转移

119. 我国肝癌最常见的病因是

A. 乙型肝炎

B. 丙型肝炎

C. 黄曲霉毒素

D. 亚硝胺类化合物

E. 血吸虫病

120. 男性，50 岁，慢性肝炎 30 年。2 月来恶心、食欲不振、乏力伴肝区疼痛。超声波显示肝右叶有一直径 5cm 的实性肿物。肝穿刺活检支持肝细胞癌的病理变化是

A. 假小叶形成

B. 肝细胞结节状增生

C. 肝细胞嗜酸性变

D. 肝细胞异型性明显、形成实性团

E. 肝细胞出现双核

121. 不符合急性出血性坏死性肠炎的描述是

A. 病变呈节段性分布

B. 主要累及大肠

C. 小儿多见

D. 引起肠穿孔

E. 肠壁出血坏死

122. 符合急性出血性坏死性肠炎的描述是

A. 主要累及小肠

B. 病变只累及粘膜层

C. 病变界限不清楚

D. 成人多见

E. 由大肠杆菌引起

123. 不符合 Crohn 病的描述是

A. 只累及回肠

B. 呈节段性分布
C. 常呈慢性经过
D. 肠粘膜呈鹅卵石状外观
E. 形成裂隙及匐行性溃疡

124. 符合溃疡性结肠炎的描述是
A. 肉芽肿形成
B. 病变多局限于粘膜和粘膜下层
C. 形成大量腺瘤性息肉
D. 由金黄色葡萄球菌感染引起
E. 溃疡圆形或椭圆形，其长轴与肠管相平行

125. 不符合溃疡性结肠炎的描述是
A. 肉芽肿形成
B. 病变多局限于粘膜和粘膜下层
C. 粘膜多量慢性炎细胞浸润
D. 多量假息肉形成
E. 隐窝脓肿形成

126. 符合 Crohn 病的描述是
A. 肠壁粘膜高度水肿呈鹅卵石状，有匐行溃疡
B. 肠粘膜出现大片坏死并形成大的溃疡
C. 溃疡呈环状与肠轴垂直，边缘不整齐
D. 溃疡边缘整齐，长轴与肠轴平行
E. 溃疡与正常粘膜相间存在

127. 与原发性吸收不良综合征密切相关的因素是
A. 脂肪
B. 动物蛋白
C. 麦胶蛋白
D. 植物纤维
E. 牛奶

128. 不符合菌群失调性假膜性肠炎的描述是
A. 与梭形芽孢杆菌有关
B. 与长期使用广谱抗生素有关
C. 与肠道正常菌群失调有关
D. 形成假膜性肠炎
E. 主要累及小肠

129. 不符合细菌性肝脓肿的描述是
A. 可为单发或多发
B. 经胆道上行感染常呈多发性
C. 经血行感染常呈单发性
D. 可引起脓胸
E. 可引起膈肌下脓肿

130. 不符合细菌性肝脓肿的描述是
A. 多经邻近器官感染直接蔓延而来
B. 死亡率高
C. 多发生于免疫功能低下的患者
D. 可经胆道上行感染
E. 可经血行感染

131. 不符合自身免疫性肝炎的描述是
A. 可发生于类风湿性关节炎患者
B. 可发生于红斑狼疮患者
C. 可发生于溃疡性结肠炎患者
D. 可发生于干燥综合征患者
E. 肝脏基本病变与病毒性肝炎不同

132. 支持自身免疫性肝炎的描述是
A. 中年男性患者多见
B. HBsAg 阳性
C. 自身抗体阳性
D. 肝细胞弥漫水肿
E. 肝细胞灶状坏死

133. 不符合药物性肝病的描述是
A. 可引起肝细胞坏死
B. 可引起淤胆
C. 可引起肝功能异常
D. 可引起肝细胞脂肪变性
E. 不引起肝肿瘤

134. 区分药物性肝病与病毒性肝炎的重要指标是
A. 肝细胞坏死
B. 淤胆
C. 肝功能异常
D. 肝细胞脂肪变性
E. 血清学病毒标志阳性

135. 不符合酒精性肝病的描述是
A. 肝细胞脂肪变性
B. 肝细胞气球样变
C. 肝细胞灶状坏死
D. 中性粒细胞浸润
E. 肝细胞透明小体形成

136. 酒精性肝病的最常见病变是
A. 肝细胞脂肪变性
B. 肝细胞气球样变
C. 肝细胞大片坏死
D. 大量淋巴细胞浸润
E. 肝细胞亚大片坏死

137. 不符合胆石症的病因发病的描述是
A. 胆汁理化性状的改变
B. 胆汁淤滞
C. 细菌感染
D. 常为单一因素引起
E. 胆色素和胆固醇析出

138. 胆道泥沙样结石是
A. 色素性结石
B. 胆固醇性结石
C. 混合性结石
D. 胆囊结石
E. 胆管结石

139. 符合我国胆石症的描述是
A. 胆固醇性结石多见
B. 混合性结石和色素性结石多见
C. 胆囊结石少见
D. 肝内胆管结石多见
E. 胆道蛔虫引起的结石多见

140. 符合胆道炎症病因的描述是
A. 多由大肠杆菌感染引起
B. 多由葡萄球菌感染引起
C. 多由链球菌感染引起
D. 多由胆道蛔虫引起
E. 多经血道、淋巴道感染

141. 符合急性胆管炎的描述是
A. 胆管以淋巴细胞浸润为主
B. 胆管壁纤维化
C. 胆管上皮变性坏死
D. 胆管上皮异型增生
E. 胆管周围纤维组织增生

142. 符合慢性胆囊炎的描述是
A. 常合并胆囊结石
B. 胆囊壁坏疽
C. 胆囊壁以中性粒细胞浸润为主
D. 胆囊粘膜溃疡形成
E. 浆膜纤维素渗出

143. 不符合肝外胆管癌的描述是
A. 可引起黄疸和胆道感染
B. 与胆道结石有关

C. 与先天性胆管扩张有关
D. 以鳞癌多见
E. 以胆囊管癌多见

144. 胆囊癌最常见的组织学类型是
A. 鳞癌
B. 腺鳞癌
C. 高分化腺癌
D. 类癌
E. 未分化癌

145. 不符合急性胰腺炎的描述是
A. 引起休克
B. 合并急性腹膜炎
C. 血清及尿中淀粉酶含量升高
D. 血清钙及钾离子浓度降低
E. 血清碱性磷酸酶活力增高

146. 不符合急性水肿性胰腺炎的描述是
A. 多局限于胰尾
B. 胰腺弥漫受累
C. 可见局灶性脂肪坏死
D. 多数预后较好
E. 少数发展为出血性胰腺炎

147. 不符合慢性胰腺炎病理特点的描述是
A. 胰腺质地变硬
B. 腺泡萎缩
C. 间质纤维组织增生
D. 可见广泛性坏死
E. 可形成假囊肿

148. 符合慢性胰腺炎的病变是
A. 间质充血水肿
B. 腺泡及胰岛组织萎缩消失，伴有纤维化
C. 胰腺组织结构模糊不清
D. 胰腺周围脂肪钙皂形成
E. 间质中性粒细胞浸润

149. 急性出血性胰腺炎的病变特点是
A. 间质充血水肿
B. 腺泡及胰岛组织萎缩消失，伴有纤维化
C. 胰腺组织大片坏死
D. 胰腺周围脂肪钙皂形成
E. 间质单核细胞浸润

150. 符合急性胰腺炎病因发病的描述是
A. 由细菌感染引起
B. 由缺血引起
C. 与胃酸分泌过高有关
D. 由胰蛋白酶自身消化引起
E. 由胆汁腐蚀引起

151. 胰腺癌最常见的好发部位是
A. 胰尾
B. 胰体
C. 胰头
D. 胰管
E. 以上都不是

152. 不符合胰腺癌的描述是
A. 腺管状腺癌多见
B. CA19-9 是较好的肿瘤标记物
C. 胰头癌、壶腹癌和胆总管下端癌治疗方案相同
D. 胰腺癌大多数预后良好
E. 起病隐匿

153. gastritis
A. 胃癌
B. 胃溃疡
C. 胃炎

D. 胃出血
E. 阑尾炎

154. peptic ulcer
A. 胃溃疡
B. 十二指肠溃疡
C. 溃疡穿孔
D. 幽门梗阻
E. 消化溃疡

155. appendictis
A. 肠炎
B. 胃炎
C. 阑尾炎
D. 肝炎
E. 食管炎

156. fulminant hepatitis
A. 急性肝炎
B. 活动性肝炎
C. 持续性肝炎
D. 重型肝炎
E. 普通型肝炎

157. portal cirrhosis
A. 肝硬化
B. 门脉性肝硬化
C. 坏死后性肝硬化
D. 胆管炎
E. 肝纤维化

158. cholelithiasis
A. 胆管炎
B. 胆囊炎
C. 胆管纤维化
D. 胆石症
E. 胆囊结石

159. pancreatitis
A. 胰腺炎
B. 胰腺癌
C. 胆管炎
D. 胆管癌
E. 胆囊炎

160. hepatocellular carcinoma
A. 胰腺癌
B. 胆囊癌
C. 胆管上皮癌
D. 肝细胞癌
E. 混合性肝癌

英文题

1. After a long weekend of binge drinking, a 44-year-old man has a sudden onset of severe abdominal pain. On physical examination the pain is centered in the mid-epigastric region with radiation to the back, though there is marked diffuse tenderness in all quadrants. Bowel sounds are absent. No masses are palpable. Laboratory studies show his serum amylase and lipase are markedly increased. His symptoms become worse, and he passes away 2 days later. An autopsy is performed and the pancreas is shown above. Which of the following microscopic findings is most likely to be present in this patient?



A. Fat necrosis
B. Prominent lymphocytes infiltration
C. Calcium soap
D. Fibrosis
E. None of above

2. A 72-year-old woman notes increasing jaundice and nausea for the past month. She reports 10 Kg weight loss during the past 5 months. On physical examination she is afebrile, but scleral icterus is present. There is no abdominal pain on palpation. She has active bowel sounds. A stool sample tested for occult blood is negative. Laboratory tests reveal normal levels of total protein and albumin, and increased levels of serum alkaline phosphatase, AST, ALT, total bilirubin, and direct bilirubin. Her serum level of CA19-9 antigen is also increased. Which of the following conditions is she most likely to have?

A. Carcinoma of the head of the pancreas
B. Chronic pancreatits
C. Chronic hepatitis D
D. holelithiasis
E. Colorectal carcinoma metastasizes to the liver

3. A 50 year-old woman comes to her physician because of colicky right upper quadrant pain for the past week. She has nausea, but no vomiting or diarrhea. On physical examination, she is afebrile. There is marked tenderness in the right upper quadrant. The liver span is normal. An abdominal ultrasound scan shows calculi within the lumen of the gallbladder, and the gallbladder wall appears thickened. The patient's gallbladder is removed by laparoscopic cholecystectomy and has the appearance shown above. Which of the following mechanisms is most likely to have played a role in development of this disease?




A. Ascaris lumbricoides within bile ducts
B. Bile supersaturated with cholesterol
C. Active gallbladder motility
D. Chronic hemolytic syndrome
E. Ingestion of foods rich in fat

4. A 45-year-old man sees his physician because he has had nausea and colicky right upper quadrant pain for the past 2 days. On physical examination, his temperature is 38.7℃. No scleral icterus or generalized jaundice is noted. Laboratory studies

show a WBC count of 10.5×10^9/L with 75% segmented neutrophils, 8% bands, 10% lymphocytes, and 7% monocytes. Serum levels of AST and ALT are within normal limits. Which of the following is the most likely diagnosis?

A. Acute HAV infection
B. Acute appendicitis
C. Acute pancreatitis
D. Acute cholecystitis
E. Acute perforation of a peptic ulcer

5. A 44-year-old farmer from south China presents with right upper quadrant abdominal pain and is found to have ascites. CT scan reveals a nodular liver. Laboratory tests reveal increased levels of AST and ALT together with decreased levels of albumin and total protein. A liver biopsy is performed and findings are as shown above. Which of the following is the most likely cause of his disease?



A. Mycobacterium tuberculosis
B. Amebic protozoa
C. Hydatid worm
D. Distomatosis hepatic
E. Schistosomiasis

6. A 18-year-old woman has had progressive malaise for the past year. She has become increasingly obtunded over the past week. On physical examination she is afebrile. On ocular examination, corneal Kayser-Fleischer rings are notices. Laboratory studies show increased level of plasma ammonia together with increased levels of serum total bilirubin, direct bilirubin, AST, and ALT. Her serum ceruloplasmin is high. The antimitochondrial antibody test is negative. A liver biopsy is performed. Which of the following microscopic findings is most likely to be present in this woman?

A. Increased copper deposition
B. Increased iron deposition
C. Increased lipid deposition
D. Increased glycogen deposition
E. Increased pigment deposition

7. A previously healthy 38-year-old woman has had become increasingly obtunded. On physical examination, she has scleral icterus and her temperature is 38.9°C. Her blood pressure is 110/55mm Hg. Laboratory findings show a prolonged prothrombin time, elevated levels of serum ALT, AST, and ammonia and decreased levels of albumin and total protein. Her symptoms become worse with continuously increasing serum levels of liver enzymes

and ammonia and dies within 2 weeks. An autopsy is performed and reveals a soft and shrunken liver. Which of the following microscopic changes of liver is most likely to be reported in this patient's report?

A. Spotty necrosis
B. Bridging necrosis
C. Submassive necrosis
D. Massive necrosis
E. "Piecemeal" necrosis

8. A 48-year-old man with fatigue is being evaluated for a 1 year history of elevated serum liver enzymes. A liver biopsy is taken and the pathology report for this specimen states there is grade 2 inflammatory activity with bridging necrosis and stage 1 fibrosis. The term "bridging necrosis" refers to which one of the following pathologic abnormalities?

A. Congo red-positive extracellular deposits surrounding necrotic hepatocytes in acinar zone 1
B. Destruction of the limiting plate with necrosis of hepatocytes surrounding the portal triad
C. Fibrosis around the central hepatic veins with apoptosis of adjacent hepatocytes
D. Necrosis of hepatocytes extending from the portal area of one hepatic lobule to the central vein of an adjacent lobule
E. Random necrosis of individual or small clusters of hepatocytes in acinar zone 3

9. A 65-year-old man presents with increasing fatigue, weakness, anorexia, and jaundice over the past several months. Physical examination shows mild ascites and gynecomastia. CT scan shows a nodular liver. A liver biopsy reveals regenerative nodules of hepatocytes surrounded by fibrosis. Which of the following is the source of the excess collagen deposited in these fibrotic bands?

A. Hepatoeytes
B. Kupffer cells
C. Perisinusoidal stellate cells
D. Endothelial cells
E. Bile duct epithelial cells

10. An epidemiologic study is conducted on patients infected with HBV in Singapore. These patients are followed for 10 years from the time of diagnosis. Historical data is collected to determine the mode of transmission of HBV. The patients receive periodic serologic testing for HBsAg, anti-HBs, and anti-HBc, as well as evaluation of serum for total bilirubin, AST, ALT, alkaline phosphatase, and prothrombin time. The study identifies a subset of patients who are positive for HBsAg and negative for anti-HBs antibody dur-


ing the 10 year period of the study. However, these patients lack symptoms and have normal total serum bilirubin, AST, ALT, alkaline phosphatase, and albumin. Which of the following modes of transmission of HBV is most likely occurred in these patients?

A. Blood transfusion
B. Heterosexual transmission
C. Homosexual transmission
D. Enterical transmission
E. Vertical transmission

11. A 53-year-old man comes to the emergency department because of marked hematemesis that has continued for the past 3 hours. On physical examination, he has a temperature of 35.9℃, pulse 112/min, respirations 26/min, and blood pressure 90/45 mm Hg. He has a distended abdomen with a fluid wave, and the spleen tip is palpable. Which of the following diseases is most likely to be present in this patient?


A. Fulminate hepatitis
B. Liver Cirrhosis
C. Hepatocellular carcinoma
D. Cholangiocarcinoma
E. Cholelithiasis

12. A 45-year-old woman from south China experienced increasing malaise and 10kg weight loss in the last year of her life. She became increasingly obtunded, lapsed into a coma, and died. At autopsy, the liver has the gross appearance shown above. Ingestion of which of the following is most likely to have played a role in the development of this condition?



A. Not fully cooked river-snail
B. Mildewed peanuts
C. Contraceptive agent
D. Ferrous sulfate
E. Nitrites

13. A 51-year-old man has had increasing malaise and swelling of the lower legs for the past 4 months. On physical examination, he is afebrile and normotensive. There is pitting edema to the knees. The abdomen is slightly distended with a fluid wave, but there is no tenderness. The liver span is increased. Laboratory studies show decreased levels of total serum protein and albumin, increased serum levels of AST, ALT, and total bilirubin. A liver biopsy is performed, and the microscopic appearance of a trichrome stained specimen is shown above. Which of the following is most likely to have caused this illness?



A. HAV infection
B. HBV infection
C. HCV infection
D. Alcohol abuse
E. Schistosome infection

14. A 53-year-old man has a long history of chronic hepatitis B and comes to the physician because of increasing malaise over the past year. He was hospitalized 1 year ago because of upper gastrointestinal hemorrhage. Physical examination reveals a firm nodular liver. Laboratory findings show a deceased level of serum albumin and prolonged prothrombin time. Which of the following additional physical examination findings is most likely to be present?
A. Spider nevi
B. Ascites
C. Caput medusae
D. Palpable spleen tip
E. All of above

15. After experiencing malaise and increasing icterus for 6 weeks, a 42-year-old man sees his physician. Physical examination shows jaundice, but there are no other significant findings. Serologic test results are negative for IgM anti-HAV and anti–HCV, and positive for HBsAg. He receives treatment and is asked to come back 6 weeks later. Which of the following serologic test results can be used as a marker for recovery from this disease?
A. Positive Anti-HBs antibody
B. Positive Anti-HBc antibody
C. Positive Anti-HBe antibody
D. Positive HBeAg and Anti-HBc antibody
E. Positive Anti-HBc antibody and Anti-HBe antibody

16. A study of patients who are infected with hepatitis virus A, B, C, D or E is conducted. The patients are categorized according to viral type of and are followed over the next 10 years. They receive periodic serologic testing to determine whether they are producing antibodies to the virus with which they were infected. Analysis of the data shows that a subset of patients progressed to chronic hepatitis and then to cirrhosis. In which of the following types of viral hepatitis are such conditions most likely to develop?
A. HAV, HBV and HCV
B. HBV, HCV and HDV
C. HCV, HDV and HEV
D. HAV, HDV and HEV

E. HAV, HBV and HEV

17. Ten years ago, a 32-year-old man saw a physician because he had experienced malaise, fever, and jaundice for 2 weeks. On physical examination there is mild right upper quadrant tenderness. Serologic test results are positive for HBsAg, HBV DNA, and IgG anti-HBc. The patient receives treatment and his symptoms are relieved but his serum transaminases are still abnormal after 6 months. During the following 7 years, the patients often felt fatigue and anorexia and was sent to an emergency department because of hematemesis and ascites. Serologic tests results were similar to those reported earlier. Sclerotherapy to treat esophageal varices was performed. Three years after this episode, he sees a physician because of a 5-kg weight loss, worsening abdominal pain, and rapid enlargement of the abdomen over the past month. Physical examination shows an increased liver span. Which of the following laboratory tests is most likely to be diagnostic of this last phase of his disease?
    A. Prolonged prothrombin time
    B. Increased blood ammonia level
    C. Elevated serum CA125 level
    D. Elevated serum alpha-fetoprotein level
    E. Elevated serum CEA level

18. A 10-year-old boy is brought to the physician because of listlessness, malaise, and mild fever of 5 days duration. Dark urine is observed by one of the caregivers. On physical examination, there is mild scleral icterus and mild right upper quadrant tenderness. Laboratory studies show increased levels of AST, ALT, total bilirubin and direct bilirubin, but total protein and albumin are within normal limits. Serologic tests are positive for anti-HBs, anti-HBc, IgM anti-HAV, and IgG anti-HAV, but negative for HBsAg, IgM anti-HBc, anti HCV and anti HDV. Based on these findings, which of the following is true?
    A. He now has an acute HAV infection
    B. He now has an acute HBV infection
    C. He inow has a chronic HAV infection
    D. He inow has a chronic HBV infection
    E. None of above

19. A 48-year-old Chinese man has noticed increasing abdominal girth and a yellowish color to his skin over the past 5 months. On physical examination, he has scleral icterus and generalized jaundice. His abdomen is distended, and a fluid wave is present. Laboratory studies include markedly elevated total serum bilirubin, direct bilirubin, AST, ALT, and a decreased albumin level. The prothrombin time is prolonged. The blood ammonia level is also elevated. No mass is noted

with ultrasound scan. Serologic tests are performed. Which of the following is the most likely serologic finding in this patient?

A. Positive HBsAg
B. Positive IgM anti-HAV
C. Positive anti-HBs
D. Elevated AFP level
E. Elevated CEA level

20. A 41-year-old man is found in an obtunded state and taken to the hospital. On admission physical examination shows he is icteric. His abdomen is enlarged with a fluid wave. An abdominal CT scan shows extensive intraperitoneal fluid and a uniformly enlarged liver. Laboratory studies show slightly decreased total protein and albumin, and obviously elevated serum levels of total bilirubin, AST, ALT, alkaline phosphatase and ammonia. Serologic test results are negative for HBs Ag, IgM anti-HAV, anti-HBs, anti-HBc and anti-HCV. A liver biopsy is performed which shows abundant Mallory's hyaline, neutrophilic infiltrates, necrosis of hepatocytes, and extensive macrovesicular steatosis. Which of the following is the most likely cause of these changes?

A. Hepatitis D virus
B. Hepatitis E virus
C. Alcoholic abuse
D. Increased serum ceruloplasmin lelvel
E. Hypercholesterolemia

21. A 54-year-old man has had malaise with a 10 kg weight loss over the past 6 months. On physical examination he has a firm, nodular liver edge. His stool is negative for occult blood. Laboratory studies are positive for hepatitis B surface antigen, but serologies for hepatitis B surface antibody, hepatitis A IgM antibody, and hepatitis C antibody are negative. His serum alpha-fetoprotein is 109 ng/mL. Which of the following abnormalities is the patient not likely to have?

A. Chronic hepatitis B
B. Liver cirrhosis
C. Ground-glass hepatocytes
D. Hepatocellular carcinoma
E. Bile duct adenocarcinoma

22. A 40-year-old man has a 20-year history of chronic alcohol abuse. On physical examination his liver edge is firm on palpation of the abdomen, but the liver does not appear to be enlarged. An abdominal CT scan reveals that the liver has cirrhosis. Which of the following consequences is the patient most likely to be at risk for?

A. Cholelithiasis
B. Hemangioma
C. Hepatic adenoma
D. Cholangiocarcinoma
E. Hepatocellular carcinoma

23. A 42-year-old man has had a fever and

a productive cough for the past 3 days. On physical examination there is dullness to percussion over the right upper lung. His temperature is 37.9 C. A chest radiograph reveals right upper lobe consolidation. Laboratory findings show a slightly decreased level of serum total protein and albumin, increased levels of AST and ALT, and a prolonged prothrombin time. He does not respond to antibiotic therapy and dies. He is found at autopsy to have an enlarged liver that microscopically shows most of the hepatocytes to be filled with large lipid droplets. Which of the following histologic changes is most likely also present in this specimen?

A. Ground-glass hepatocytes
B. Prominent lymphoid aggregates in the portal tract
C. Mallory bodies
D. Massive necrosis
E. Hepatocyte dysplasia

24. A group of undergraduate students plan to donate blood. Among them, a 20-year-old girl is found to be positive for HBsAg and is therefore excluded as a blood donor. Further work-up reveals no significant physical examination findings. Laboratory values for total serum bilirubin, AST, ALT, alkaline phosphatase, and albumin are normal. Serologic test results are negative for IgM anti-HAV, anti-HBc, and anti-HCV. Repeat testing 6 months later yields the same results. Which of the following is true for this patient?

A. She acquired this infection through intravenous drug use
B. She acquired this infection from enteric transmission
C. She will soon develop chronic hepatitis
D. This is probably a chronic carrier state
E. She should get a hepatitis B vaccination series

25. A 41-year-old woman has had increasing malaise and nausea for the past 2 weeks. On physical examination, she has icterus and mild right upper quadrant tenderness. A scar is noted on the left upper quadrant. The patient reports that she had her spleen removed 2 years ago following a traffic accident. Laboratory studies show increased levels of serum AST and ALT, as well as increased total and direct bilirubin. She continues to have malaise for the next year. A liver biopsy is performed. The specimen shows spotty necrosis, mild steatosis of hepatocytes, and lymphoid aggregates in the portal tract. An infection with which of the following viruses is most likely to produce these findings?

A. HAV
B. HBV
C. HCV

D. HDV

E. HEV

26. A 12-year-old boy presents with malaise, fatigue, and loss of appetite for 4 days. His mother reports that he has had a meal with fried river snail at lake side 3 weeks ago. On physical examination, he has mild scleral icterus. Laboratory studies show slightly elevated serum AST and ALT. The total bilirubin concentration and the direct bilirubin concentration are also increased. His symptoms gradually subside over the next 3 weeks. Which of the following serologic test results is most likely to be positive in this patient?

A. Anti-HAV


B. Anti-HBs

C. Anti-HBc

D. Anti-HCV

E. Anti-HDV

27. A 56-year-old man from Henan province has experienced fatigue and a 15-kg weight loss over the past 5 months. Physical examination yields no remarkable findings. Laboratory test results are positive for HBsAg and negative for anti-HCV and anti-HAV and there is an increased level of serum alpha fetoprotein. An abdominal CT scan shows a 10-cm solid mass in the left lobe of the liver which is nodular. A liver biopsy of the lesion is performed and shown above. Which of the following is not true for this patient?



A. The mass is a primary hepatocellular carcinoma

B. The mass developed in the setting of cirrhosis

C. High fat diet is an accelator for the development of this mass

D. Repeated cycles of liver cell necrosis and regeneration caused by HBV is the mechanism for the development of the mass

E. The prognosis is very poor

28. To your knowledge, for patients known to be infected with various forms of infectious hepatitis, which of the following is the best predictor for the development of chronic liver disease and progression to cirrhosis?

A. Form of the infected hepatitis virus (*HERE DO YOU MEAN "TYPE" OF VIRUS??*)

B. Degree of transaminase enzymes are increased

C. Length of time that transaminase enzymes remain elevated

D. Presence of chronic inflammatory cells in the portal tract

E. Presence of chronic inflammatory cells in the sinusoids

29. A 44-year-old man has been feeling fatigued for the past 4 months. He remembers that he had jaundice about 10 years ago, but that resolved 2 months later and he has been healthy since. On physical examination there are no remarkable findings. Laboratory studies show his hemoglobin is 14.0 g/dL. A chemistry panel reveals normal serum electrolytes, but he has a decreased level of total protein and albumin, an increased level of ALT and AST, as well as a slightly elevated level of total bilirubin and direct bilirubin. A liver biopsy is performed and microscopic examination shows disruption of the limiting plate of hepatocytes with extension of inflammation into the lobules from the triads. There is focal ballooning degeneration of hepatocytes. Which of the following laboratory test findings is most characteristic of his disease?

A. Increased serum alpha fetoprotein

B. Positive hepatitis B surface antibody

C. Positive hepatitis B surface antigen

D. Positive hepatitis B core antigen

E. Positive hepatitis A antigen

30. A 30-year-old man has had nausea and vague epigastric abdominal pain for the past 24 hours, but now the pain has become more severe and localized to right lower abdomen. On physical examination, his temperature is 38.5°C. There is haphalgesia and rebound tenderness in the right lower quadrant. An appendectomy is performed. On gross examination, the serosal surface of appendix is congested and is covered by exudate. The appendix wall is slightly thickened at the base and the lumen is filled with pus but no fecalith is noted. Microscopically, the mass is composed of nests of cells with uniform small round nuclei and cytoplasm with small purple granules. The cytoplasm is positive with antibody to chromogranin on immunohistochemical staining. Which of the following is the most likely diagnosis?

A. Acute appendicitis

B. Chronic appendicitis

C. Acute appendicitis accompanied with perforation

D. Chronic appendicitis with luminal obliteration

E. Acute appendicitis secondary to carcinoid tumor

31. A 25-year-old man has had nausea and vague epigastric abdominal pain for the past 2 days, but now the pain is localizing to the right lower quadrant. His tem-


perature is 38.7°C. On palpation, there is diffuse abdominal tenderness and rebound tenderness. Paracentesis yields 300mL of cloudy yellow fluid. A Gram stain of the fluid shows gram-negative rods. An emergency surgery is performed and the appendix is removed. Which of the following is the most likely diagnosis?

A. Acute appendicitis
B. Chronic appendicitis
C. Acute appendicitis accompanied by perforation
D. Chronic appendicitis with luminal obliteration
E. Carcinoid tumor

32. A 20-year-old woman has had nausea and vague epigastric abdominal pain for the past 24 hours, but now the pain has become more severe and localized to right lower abdomen. On physical examination, her temperature is 38.5°C. There is haphalgesia and rebound tenderness in the right lower quadrant. A stool sample is negative for occult blood. Abdominal plain film radiographs show no free air. The result of a serum pregnancy test is negative. Which of the following laboratory findings is most likely present in this patient?

A. Hyperamylasemia
B. Increased serum carcinoembryonic antigen
C. Positive urea breath test
D. Leukocytosis
E. Clostridium difficile toxin in stool sample

33. A 53-year-old woman has been feeling increasingly tired for 3 months. She has had vague abdomimal pain for the past month. On physical examination, her stool is positive for occult blood. Abnormal laboratory findings include an elevated CEA level. Colonoscopy is performed and biopsy is taken from a 4 cm ulcerative mass in the cecum. The microscopic features are shown above. Which of the following is the most likely diagnosis?



A. Mucinous carcinoma
B. Signet ring cell carcinoma
C. Medullary carcinoma
D. Well differentiated adenocarcinoma
E. Small cell carcinoma

34. A 65-year-old woman has had increasing abdominal distention for the past 6 weeks. On physical examination, there is an abdominal fluid wave and bowel sounds are present. Paracentesis yields 1000mL

of slightly cloudy serous fluid. Cytologic examination of the fluid shows malignant cells consistent with adenocarcinoma. Further workup reveals a 7 cm circumferential mass in the transverse colon. Which of the following conditions has most likely occurred in this patient?

A. Lymphangitic spread of colon adenocarcinoma
B. Hematogenous dissemination of colon adenocarcinoma
C. Implantation metastasis of colon adenocarcinoma
D. Local spreads of colon adenocarcinoma to the serosal surface
E. Both C and D

35. A 53-year-old woman saw her physician because she felt increasingly tired and listless over the past 5 months. On physical examination, there were no remarkable findings except for a positive result on stool guaiac testing. Laboratory tests showed an elevated CEA level. Colonoscopy was performed, followed by partial colectomy. After this operation, the CEA level decreased back to a normal level. The patient refused further treatment. Three years latter, in a follow-up study, it is noted that the serum CEA level of the patient is elevated again. An ultrasound scan reveals two masses in the liver of 3 and 5 cm in greatest dimension. Which of the following conditions has most likely occurred in this patient?

A. Lymphangitic spread of colon adenocarcinoma
B. Hematogenous dissemination of colon adenocarcinoma
C. Implantation metastasis of colon adenocarcinoma to the liver
D. Heterochronic carcinoma of colon and liver
E. None of above

36. A 70-year-old man sees his physician for change in bowel habits. On physical examination, there are no remarkable findings. A colonoscopy is performed and shows a 5-cm sessile mass in the upper portion of the descending colon. The histologic examination of a biopsy specimen from the lesion confirms it is a benign neoplasm with villiform structure and moderate to severe epithelial dysplasia. The patient refuses further workup and treatment. Five years later, he is see in the emergency room because of intestinal obstruction. On surgical exploration, there is a 7 cm mass encircling the descending colon and multiple enlarged lymph nodes are detected in the mesentery. Which of the following neoplasms is he now most likely to have?

A. Villous adenoma
B. Squamous cell carcinoma


C. Adenocarcinoma
D. Gastrointestinal stromal tumor
E. Carcinoid tumor

37. A 19-year-old man is advised to see his physician because genetic screening has detected a disease in other family members. On physical examination, a stool sample is positive for occult blood. A colonoscopy is performed, followed by a colectomy. The gross appearance of the mucosal surface of the colectomy specimen is shown above. Which of the following genetic changes is the patient most likely to have?



A. Germ-line mutation of APC gene
B. p53 gene mutation
C. K-RAS gene mutation
D. Defection of DNA mismatch repair genes
E. Germ-line mutation of p16 gene

38. A 45-year-old man has a routine health maintenance examination. A stool sample is positive for occult blood. On colonoscopy, an ulcerative lesion is seen projecting into the cecum. The microscopic appearance of a section of the excised lesion is shown in the figure above. Which of the following molecular biologic events is thought to be most critical in the development of such lesions?



A. Mutation of p53 gene
B. Microsatellites instability
C. Germ-line transmission of a defective RB gene
D. Amplification of ERBB2 gene
E. Germ-line transmission of a defective APC gene


39. A 55-year-old man has a routine health maintenance examination. Physical examination is unremarkable except that the stool sample is positive for occult blood. Colonoscopy is performed and a 1.5 cm pedunculated polyp with erosion is noted in the sigmoid colon. Microscopic examination reveals a tubulovillous adenoma. Which of the following is not true?

A. This is a precurosor lesion of colon cancer
B. High fat diet is a risk factor
C. Polypectomy is adequate
D. Colonoscopy should be performed on his family members to exclude colon

cancer

E. It is more common in the left-side colon than that in right-side colon

40. A 20-year-old man is healthy but has a family history of colon cancer at a young age. There are no abnormal physical examination findings. He undergoes colonoscopy and there are countless polyps ranging in size from 0.2 to 1cm. A representative microscopic picture of these polyps is shown above. Which of the following diseases is he most likely to have?



A. Familial adenomatous polyposis

B. Hereditary non-polyposis colon carcinoma syndrome

C. Carcinoid syndrome

D. Down syndrome

E. Multiple endocrine neoplasia

41. A 57-year-old woman comes to her physician for a routine health maintenance examination. The only abnormal finding is a stool specimen that contains occult blood. Colonoscopy shows two polyps, one is in the descending colon and is less than 0.5 cm in diameter with a smooth surface. The other is in the sigmoid colon, 1.5-cm in diameter with a 0.5-cm stalk. The polyp is removed, and its histology shows a gland-forming neoplasm with severe epithelial dysplasia. There is no evidence that the tumor has extended beyond the muscularis mucosa. When the physician discusses these findings with the patient, which of the following statements is most appropriate?

A. You have genomic mutation of the APC gene

B. Your family members are in great risk of developing colon cancer

C. You will develop many polyps within the next few years

D. You are in great risk of developing endometrial cancer

E. You need not to have additional examination to look for metastases from this lesion

42. A 70-year-old previously healthy man sees his physician for blood streaked stool. Physical examination is unremarkable. His stool is positive for blood but no lesion is palpated on digital rectal examination. There is no tenderness and bowel sounds are active. A cholecystectomy scar is present in the right upper quadrant. A colonoscopy is performed. About 20 cm above the anal verge，there is a 4.5 cm mass with superficial ulceration pro-


truding into the lumen. Which of the following risk factors was most likely to have been present for development of this lesion?

A. Prior abdominal surgery
B. High carbohydrate diet
C. High fat diet
D. Ulcerative colitis
E. Crohn's disease

43. A 45-year-old man has had recurrent low volume, bloody, mucoid diarrhea for 13 years accompanied by lower abdominal pain. Physical examination only reveals mild diffuse lower abdominal tenderness but no masses. His stool is positive for occult blood. Colonoscopy reveals erythematous, friable colonic mucosa extending from the rectum to the splenic flexure with formation of pseudopolyps. Colonic biopsies reveal mucosal ulceration with crypt abscesses and low grade dysplasia. Which of the following complications is he most likely to develop?

A. Bowel perforation and peritonitis
B. Perirectal fistula formation
C. Ischemic bowel necrosis
D. Intestinal lymphoma
E. Colonic adenocarcinoma

44. A 62-year-old man comes to his physician because of rectal bleeding. No other symptoms are present. Physical examination is unremarkable. Laboratory tests show an elevated CEA level. Colonoscopy is performed and there is a 3 cm ulcerative mass which occupies up to 3/4 of the colonic wall. The mass is resected and the gross findings are as shown above. Which of the following is the most likely diagnosis?



A. Hyperplastic polyp
B. Tubular adenoma
C. Villous adenoma
D. Adenocarcinoma
E. Squamous cell carcinoma

45. A 45-year-old man is having a routine physical examination because of a history of colon cancer in his family. He has no symptoms and physical examination is unremarkable. A laboratory study shows his stool is positive for occult blood. Colonoscopy is performed. Six polyps are found in the ascending and transverse colon. Two of them are pedunculated tubular adenomas less than 0.5 cm in diameter, three are 1 cm tubulovillous adenomas, and one is a 2.5 cm sessile villous adenoma in the ascending colon containing a focus of well differentiated ade-

nocarcinoma. Which of the following is the most likely diagnosis?

A. Sporadic colon adenocarcinoma
B. Hereditary non-polyposis colon carcinoma
C. Adenocarcinoma arising in a background of chronic ulcerative colitis
D. Adenocarcinoma arising in a background of familial adenomatous polyposis
E. None of above

46. A 27-year-old man presents with intermittent cramping abdominal pain and low-volume diarrhea for several weeks. On physical examination, he is afebrile, and there is mild lower abdominal tenderness but no masses, and bowel sounds are present. A stool sample is positive for occult blood. The symptoms subside within 1 week. Six months later, the abdominal pain recurs together with perirectal fistula formation. Colonoscopy shows areas of mucosal edema and ulceration in the transverse colon and rectum, but the mucosa between these affected sites appears normal. Microscopic examination of a biopsy specimen from an ulcerated area shows a patchy acute and chronic inflammatory infiltrate, crypt abscesses, and occasional noncaseating granulomas. Which of the following disease processes best explains these findings?

A. Sarcoidosis
B. Amebiasis
C. Tuberculosis
D. Crohn disease
E. Ulcerative colitis

47. A 25-year-old man has noted cramping abdominal pain for the past week associated with fever and low-volume diarrhea. On physical examination, there is right lower quadrant tenderness. Bowel sounds are present. His stool is positive for occult blood. A colonoscopy reveals mucosal edema and ulceration in the ascending colon, but the transverse and descending portions of the colon are not affected. A biospy specimen reveals chronic inflammation with non-caseating granulomas. Which of the following is the most likely diagnosis?


A. Ulcerative colitis
B. Ulcerative type adenocarcinoma
C. Regional enteritis
D. Tuberculosis
E. Amebiasis

48. A 31-year-old man presents with a history of intermittent bloody diarrhea for 3 years. On physical examination, there is no abdominal tenderness or masses. Laboratory studies show no ova or parasites in the stool but it is positive for occult blood. Colonoscopy shows friable, erythematous mucosa with numerous poly-

poid lesions extending from the rectum to the mid-transverse colon. A biopsy specimen shows acute mucosal inflammation with crypt abscesses and epithelial cell necrosis. Which of the following is the most likely diagnosis?

A. Ulcerative colitis with pseudopolyps
B. Crohn disease with pseudopolyps
C. Familial adenomatous polyposis
D. Bacillary dysentery
E. Ischemic colitis

49. A 25-year-old woman has had bloody, mucoid, low-volume diarrhea for the past 7 weeks. She has 5 -7 stools per day. She has no other major medical problems. On physical examination there are no lesions palpable on digital rectal examination, and only mucus and blood with few leukocytes are found. Colonoscopy reveals a friable, erythematous mucosa with focal ulceration that extends from the rectum to the sigmoid colon. Biopsies are taken and a representative photo is shown above. Which of the following histologic terms should be used to describe the morphologic changes highlighted by the stars?



A. Pseudomembrane
B. Crypt abscess
C. Noncaseating granuloma
D. Caseating granuloma
E. Typhoid granuloma

50. A 62-year-old woman with a long history of hypertension is hospitalized with severe congestive heart failure. Several days after admission, she develops abdominal distention followed by diarrhea. On physical examination, she is afebrile. Colonoscopy shows patchy areas of erosion in the ascending and descending colon, some with overlying tan exudate. No polyps or masses are found. Which of the following is the most likely diagnosis?

A. Ulcerative colitis
B. Pseudomembranous enteritis
C. Bacillary dysentery
D. Typhoid fever
E. Ischemic colitis

51. A 50-year-old woman was treated in the hospital for servere bronchopneumonia. She required multiple antibiotics and was intubated and mechanically ventilated. Three weeks after hospitalization, she has abdominal distention. Bowel sounds are absent, and an abdominal radiograph shows dilated loops of small bowel suggestive of ileus. She has a low volume of bloody stool. At laparotomy, a por-

tion of distal ileum and the cecum are resected. The gross appearance of the mucosal surface is shown in the figure above. Which of the following is the most likely cause of her diarrhea?



A. Ischemia
B. Entamoeba histolytica
C. Salmonella enteritidis
D. Clostridium difficile
E. Dysentery bacterium

52. A 67-year-old woman sees her physician because she has had fever and back pain for the past 2 days. Physical examination reveals tenderness at the right costovertebral angle. Laboratory studies show leukocytosis and pyuria with WBC casts. She has been receiving broad spectrum antibiotics for the past 14 days. She now develops lower abdominal pain and a severe diarrhea. Clostridium difficile toxin is identified in a stool specimen. Which of the following conditions does she most likely have now?
A. Appendicitis
B. Ischemic colitis
C. Pseudomembranous enteritis
D. Typhoid fever
E. Bacillary dysentery

53. A 19-year-old man experiences cramping abdominal pain with fever and mucinous bloody diarrhea two days after eating an unwashed fresh cucumber when enjoying his summer holiday in a small village. Physical examination shows mild diffuse abdominal pain on palpation, but there are no masses. Bowel sounds are present. He recovers completely within two week after treatment with antibiotics. Which of the following infectious organisms is most likely to have produced these findings?
A. Yersinia enterocolitica
B. Entamoeba histolytica
C. Salmonella enteritidis
D. Staphylococcus aureus
E. Dysentery bacterium

54. A 21-year-old man who is enjoying his summer holiday experiences cramping abdominal pain with fever and diarrhea two days after eating contaminated food. Physical examination shows mild diffuse abdominal pain on palpation, but there are no masses. Stool cultrure confirms the present of dysentery bacterium. Which is the most likely morphologic change in his intestine?
A. Pseudomembrane
B. Pseudopolyps
C. Linear ulcer
D. Flask ulcer
E. Noncaseating granulomas

55. A 24-year-old man experiences episodes of abdominal pain with fever and loose stools for nearly 6 months. His temperature is 37.6℃. Radiologic study reveals calcification at the right upper lobe and hilar lymph nodes. Colonoscopy reveals an ileocecal ulcer with stricture. Biopsy from this ulcer shows granulomatous inflammation. Which of the following is true for this patient?
    A. The major axis of the ulcer is probably parallel with the long axis of the intestine
    B. PPD test is probably positive
    C. The granulomas will be noncaseating granulomas
    D. The acid-fast staining will be negative
    E. He should be treated with steroids
56. 31-year-old woman sees her physician after becoming febrile and having pain in the right upper quadrant of the abdomen. She reports she previously had diarrhea after she came back from a trip to South East Asia. The diarrheal illness subsided within a couple of weeks. The stools showed mucus and streaks of blood. An abdominal ultrasound scan shows an 8-cm irregular, partly cystic mass in the right hepatic lobe. Which of the following infectious organisms is most likely to produce these findings?
    A. Tubercule bacillus
    B. Schistosoma
    C. Entamoeba histolytica
    D. Clostridium difficile
    E. Escherichia coli
57. A 36-year-old man works in an epidemiologic laboratory studying infectious agents of the gastrointestinal tract. One day, after doing experimental work, he did not follow the safety rules because he was tired. Two weeks later, he has spiking fevers and crampy abdominal pain with diarrhea. On physical examination his temperature is 38.3, pulse 100/minute, respiratory rate 19/minute, and blood pressure 100/60 mm Hg. On palpation, he has an enlarged spleen and diffuse abdominal pain without masses. Laboratory studies show a WBC count of 2330/microliter, Hb 13.8 g/dL, platelet count 280,000/microliter and elevations in serum AST and ALT levels. Which of the following organisms was he most likely using in his experimental work?
    A. Shigella flexneri
    B. Entameba histolytica
    C. Salmonella typhi
    D. Tubercule bacillus
    E. Clostridium difficile
58. A 27-year-old woman sees her physician because of abdominal pain and increasing fatigue that has developed over the past 6 months. On physical examina-

tion, her temperature is 36.6°C and she appears pale. On palpation, there is pain in the right lower quadrant of the abdomen but no rebound tenderness. There are no masses, and bowel sounds are active. Laboratory studies show a hemoglobin of 8.6g/dL, platelet count 275,000/mm^3 and WBC count 6800/mm^3. No lesions are found on upper gastrointestinal endoscopy and colonoscopy. One month later, the patient continues to experience the same abdominal pain. Which of the following is most likely to cause this patient's illness?

A. Acute appendicitis
B. Ischemic colitis
C. Typhoid fever
D. Meckel's diverticulum
E. Carcinoid

59. A 4 month old male infant is brought to the emergency room by her mother because of severe vomiting. The infant was born at term with no congenital anomalies. His mother has noted marked abdominal enlargement along with infrequent bowel movements for the past week. On physical examination his abdomen is distended but there is no tenderness. A plain film abdominal radiograph reveals marked colonic dilation. A biopsy specimen from the rectum distal to the region of dilation shows an absence of ganglion cells in the muscle wall and submucosa. Which of the following is most likely to produce these findings?

A. Constipation
B. Hirschsprung's disease
C. Intussusception
D. Volvulus
E. Crohn's disease

60. A 60-year-old man has experienced nausea and epigastric pain over the past 3 months. Physical examination is unremarkable. Upper gastrointestinal endoscopy reveals a slight elevated submucosal mass with erosions of the overlying mucosa. CT scan confirms the present of a well circumscribed mass within the gastric wall which extends to the serosa. The mass is removed and pathologic examination reveals a spindle cell tumor 8 cm in maximum dimension which is positive for CD117 staining. Which of the following neoplasm does this patient have?

A. Leiomyosarcoma
B. Gastrointestinal stromal tumor
C. Lymphoma
D. Liposarcoma
E. Schwannoma

61. A 50-year-old man has had persistent nausea for 3 years with occasional vomiting. On physical examination there are no abnormal findings. He undergoes an upper GI endoscopy, and a small area of gas-

tric mucosa is noted in the antrum which has loss of rugal folds. Biopsy from this site reveals a monoclonal proliferation of small lymphoid cells. Immunohistochemical stains show these lymphoid cells are of B cell origin. An upper GI endoscopy performed 5 years previously showed gastritis and microscopically there was chronic inflammation with the presence of Helicobacter pylori. Which of the following treatments is most likely to be the best choice for his patient?

A. Partial gastrectomy
B. Chemotherapy
C. Radiotherapy
D. Antibiotics
E. All of above

62. A 57-year-old female complaints of pelvic pain accompanied by weight loss of 6 kg over the past 5 months. Ultrasound examination reveals that both ovaries are enlarged. Serologic tests shows a markedly elevated CEA level and mildly elevated CA125 and CA19-9 levels. Exploratory laparotomy is performed. Biopsies are taken from both ovaries and sent for intraoperative analysis. A metastatic signet ring cell carcinoma is revealed on frozen section. Which of the following organs is most likely the site of the primary tumor?


A. Stomach
B. Appendix
C. Esophagus
D. Liver
E. Pancreas

63. A 55-year-old man has experienced severe epigastric pain, vomiting, and a 10-kg weight loss over the past 3 months. Physical examination reveals fixed left supraclavicular lymphadenopathy. Upper gastrointestinal endoscopy is done and confirms the present of a moderately differentiated adenocarcinoma. A palliative operation is performed to relieve his symptoms. A 6-cm ulcerative mass is located in the gastric antrum. The tumor has penetrated through the gastric wall is adherent to the transverse colon. The liver appears normal. There are numerous small white nodules on the serosal surface of intestine, spleen and peritoneum. Which of the following conditions may not have occurred in this patient's tumor?

A. Local invasion
B. Lymphatic spread
C. Hematgenous spread
D. Seeding within abdominal cavity
E. Perforation

64. A 60-year-old man has experienced severe nausea, vomiting, and a 5kg weight loss over the past 3 months. Physical examination is unremarkable. Upper gastrointestinal endoscopy is done followed by partial gastrectomy. The gross picture of the tu-

mor is shown above. Which of the following gross types does this tumor belong to?



A. Polypoid type
B. Fungating type
C. Ulcerative type
D. Infiltrative type
E. Elevated type

65. A 63-year-old man has experienced severe nausea, vomiting, and an 8-kg weight loss over the past 3 months. Physical examination reveals fixed left supraclavicular lymphadenopathy. Upper gastrointestinal endoscopy shows a 4 cm elevated mass with an irregular central ulceration. The rest of the mucosa appears pale and thin with areas of erythema. Which of the following conditions is this patient most likely to have?

A. Peptic ulcer
B. Preexisting chronic atrophic gastritis
C. Diffuse type adenocarcinoma
D. Early stage gastric carcinoma
E. Linitis plastica

66. A 50-year-old woman has a 15 year history of chronic atrophic gastritis. She has undergone endoscopic surveillance every 2-3 years. During the last endoscopic examination, a polypoid lesion of less than 1 cm is found on the lesser curvature of the antrum. A biopsy is taken and a partial gastrectomy is performed. Microscopically, it is an adenocarcinoma and is within the mucosa. Which of the following gross types does this tumor likely belong to?

A. Protruding type
B. Elevated type
C. Flat type
D. Depressed type
E. Excavated type

67. A 60-year-old man has had persistent gastric distention for 5 years with occasional vomiting. On physical examination there are no abnormal findings. He undergoes an upper GI endoscopy followed by partial gastrectomy. Pathologic examination confirms the presence of an early gastric adenocarcinoma. Which of the following conditions will most likely not be seen in this neoplasm?

A. The lamina propria is involved
B. The muscularis mucosa is involved
C. The submucosa is involved
D. The muscularis propria is involved
E. The lymph nodes are involved

68. A 60-year-old man has had anorexia,

vomiting, and vague abdominal pain accompanied by weight loss of 10 kg over the past 5 months. Physical examination reveals supraclavicular non-tender lymphadenopathy. He becomes progressively cachectic and dies. At autopsy the stomach is shrunken with thickening of the gastric wall to 1 cm and extensive overlying mucosal erosions. Multiple tan umbilicated masses from 1 to 4 cm in size are scattered over the surface of the liver. Which of the following conditions most likely preceded development of his terminal illness?


A. Human immunodeficiency virus infection
B. Use of NSAIDS
C. Chronic alcoholism
D. Autoimmune gastritis
E. A diet rich in Vitamin C



69. A 62-year-old woman has had nausea, vomiting, and mid-epigastric pain for 6 months. On physical examination, there are no significant findings. An upper gastrointestinal radiographic series shows gastric outlet obstruction. Upper gastrointestinal endoscopy shows a huge ulcerated mass at the antrum. The resection specimen is shown as above. Which of the following neoplasms is most likely to be seen under microscope?

A. Adenocarcinoma
B. Gastrointestinal stromal tumor
C. Squamous cell carcinoma
D. Non-Hodgkin lymphoma
E. Leiomyosarcoma

70. A 55-year-old man has had persistent nausea for 5 years with occasional vomiting. On physical examination there are no abnormal findings. He undergoes an upper GI endoscopy, and a small area of gastric mucosa is noted in the fundus which shows loss of rugal folds. Partial gastrectomy reveals the microscopic change shown above and the lesion is confined to submucosa. An upper GI endoscopy performed 5 years previously showed a pattern of gastritis and microscopically there was chronic inflammation with the presence of Helicobacter pylori. Which of the following interpretations is most applicable to his neoplasm?



A. Metastases are limited to regional lymph nodes
B. The incidence is high in East Asia
C. The 5-year survival rate following resection is <80%
D. It has a gross appearance of linitis plastica
E. A signet ring cell pattern is seen with light microscopy

71. A 67-year-old woman has experienced severe nausea, vomiting, early satiety, and a 5-kg weight loss over the past 3 months. Upper gastrointestinal endoscopy shows that the entire gastric mucosa is eroded and has an erythematous, cobblestone appearance. Upper gastrointestinal radiographs show that the stomach is small and shrunken with reduced movement. Which of the following is most likely to be present in this patient?
A. Advanced stage gastric carcinoma
B. Diffuse type adenocarcinoma
C. Signet-ring cell adenocarcinoma
D. Linitis plastica
E. All of above

72. A 38-year-old man has had repeated upper abdominal pain for a half year and it has become worse over the past 2 weeks. On physical examination a stool sample is positive for occult blood. An upper GI endoscopy reveals no esophageal lesions, but there is a solitary 2 cm diameter shallow, sharply demarcated ulceration of the stomach. Which of the following statements regarding this lesion is most appropriate?
A. It is probably located in the cardia
B. A gastrinoma of the pancreas is probably present
C. It is probably associated with increased gastric acid production
D. Because of its small size, a biopsy is not necessary
E. It is probably benign

73. A 46-year-old woman is sent to emergency department because of a sudden onset of abdominal pain. She has a long history of heartburn and dyspepsia. On palpation, she has severe mid-epigastric pain with guarding. Bowel sounds are reduced. An abdominal plain film radiograph shows free air under the left side of the diaphragm. The patient is immediately taken to surgery, and a perforated duodenal ulcer is repaired. Which of the following organisms is most likely to have produced these findings?
A. Campylobacter jejuni
B. Shigella flexneri
C. Entamoeba histolytica
D. Helicobacter pylori
E. Salmonella typhi

74. A 28-year-old taxi driver has had epigastric pain for over 1 year. The pain tends

to occur 2 to 3 hours after a meal and is relieved if he takes antacids or eats more food. He smokes and drinks about 150 mL of wine daily. The result of a urea breath test is positive, and a gastric biopsy specimen contains urease. He begins a 2-week course of antibiotics, but on day 7, he feels better and discontinues treatment. Several weeks later, the epigastric pain recurs. If the patient does not seek further treatment, which of the following is he less likely to develop?

A. Hematemesis
B. Pyloric obstruction
C. Melena
D. Adenocarcinoma
E. Perforation

75. A 54-year-old man has had upper abdominal pain accompanied by nausea for 5 months. He has no hematemesis. On physical examination, the stool sample is positive for occult blood. Upper GI endoscopy is performed and gastric biopsies are taken. On microscopic examination the biopsies reveal acute and chronic mucosal inflammation together with the presence of Helicobacter pylori organisms. The presence of these organisms is most likely to be associated with which of the following?

A. Duodenal peptic ulceration
B. Hypochlorhydria with atrophic gastritis
C. Development of pernicious anemia
D. Development of diffuse large B cell lymphoma
E. None of above

76. A 40-year-old man has complained for 3 months of mid-epigastric pain and nausea. On physical examination he has no abnormal findings. On upper GI endoscopy a solitary sharply demarcated 2 cm shallow gastric antral ulcer is seen. Which of the following is most likely to be present in this man?


A. Gastric achlorhydria
B. Positive serology for antinuclear antibody
C. Pernicious anemia
D. Positive urea breath test
E. Elevated serum gastrin

77. A 42-year-old woman has had nausea and epigastirc pain for the past two years. She has been feeling increasingly tired in the recent 6 months. Physical examination is unremarkable. Laboratory studies show a hemoglobin of 9.2 g/dL, hematocrit 28.5%, MCV 122 μm^3, platelet count 247,000/mm^3, and WBC count 5500/mm^3. The reticulocyte index is low. Upper GI endoscopy is performed and multiple biopsies are taken. Microscopic examination shows chronic atrophic gastritis in some of the biopsy specimens and

silver stains do not show any positive organisms. The biopsies that show atrophic gastritis are most likely taken from which portion of the stomach?

A. Pylorus
B. Antrum
C. Body
D. Cardia
E. Gastroesophageal junction

78. A middle aged man complains for 2 month of gastric distention and vague pain. He has a history of acute gastritis and had a positive urea breath test 4 years ago. His physician suggested he have a course of antibiotic treatment but he did not do so. Physical examination is unremarkable. Upper GI endoscopy shows the antral mucosa is pale，thin and interspersed with erythema. No ulceration or mass is present. A biopsy is taken and a tissue section with a special stain is shown above. Which of the following is correct?



A. It is a silver stain
B. The purple stain represents intestinal mucin
C. The blue stain represents neutral mucin
D. It is closely related to Helicobacter pylori infection
E. All of above

79. A middle aged woman has had vague abdominal pain and nausea for the past year. She has tried antacid medications for two weeks with no relief of symptoms. On physical examination there are no abnormal findings. On upper GI endoscopy, the antral mucosal is pale without ulcerations or masses. Biopsies are taken. No autoimmune antibodies are found in serologic tests and a urea breath test is positive. Which of the following microscopic changes is most likely to be present in this woman?

A. Reduced number of parietal cells
B. Reduced number of chief cells
C. Intestinal metaplasia
D. Squamous metaplasia
E. Granulomas

80. A 44-year-old woman has been feeling increasingly tired for the past 9 months. There are no remarkable findings on physical examination. Laboratory studies show a hemoglobin of 8.5g/dL, hematocrit 27.1%, MCV 120μm^3, platelet count 252,000/mm^3, and WBC count 6350/mm^3. The reticulocyte index is low. Antibodies to which of the following are most likely to be found in this patient?

A. Gliaden
B. Rheumatoid factor
C. Helicobacter pylori
D. Na+K+-ATPase
E. Intrinsic factor

81. A 45-year-old man has had vague abdominal pain and nausea for the past 2 years. This pain is unrelieved by antacid medications. On physical examination there are no abnormal findings. Upper GI endoscopy reveals antral mucosal erythema, but no ulcerations or masses. Biopsies are taken, and microscopically there is a chronic non-specific gastritis. Which of the following conditions is not likely to be present in this man?
A. Positive urea breath test
B. Normal serum gastrin level
C. Helicobacter pylori infection
D. Increasing risk of developing gastric adenocarcinoma
E. Pernicious anemia

82. A 65-year-old woman takes large quantities of Aspirin because of chronic arthritis. Recently, she has had epigastric pain with nausea and vomiting and an episode of hematemesis. There are no remarkable findings on physical examination. A gastric biopsy specimen is most likely to show which of the following lesions?
A. Epithelial dysplasia
B. Acute gastritis
C. Atrophic gastritis
D. Adenocarcinoma
E. Hypertrophic gastritis

83. A 45-year-old woman has had nausea and vomiting for several months. On physical examination, there is no abdominal tenderness, and bowel sounds are present. Upper gastrointestinal endoscopy shows erythematous areas of mucosa with thickening of the rugal folds in the gastric antrum. The microscopic appearance of a gastric biopsy specimen with a Steiner silver stain is shown in the preceding figure. Which of the following organisms is most likely to be present on the luminal surface?



A. Clostridiun difficile
B. Herpes simplex
C. Staphylococcus aureus
D. Helicobacter pylori
E. Salmonella enteritidis

84. A 25-year-old man has an acute onset of nausea and vomiting two hours after drinking one bottle of “二锅头”. Phys-

ical examination shows no remarkable findings. Endoscopically, diffuse gastric mucosal erythema with focal mucosal erosion and hemorrhage are present. Microscopic examination of gastric biopsy specimens shows increased infiltrates in the lamina propria, consisting predominately of neutrophils. No organisms are found by sliver stain. Laboratory studies show a normal serum gastrin level. Which of the following description is proper for this patient?

A. He will recover within days without further treatment
B. He will develop chronic gastritis if not treated properly
C. The complications of this disease includes pernicious anemia
D. Intestinal metaplasia is commonly seen in this disease
E. The patient is at risk for developing gastric lymphoma

85. A middle aged man has had difficulty swallowing for the past 8 months and has had a 7-kg weight loss within 3 months. Physical examination reveals fixed supraclavicular lymph nodes. Upper gastrointestinal endoscopy reveals an ulcerative mass in the mid-esophagus which almost obstructs the entire lumen. Biopsy is performed and microscopic examination confirms the presence of a squamous cell carcinoma. Which of the following statements is most likely correct?


A. The tumor is restricted to the submucosa
B. The progression is excellent
C. The tumor has metastasized via lymph vessels
D. The precursor lesion is Barrett's esophagus with dysplasia
E. None of above

86. A 53-year-old male presents with increasing dysphagia and retrosternal pain. He has also noticed a 7-kg weight loss over the past 3 months. On physical examination, there are no remarkable findings. Upper gastrointestinal endoscopy reveals a 4 cm length of narrowing of the mid-esophagus lined by coarse mucosa. Esophagectomy is performed and microscopic examination confirms the presence of an advanced stage carcinoma. Which of the following appropriately describes the gross appearance of this carcinoma?

A. Polypoid pattern
B. Fungating pattern
C. Ulcerative pattern
D. Infiltrative pattern
E. Depressed pattern

87. A 55-year-old man has had dysphagia for 6 months. On physical examination there are no abnormal findings. He undergoes an upper GI endoscopy, and a small shal-

low ulcer is noted in the middle portion of the esophagus. Biopsies from this area show a moderately differentiated squamous cell carcinoma. Esophagectomy is performed. The microscopic changes are as shown above and no metastases are found. Which of the following interpretations is most applicable to his neoplasm?




A. It has a high potential to metastasize to the liver
B. The incidence is high in Western countries
C. The 5-year survival following resection is >80%
D. It is associated with a high fruit diet
E. Its precursor lesion is Barrett's esophagus

88. A 58-year-old man has had increasing difficulty swallowing for the past 6 months. He has lost 5 kg in the past 2 months. No abnormal physical examination findings are noted. Upper GI endoscopy reveals a nearly circumferential mass with overlying ulceration in the mid esophageal region. Biopsy of the mass reveals pink polygonal cells with marked hyperchromatism and pleomorphism. Which of the following is the most likely risk factor for development of his disease?

A. Iron deficiency
B. Reflux esophagitis
C. Helicobacter pylori infection
D. Ice-cold beverage or food
E. Chronic alcoholism

89. A 50-year-old male farmer from Henan province presents with increasing dysphagia and has noticed a 6kg weight loss over the past 2 months. His grandfather and father both died in their late 50's with similar symptoms. On physical examination, there are no remarkable findings. Upper gastrointestinal endoscopy shows a 4cm ulcerative mass in the mid-esophagus that occupies 1/2 of the esophageal lumen. Esophagectomy is performed, and the gross appearance of the lesion is shown in the preceding figure. Which of the following is most likely to be seen on microscopic sections of this mass?




A. Columnar epithelium with goblet cells
B. Adenocarcinoma
C. Dense collagenous scar
D. Squamous cell carcinoma
E. Ectatic vascular channels

90. A 58-year-old man has had sour brash and substernal pain after meals for many years. For the past year, he has had increased dysphagia even when drinking liquids. On physical examination, there are no remarkable findings. Upper gastrointestinal endoscopy shows a lower esophageal mass that nearly occludes the lumen of the esophagus. Biopsy of this mass is most likely to show which of the following neoplasms?
    A. Malignant neoplasm which forms glandular structure
    B. Malignant neoplasm which is composed of spindle cells
    C. Malignant neoplasm which forms "keratin pearls"
    D. Malignant neoplasm which has "Reed-Sternberg" cells
    E. Malignant neoplasm which has neuroendocrine granules within the cytoplasm


91. A 49-year-old man has complained of "heartburn" after meals for the past decade. There are no remarkable findings on physical examination. Upper gastrointestinal endoscopy is performed, and a biopsy specimen is taken above the gastroesophageal junction. Microscopically, the mucosa shows columnar metaplasia with goblet cells. Which of the following description is not correct for this disease?
    A. It is caused by gastroesophageal reflux
    B. The diagnosis for this condition is Barrett's esophagus
    C. The morphologic changes referred to as squamous metaplasia
    D. The gross appearance of the lesion is usually velvety and erythematous
    E. The patient is at risk for developing adenocarcinoma

92. A 49-year-old man has had progressive difficulty on swallowing during the past 6 months. No significant findings are observed on physical examination. Upper gastrointestinal endoscopy shows areas of velvety erythematous mucosa above the Z-line. A biopsy specimen from the lower esophagus has the microscopic appearance shown in the preceding figure. Which of the following is the most likely consequence of this disease?



    A. Esophageal stricture
    B. Adenocarcinoma
    C. Hematemesis
    D. Esophageal laceration
    E. Squamous cell carcinoma

93. A 57-year-old woman has had burning epigastric pain after meals for more than 1 year. Physical examination shows no abnormal findings. Upper gastrointestinal endoscopy shows an erythematous patch in the lower esophageal mucosa. A biopsy specimen shows basal squamous epithelial hyperplasia, elongation of lamina propria papillae, and scattered intraepithelial neutrophils with some eosinophils. As a consequence of this patient's condition, which of the following complications is not likely to occur?
    A. Intestinal metaplasia
    B. Barrett's esophagus
    C. Squamous cell carcinoma
    D. Esophageal stricture
    E. Adenocarcinoma
94. A 35-year-old woman has experienced "heartburn" and sour regurgitation following meals for the past 2 years. There are no abnormal findings on physical examination. Upper GI endoscopy is performed. There are 2 to 3 cm long tongues of erythematous mucosa extending from the gastroesophageal junction at the Z line upward into the lower esophagus. Biopsies of this region are performed and the microscopic pictures is shown above. Which of the following interpretations is most appropriate for this woman's findings?



    A. She has a congenital anomaly
    B. Her risk for squamous cell carcinoma is increased
    C. Her consumption of alcohol is high
    D. She has chronic gastroesophageal reflux
    E. Formation of a diverticulum may occur
95. A 28-year-old HIV positive woman has had pain on swallowing for the past week. No abnormal physical examination findings are noted. Upper GI endoscopy is performed. There are multiple small ulcers in the lower esophagus. Inclusions are found in the biopsies from these ulcers. She is most likely to have infection with which of the following organisms?
    A. Helicobacter pylori
    B. Candida albicans
    C. Aspergillus
    D. Mycobacterium avium-complex
    E. Cytomegalovirus
96. A 16-year-old boy sees the physician because he has had pain for 1 week when

he swallows food. Physical examination shows no abnormal findings. Upper gastrointestinal endoscopy shows 0.5- to 0.8-cm mucosal ulcers in the region of the middle to lower esophagus. The shallow ulcers are round, sharply demarcated, and have an erythematous base. The boy has a history of acute lymphoblastic leukemia and is receiving chemotherapy. Which of the following is most likely to produce these findings?

A. Helicobacter pylori infection
B. Herpes simplex virus infection
C. Gastroesophageal reflux disease
D. Barrett's esophagus
E. Mycobacterium tuberculosis infection

97. An 18-year-old woman has had several episodes of aspiration of food associated with difficulty swallowing during the past year. On auscultation, crackles are heard at the base of the right lung. A barium swallow shows marked esophageal dilation above the level of the lower esophageal sphincter. No myenteric ganglia are present in a biopsy specimen from the lower esophagus. Which of the following is the most likely diagnosis?

A. Achalasia
B. Esophageal stricture
C. Barrett's esophagus
D. Reflux esophagitis
E. Hiatal hernia

98. A 51-year-old man has a sudden onset of massive emesis of bright red blood. On physical examination, his temperature is 36.9°C, pulse 103/min, respirations 19/min, and blood pressure 85/50mm Hg. Laboratory studies show AST 137U/L and ALT 108U/L. The serologic test result for HBsAg is positive. He has had no prior episodes of hematemesis. The hematemesis is most likely to be a consequence of which of the following?

A. Barrett esophagus
B. Esophageal laceration
C. Esophageal varices
D. Reflux esophagitis
E. Squamous cell carcinoma

99. A 35-year-old man experiences a bout of prolonged vomiting, followed by massive hematemesis. On physical examination in the emergency room, he has vital signs of T 36.7°C, P 108, RR 22, and BP 80/40 mm Hg. His heart has a regular rate and rhythm with no murmurs and his lungs are clear to auscultation. There is no abdominal tenderness or distension and bowel sounds are present. His stool is negative for occult blood. Which of the following is the most likely diagnosis?

A. Hiatal hernia
B. Esophageal laceration
C. Esophageal pulsion diverticulum
D. Reflux esophagitis
E. Esophageal squamous cell carcinoma

答案

中文题

1.D	2.E	3.A	4.C	5.B	6.B	7.B	8.E	9.E	10.B
11.E	12.D	13.C	14.D	15.B	16.C	17.B	18.D	19.E	20.E
21.C	22.D	23.C	24.D	25.D	26.C	27.A	28.B	29.C	30.A
31.D	32.C	33.D	34.B	35.C	36.A	37.A	38.B	39.B	40.C
41.D	42.D	43.A	44.E	45.A	46.E	47.D	48.B	49.D	50.A
51.E	52.A	53.D	54.C	55.C	56.A	57.C	58.B	59.A	60.D
61.B	62.A	63.D	64.A	65.B	66.A	67.C	68.D	69.C	70.A
71.E	72.A	73.D	74.B	75.A	76.E	77.B	78.C	79.D	80.E
81.E	82.B	83.A	84.D	85.D	86.A	87.E	88.C	89.D	90.A
91.C	92.E	93.A	94.C	95.A	96.A	97.B	98.A	99.A	100.B
101.A	102.A	103.C	104.A	105.E	106.A	107.D	108.E	109.B	110.E
111.C	112.B	113.C	114.B	115.B	116.D	117.C	118.E	119.A	120.D
121.B	122.A	123.A	124.B	125.A	126.A	127.C	128.E	129.C	130.A
131.E	132.C	133.E	134.E	135.B	136.A	137.D	138.A	139.B	140.A
141.C	142.A	143.A	144.C	145.E	146.B	147.D	148.B	149.C	150.D
151.C	152.D	153.C	154.E	155.C	156.D	157.B	158.D	159.A	160.D

英文题

1.A	2.A	3.B	4.D	5.E	6.A	7.D	8.D	9.C	10.E
11.B	12.B	13.B	14.E	15.A	16.B	17.D	18.A	19.A	20.C
21.E	22.E	23.C	24.D	25.?	26.?	27.C	28.A	29.C	30.E
31.C	32.D	33.A	34.E	35.B	36.C	37.A	38.B	39.D	40.A
41.E	42.C	43.E	44.D	45.B	46.D	47.C	48.A	49.B	50.E
51.D	52.C	53.E	54.A	55.B	56.C	57.C	58.D	59.B	60.B
61.D	62.A	63.C	64.A	65.B	66.A	67.D	68.D	69.A	70.B
71.E	72.E	73.D	74.D	75.A	76.D	77.C	78.D	79.C	80.E
81.E	82.B	83.D	84.A	85.D	86.D	87.D	88.E	89.D	90.A
91.C	92.B	93.C	94.D	95.E	96.B	97.A	98.C	99.B	

8　泌尿系统疾病

中文题

1. 肾小球肾炎属于
   A. 化脓性炎
   B. 纤维素性炎
   C. 肉芽肿性炎
   D. 出血性炎
   E. 非化脓性变态反应性炎
2. 不符合肾小球肾炎的描述是
   A. 可由循环免疫复合物引起
   B. 可由原位免疫复合物引起
   C. 可由肾小球固有抗原引起
   D. 可由肾小球植入抗原引起
   E. 可由细菌感染引起
3. 肾小球肾炎属于
   A. I 型变态反应
   B. II 型变态反应
   C. III 型变态反应
   D. IV 型变态反应
   E. 以上都不是
4. 不符合肾小球的描述是
   A. 再生能力强
   B. 由毛细血管丛和肾球囊构成
   C. 滤过膜由内皮细胞、基底膜和上皮细胞构成
   D. 系膜细胞具有吞噬功能
   E. 滤过膜带负电荷
5. 非免疫复合物性肾小球肾炎时，引起肾小球损伤的因素是
   A. 补体
   B. 血小板
   C. 中性粒细胞
   D. 巨噬细胞
   E. T 细胞
6. 不属于内源性抗原的是
   A. 肿瘤抗原
   B. 基底膜抗原
   C. 异种血清
   D. 核抗原
   E. Heymann 抗原
7. 可引起原位免疫复合物性肾小球肾炎的植入性抗原是
   A. 基底膜抗原
   B. Heymann 抗原
   C. 聚合性 IgG
   D. 内皮细胞膜抗原
   E. 系膜细胞膜抗原
8. 不符合循环免疫复合物引起的肾小球肾炎的描述是
   A. 由肾小球性抗原引起
   B. 由非肾小球性抗原引起
   C. 抗原抗体在血液中形成复合物
   D. 抗原抗体复合物沉积于肾小球引起炎症
   E. 免疫复合物的大小、所带电荷在发病中起重要作用
9. 急性链球菌感染后肾小球肾炎属于
   A. 弥漫性毛细血管内增生性肾小球肾炎
   B. 弥漫性新月体性肾小球肾炎

C. 弥漫性膜性肾小球肾炎
D. 弥漫性膜性增生性肾小球肾炎
E. 轻微病变性肾小球肾炎

10. 弥漫性毛细血管内增生性肾小球肾炎增生的细胞是
A. 肾小球球囊壁层上皮细胞
B. 肾小球球囊脏层上皮细胞
C. 肾小球系膜细胞和内皮细胞
D. 肾小球内皮细胞
E. 肾小球系膜细胞

11. 不符合弥漫性毛细血管内增生性肾小球肾炎的是
A. 肾小球内皮细胞增生
B. 肾小球系膜细胞增生
C. 肾小球基底膜弥漫性增厚
D. 肾小球毛细血管壁显示 IgG 和 C3 的颗粒荧光
E. 肾小球基底膜外侧有驼峰状电子致密物

12. 不符合弥漫性毛细血管内增生性肾小球肾炎的描述是
A. 可引起大红肾
B. 双肾弥漫受累
C. 发生于链球菌感染后
D. 表现为急性肾炎综合征
E. 预后差

13. 弥漫性毛细血管内增生性肾小球肾炎电镜下病变特点是
A. 毛细血管基底膜内皮细胞下见致密沉积物
B. 毛细血管基底膜上皮侧见驼峰状致密沉积物
C. 脏层上皮细胞足突融合
D. 毛细血管基底膜严重缺损
E. 毛细血管基底膜内见致密沉积物

14. 不符合弥漫性毛细血管内增生性肾小球肾炎尿液变化的是
A. 少尿无尿
B. 血尿
C. 蛋白尿
D. 管型尿
E. 脓尿

15. 绝大多数能治愈的肾小球肾炎是
A. 弥漫性膜性增生性肾小球肾炎
B. 弥漫性新月体性肾小球肾炎
C. 弥漫性毛细血管内增生性肾小球肾炎
D. 弥漫性膜性肾小球肾炎
E. 肺出血肾炎综合征

16. 弥漫性毛细血管内增生性肾小球肾炎的临床表现常为
A. 肾病综合征
B. 急性肾炎综合征
C. 慢性肾炎综合征
D. 隐匿性肾炎综合征
E. 急进性肾炎综合征

17. 弥漫性新月体性肾小球肾炎中新月体的细胞是
A. 肾小球球囊壁层上皮细胞
B. 肾小球球囊壁层上皮细胞和单核巨噬细胞
C. 肾小球球囊脏层上皮细胞和单核巨噬细胞
D. 肾小球系膜细胞和内皮细胞

E. 肾小球系膜细胞

18. 弥漫性毛细血管外增生性肾小球肾炎的主要病变为

A. 肾小球毛细血管基底膜增厚

B. 肾小球内皮细胞增生

C. 肾小球系膜细胞

D. 肾小球硬化

E. 新月体形成

19. 弥漫性新月体性肾小球肾炎常见的临床症状是

A. 肾病综合征

B. 急性肾炎综合征

C. 慢性肾炎综合征

D. 隐匿性肾炎综合征

E. 急进性肾炎综合征

20. 常导致急性肾功能衰竭的肾小球肾炎是

A. 弥漫性新月体性肾小球肾炎

B. 弥漫性系膜增生性肾小球肾炎

C. 弥漫性膜性肾小球肾炎

D. 轻微病变性肾小球肾炎

E. 局灶性肾小球肾炎

21. 不易演变成新月体性肾小球肾炎的是

A. 弥漫性毛细血管增生性肾小球肾炎

B. 抗基底膜性肾小球肾炎

C. 弥漫性膜性增生性肾小球肾炎

D. 弥漫性系膜增生性肾小球肾炎

E. 轻微病变性肾小球肾炎

22. 肺出血肾炎综合征肾脏的主要病变是

A. 新月体形成

B. 肾小球毛细血管基底膜增厚

C. 肾小球内皮细胞和系膜细胞弥漫性增生

D. 肾小球系膜细胞及基质增生

E. 系膜细胞增生

23. 免疫荧光沿基底膜呈线型分布的肾小球肾炎是

A. 弥漫性膜性肾小球肾炎

B. 轻微病变性肾小球肾炎

C. 肺出血肾炎综合征

D. 弥漫性膜性增生性肾小球肾炎

E. IgA 肾病

24. 不符合新月体性肾小球肾炎的描述是

A. 病变严重，进展快，预后差

B. 多数肾小球新月体形成

C. 新月体由肾球囊壁层上皮和巨噬细胞构成

D. 免疫荧光沿基底膜呈线性分布

E. 基底膜变性断裂

25. 弥漫性膜性肾小球肾炎的主要病理特点是

A. 肾小球系膜细胞和内皮细胞增生

B. 肾小球基底膜弥漫性增厚并有钉突形成

C. 肾小球系膜细胞增生、基底膜增厚及分层

D. 肾小球硬化

E. 肾小球无明显病变

26. 符合弥漫性膜性肾小球肾炎的描述是

A. 肾小球系膜区免疫复合物沉积

B. 肾小球基底膜内免疫复合物沉积

C. 肾小球基底膜和上皮细胞间驼峰状免疫复合物沉积

D. 肾小球基底膜和内皮细胞间免疫复

合物沉积

E. 肾小球内无免疫复合物沉积

27. 不符合弥漫性膜性肾小球肾炎的描述是

A. 肾小球基底膜弥漫性增生

B. 肾小球基底膜有钉突形成

C. 肾小球毛细血管壁有颗粒状 IgG 和 C3 荧光

D. 肾小球系膜区有团块状 IgG 荧光

E. 肾小球基底膜外侧有多数电子致密物

28. 弥漫性膜性肾小球肾炎光镜病变特点是

A. 肾球囊壁层上皮细胞增生

B. 肾小球内皮细胞增生

C. 小球毛细血管壁增厚系膜细胞和基质增生

D. 脏层上皮细胞足土消失融合

E. 肾小球毛细血管壁弥漫性增厚

29. 男性，38 岁，面部及下肢浮肿两月余，尿液检查蛋白 ++++，肾穿刺组织光镜观察肾小球体积增大，肾小球毛细血管基底膜弥漫性增厚。此例肾炎的类型最大可能是

A. 弥漫性膜性增生性肾小球肾炎

B. 弥漫性膜性肾小球肾炎

C. 轻微病变性肾小球肾炎

D. 弥漫性新月体性肾小球肾炎

E. 弥漫性毛细血管内增生性肾小球肾炎

30. 男性，38 岁，面部及下肢浮肿两月余，尿液检查蛋白 ++++，肾穿刺组织光镜观察肾小球体积增大，肾小球毛细血管基底膜弥漫性增厚。此例患者肾脏呈

A. 大红肾

B. 固缩肾

C. 大白肾

D. 蚤咬肾

E. 以上都不是

31. 不符合弥漫性膜性肾小球肾炎的描述是

A. 起病缓慢，病程长

B. 对皮质激素治疗效果不佳

C. 晚期导致肾功能衰竭

D. 常引起肾病综合征

E. 儿童患者多见

32. 弥漫性膜性肾小球肾炎临床上常表现为

A. 肾病综合征

B. 急性肾炎综合征

C. 慢性肾炎综合征

D. 隐匿性肾炎综合征

E. 急进性肾炎综合征

33. 弥漫性膜性增生性肾小球肾炎增生的成分是

A. 球囊壁层上皮细胞

B. 球囊脏层上皮细胞

C. 系膜细胞和内皮细胞

D. 系膜细胞和基质

E. 内皮细胞

34. 不符合弥漫性膜性增生性肾小球肾炎的描述是

A. 系膜细胞和系膜基质弥漫性增生

B. 肾小球基底膜弥漫性增厚

C. 肾小球系膜区 IgG 和 C3 沉积
D. 肾小球系膜区电子致密物
E. 毛细血管壁呈车轨状

35. 不符合弥漫性膜性增生性肾小球肾炎病变特点的是
A. 系膜细胞及基质增生
B. 系膜区增宽
C. 系膜硬化
D. 系膜内无炎性细胞浸润
E. 毛细血管壁呈车轨状

36. 低补体血症性肾小球肾炎属于
A. 弥漫性新月体性肾小球肾炎
B. 慢性硬化性肾小球肾炎
C. 轻微病变性肾小球肾炎
D. 弥漫性膜性肾小球肾炎
E. 弥漫性膜性增生性肾小球肾炎

37. 符合弥漫性膜性增生性肾小球肾炎的描述是
A. 系膜区致密物沉积
B. 基底膜内致密物沉积
C. 基底膜和上皮细胞间驼峰状致密物沉积
D. 基底膜内和基底膜内、外侧致密物沉积
E. 无致密物沉积

38. 致密物沉积病属于
A. 弥漫性新月体性肾小球肾炎
B. 慢性硬化性肾小球肾炎
C. 轻微病变性肾小球肾炎
D. 弥漫性膜性肾小球肾炎
E. 弥漫性膜性增生性肾小球肾炎

39. 不符合膜性增生性肾小球肾炎的描述是
A. 多由高分子量循环免疫复合物引起
B. 多数患者预后良好
C. IgG 和 C3 沉积于系膜区和基底膜
D. 系膜细胞和基质增生
E. 可引起肾病综合征

40. 符合致密物沉积病的描述是
A. C3 颗粒沉积于基底膜内
B. 患者预后良好
C. 毛细血管基底膜正常
D. IgG 颗粒沉积于基底膜内
E. 临床表现为急性肾炎综合征

41. 弥漫性系膜增生性肾小球肾炎的主要病理特点是
A. 肾小球系膜细胞和内皮细胞增生
B. 肾小球基底膜弥漫增厚并有钉突形成
C. 肾小球系膜细胞及基质增生
D. 肾小球硬化
E. 肾小球无明显病变

42. 不符合弥漫性系膜增生性肾小球肾炎的病变是
A. 系膜细胞和基质增生
B. 毛细血管无明显改变
C. 系膜区电子致密物沉积
D. 系膜区 IgM 、IgG 和 C3 沉积
E. 系膜细胞和内皮细胞弥漫性增生

43. 符合弥漫性系膜增生性肾小球肾炎的临床表现是
A. 发病率高
B. 青少年多见
C. 可表现为急性肾炎综合征

D. 可表现为隐匿性肾炎综合征
E. 不表现为肾病综合征

44. 符合弥漫性系膜增生性肾小球肾炎的电镜下变化是
A. 系膜区致密物沉积
B. 基底膜内致密物沉积
C. 基底膜和上皮细胞间驼峰状致密物沉积
D. 基底膜内和基底膜内、外侧致密物沉积
E. 无致密物沉积

45. IgA 肾病属于
A. 毛细血管增生性肾小球肾炎
B. 抗基底膜性肾小球肾炎
C. 弥漫性膜增生性肾小球肾炎
D. 弥漫性系膜增生性肾小球肾炎
E. 轻微病变性肾小球肾炎

46. 不符合 IgA 肾病的描述是
A. 约占弥漫性系膜增生性肾小球肾炎的 50%
B. 青少年多见
C. 可引起肉眼血尿
D. 预后差
E. 可引起肾病综合征

47. 不符合弥漫性系膜增生性肾小球肾炎的描述是
A. 约占原发性肾小球肾炎的 50%
B. 可分为 IgA 肾病及非 IgA 肾病
C. 部分病例可发展为硬化性肾小球肾炎
D. 皮质激素和细胞毒药物治疗无效
E. 可引起肾病综合征

48. 符合弥漫性系膜增生性肾小球肾炎的描述是
A. 肾小球系膜区免疫复合物沉积
B. 肾小球基底膜内免疫复合物沉积
C. 肾小球基底膜和上皮细胞间驼峰状免疫复合物沉积
D. 肾小球基底膜和内皮细胞间免疫复合物沉积
E. 肾小球内无免疫复合物沉积

49. 轻微病变性肾小球肾炎的主要病理特点是
A. 肾小管上皮脂肪变性和坏死
B. 肾小球系膜细胞增生
C. 肾小球内皮细胞增生
D. 肾小球球囊壁层上皮细胞增生
E. 肾小球球囊脏层上皮细胞足突融合

50. 轻微病变性肾小球肾炎的免疫病理变化是
A. 肾小球基底膜上皮下有免疫复合物沉积
B. 肾小球基底膜内有免疫复合物沉积
C. 肾小球基底膜内皮细胞下有免疫复合物沉积
D. 肾小球系膜区有免疫复合物沉积
E. 以上都不是

51. 不符合轻微病变性肾小球肾炎的临床表现是
A. 高度水肿
B. 血尿
C. 低蛋白血症
D. 高胆固醇血症
E. 大量蛋白尿

52. 符合轻微病变性肾小球肾炎的病因发病的描述是
    A. 由循环免疫复合物引起
    B. 属抗基底膜肾小球肾炎
    C. 由植入性非肾性抗原引起
    D. 通过补体旁路激活引起
    E. 由 T 细胞介导引起
53. 与免疫复合物无关的肾小球肾炎是
    A. 弥漫性膜性肾小球肾炎
    B. 弥漫性新月体性肾小球肾炎
    C. 轻微病变性肾小球肾炎
    D. 弥漫性毛细血管内增生性肾小球肾炎
    E. 弥漫性膜性增生性肾小球肾炎
54. 多发生于儿童的肾小球肾炎是
    A. 弥漫性新月体性肾小球肾炎
    B. 弥漫性膜性肾小球肾炎
    C. 弥漫性膜性增生性肾小球肾炎
    D. 弥漫性毛细血管内增生性肾小球肾炎
    E. 脂性肾病
55. 激素治疗效果较好的肾小球肾炎是
    A. 弥漫性新月体性肾小球肾炎
    B. 轻微病变性肾小球肾炎
    C. 弥漫性膜性肾小球肾炎
    D. 弥漫性膜性增生性肾小球肾炎
    E. 弥漫性系膜增生性肾小球肾炎
56. 轻微病变性肾小球肾炎常表现为
    A. 肾病综合征
    B. 急性肾炎综合征
    C. 慢性肾炎综合征
    D. 隐匿性肾炎综合征
    E. 急进性肾炎综合征
57. 不符合局灶性节段性肾小球肾炎的描述是
    A. 可继发于感染性心内膜炎
    B. 肾小球毛细血管袢节段性坏死
    C. 可引起血尿
    D. 有复发倾向
    E. 多引起肾功能衰竭
58. 局灶性节段性肾小球肾炎常表现为
    A. 肾病综合征
    B. 急性肾炎综合征
    C. 慢性肾炎综合征
    D. 隐匿性肾炎综合征
    E. 急进性肾炎综合征
59. 局灶性节段性肾小球肾炎的病变特点为
    A. 肾小管上皮脂肪变性和坏死
    B. 肾小球系膜细胞增生
    C. 肾小球内节段性病变
    D. 肾小球球囊壁层上皮细胞增生
    E. 肾小球球囊脏层上皮细胞足突融合
60. 符合局灶性节段性肾小球肾炎结局的描述是
    A. 多数转变为慢性硬化性肾小球肾炎
    B. 多数转变为新月体性肾小球肾炎
    C. 多数可自行消退
    D. 多数严重影响肾功能
    E. 多数 10 年以内死亡
61. 符合原发性局灶性节段性肾小球肾炎病因的描述是
    A. 由循环免疫复合物引起
    B. 属抗基底膜肾小球肾炎

C. 由植入性非肾性抗原引起
D. 病因不明
E. 由 T 细胞介导引起

62. 不符合局灶性节段性肾小球肾炎病变特点的是
A. 部分肾小球受累
B. 肾小球节段性增生
C. 肾小球节段性硬化
D. IgM 和 C3 局灶性沉积
E. 基底膜增厚呈双轨状

63. 继发性局灶性节段性肾小球肾炎常表现为
A. 可继发于系统性红斑狼疮
B. 可继发于过敏性紫癜
C. 可继发于细菌性心内膜炎
D. 可继发于结节性动脉炎
E. 不继发于艾滋病

64. 不符合局灶性节段性肾小球肾炎的描述是
A. 不转变为慢性硬化性肾小球肾炎
B. 多数肾小球无病变
C. 多数可自行消退
D. 多数对肾功能影响较小
E. 多数呈隐匿性肾炎综合征

65. 硬化性肾小球肾炎的主要病理特点是
A. 肾间质纤维化
B. 肾小管萎缩
C. 部分肾小球硬化、肾小管萎缩和肾间质纤维化
D. 大部分肾小球硬化、肾小管萎缩和肾间质纤维化
E. 入球小动脉玻璃样变性

66. 不符合硬化性肾小球肾炎的是
A. 部分肾单位代偿肥大
B. 大部分肾单位硬化和萎缩
C. 颗粒性萎缩肾
D. 肾盂扩张
E. 肾间质纤维化

67. 慢性硬化性肾小球肾炎特征性的尿变化是
A. 蛋白尿
B. 少尿无尿
C. 多尿夜尿
D. 管型尿
E. 血尿

68. 很少发展为慢性硬化性肾小球肾炎的是
A. IgA 肾病
B. 膜性增生性肾小球肾炎
C. 膜性肾小球肾炎
D. 毛细血管内增生性肾小球肾炎
E. 新月体性肾小球肾炎

69. 慢性硬化性肾小球肾炎的肾脏呈
A. 大红肾
B. 颗粒性固缩肾
C. 大白肾
D. 蚤咬肾
E. 大瘢痕性固缩肾

70. 符合慢性硬化性肾小球肾炎的描述是
A. 是各型肾小球肾炎的终末期
B. 很少由毛细血管内增生性肾小球肾炎发展而来
C. 可由新月体性肾小球肾炎转化而来
D. 可由膜性增生性肾小球肾炎转化而

来

E. 患者皆有肾小球肾炎病史

71. 慢性硬化性肾小球肾炎引起肾功能衰竭的主要原因是

A. 肾小球代偿性肥大

B. 肾间质纤维化

C. 肾间质慢性炎细胞浸润

D. 大部分肾小球纤维化

E. 肾小动脉硬化

72. 不符合慢性硬化性肾小球肾炎的描述是

A. 可引起贫血

B. 可引起高血压

C. 可引起严重的蛋白尿

D. 可引起水电解质紊乱

E. 可引起氮质血症

73. 不引起尿毒症的疾病是

A. 慢性硬化性肾小球肾炎

B. 肾细胞癌

C. 肾结核

D. 高血压肾硬化

E. 肾盂肾炎

74. 引起尿毒症的病理基础是

A. 肾小球代偿性肥大

B. 肾间质纤维化

C. 肾间质慢性炎细胞浸润

D. 大部分肾单位破坏

E. 肾小动脉硬化

75. 引起尿毒症的主要代谢产物是

A. 肌酐

B. 酚类及其代谢产物

C. 肌酸

D. 尿素

E. 胍类及其代谢产物

76. 尿毒症性肺炎的病变特点是

A. 肺泡腔内透明膜形成

B. 肺泡腔内大量中性白细胞

C. 肺泡腔内大量纤维素

D. 肺泡腔内大量巨噬细胞浸润

E. 肺泡上皮细胞弥漫性增生

77. 尿毒症时，引起纤维素性胸膜炎的是

A. 肌酐

B. 酚类及其代谢产物

C. 肌酸

D. 尿素

E. 胍类及其代谢产物

78. 不符合尿毒症患者贫血、出血的原因是

A. 尿素的毒性作用

B. 酚类及其衍生物抑制骨髓造血功能

C. 胍类及其衍生物引起血小板功能障碍

D. 促红细胞生成素减少

E. 胍类及其衍生物加速红细胞破坏

79. 不符合尿毒症引起的肾性骨营养不良的描述是

A. 与低血钙有关

B. 引起甲状旁腺功能亢进

C. 引起骨质疏松

D. 1,25-二羟胆钙化醇及其类似药物治疗无效

E. 与肾排泄磷酸盐功能下降有关

80. 不符合尿毒症引起的心血管疾病的描述是

A. 引起高血压

B. 引起心脏肥大
C. 引起心肌变性坏死
D. 引起特发性心肌病
E. 引起纤维素性心外膜炎

81. 肾盂肾炎的基本病变是
A. 化脓性炎
B. 变态反应性炎
C. 增生性炎
D. 变质性炎
E. 纤维素性炎

82. 肾盂肾炎主要累及
A. 肾小球
B. 肾小管
C. 肾间质
D. 肾盂
E. 肾小管、肾间质和肾盂

83. 肾盂肾炎最常见的病原体是
A. 变形杆菌
B. 产气杆菌
C. 大肠杆菌
D. 葡萄球菌
E. 链球菌

84. 不符合肾盂肾炎的病因发病的描述是
A. 多为上行性感染
B. 多为血源性感染
C. 多为大肠杆菌感染
D. 与尿路阻塞有关
E. 多发生于女性

85. 上行性感染引起的急性肾盂肾炎最先累及
A. 肾小球
B. 肾小管
C. 集合管
D. 肾乳头
E. 肾盂粘膜

86. 血源性感染引起的急性肾盂肾炎的病变特点是
A. 肾盂粘膜充血水肿
B. 肾小管内充满脓细胞
C. 肾脏散在的多发性小脓肿
D. 肾小球纤维化
E. 肾乳头坏死

87. 急性肾盂肾炎的合并症不包括
A. 急性坏死性乳头炎
B. 肾盂积脓
C. 肾周围脓肿
D. 肾结石
E. 急性膀胱炎

88. 急性肾盂肾炎一般不引起
A. 发热、寒战、白细胞增高
B. 腰痛
C. 脓尿、菌尿、蛋白尿
D. 尿频、尿急、尿痛
E. 尿毒症

89. 慢性肾盂肾炎的基本病变属于
A. 纤维素性炎
B. 肉芽肿性炎
C. 卡他性炎
D. 慢性化脓性炎
E. 增生性炎

90. 慢性肾盂肾炎的主要病因是
A. 变态反应
B. 血循环障碍
C. 细菌感染

D. 先天性发育异常

E. 代谢障碍

91. 慢性肾盂肾炎的主要病变特点是

A. 肾小球硬化

B. 肾小球新月体形成

C. 肾小管坏死

D. 肾间质慢性化脓性炎

E. 肾间质纤维化

92. 符合慢性肾盂肾炎临床症状的是

A. 肾病综合征

B. 急性肾炎综合征

C. 蛋白尿、脓尿、菌尿

D. 慢性肾炎综合征

E. 急进性肾炎综合征

93. 慢性肾盂肾炎的肾脏呈

A. 凹陷性瘢痕伴脓肿形成

B. 颗粒性固缩肾

C. 大白肾

D. 蚤咬肾

E. 大瘢痕性固缩肾

94. 支持慢性肾盂肾炎的尿的变化是

A. 血尿

B. 蛋白尿

C. 管形尿

D. 多尿

E. 细菌尿

95. 不符合慢性肾盂肾炎的描述是

A. 常反复发作

B. 可引起肾功能衰竭

C. 可引起高血压

D. 可出现多尿和夜尿

E. 常引起慢性肾炎综合征

96. 慢性肾盂肾炎不同于慢性硬化性肾小球肾炎的病变是

A. 肾小球代偿性肥大

B. 肾内小动脉硬化

C. 肾间质纤维化

D. 肾小球囊周围纤维化

E. 肾小管扩张

97. 不符合膀胱乳头状瘤的描述是

A. 手术后易复发

B. 乳头被覆腺上皮

C. 乳头易折断、脱落

D. 易发生癌变

E. 乳头有蒂与粘膜相连

98. 膀胱癌最常见的组织学类型是

A. 腺癌

B. 鳞癌

C. 移行细胞癌

D. 未分化癌

E. 腺鳞癌

99. 女性，60 岁，送检组织为膀胱肿瘤标本，肉眼检查：肿瘤直径约 3cm，呈乳头状，表面溃疡形成，切面灰白、质硬，侵及膀胱肌层。符合此肿瘤的诊断是

A. 腺癌

B. 鳞癌

C. 移行细胞癌

D. 未分化癌

E. 平滑肌瘤

100. 膀胱癌的早期临床症状是

A. 无痛性血尿

B. 菌尿

C. 管形尿

D. 脓尿
E. 蛋白尿

101. 移行细胞癌最常见的好发部位是
A. 膀胱三角区和侧壁
B. 输尿管
C. 肾盂
D. 膀胱前壁
E. 膀胱颈部

102. 膀胱癌不引起
A. 肾盂积水
B. 尿频、尿急、尿痛
C. 血尿
D. 肾盂肾炎
E. 肾小球肾炎

103. 预后较好的膀胱癌是
A. 移行细胞癌 I 级
B. 移行细胞癌 II 级
C. 移行细胞癌 III 级
D. 鳞状细胞癌
E. 未分化癌

104. 预后最好的膀胱癌是
A. 原位癌
B. 粘膜下癌
C. 侵及膀胱浅肌层的癌
D. 侵及膀胱深肌层的癌
E. 累及膀胱周围器官的癌

105. 肾脏最常见的肿瘤是
A. 肾盂乳头状瘤
B. 血管平滑肌脂肪瘤
C. 肾母细胞瘤
D. 肾细胞癌
E. 肾腺瘤

106. 肾细胞癌来源于
A. 移行上皮
B. 鳞状上皮
C. 肾小管上皮
D. 肾球囊壁层上皮
E. 足突细胞

107. 与肾细胞癌发生关系密切的基因是
A. APC 基因
B. Rb 基因
C. WT-1 基因
D. VHL 基因
E. NF-1 基因

108. 肾细胞癌最常见的扩散途径是
A. 沿腹膜后静脉转移到肝
B. 种植转移到盆腔
C. 沿淋巴道转移到主动脉旁淋巴结
D. 沿肾静脉转移到肺
E. 沿精索静脉转移到生殖器官

109. 不符合肾细胞癌的描述是
A. 早期发生血道转移
B. 发病隐匿
C. 不引起副肿瘤综合征
D. 多见于肾脏两极
E. 可引起肾盂积水

110. 肾细胞癌最常见的组织学类型是
A. 透明细胞癌
B. 乳头状癌
C. 颗粒细胞癌
D. 未分化癌
E. 移行细胞癌

111. 肾细胞癌危险因素不包括
A. 吸烟

B. 肥胖
C. 遗传
D. 糖尿病
E. 高血压

112. 不符合肾细胞癌的描述是
A. 肿瘤间质丰富，血管少
B. 发病隐匿
C. 可引起副肿瘤综合征
D. 多见于肾脏上极
E. 肿瘤与邻近肾组织分界清楚

113. 肾母细胞瘤是
A. Wilms' tumor
B. Ewing tumor
C. Krukenberg tumor
D. Borderline tumor
E. Hodgkin diseases

114. 肾母细胞瘤起源于
A. 肾小管
B. 移行上皮
C. 原始肾组织
D. 球囊壁层上皮
E. 肾间质

115. 儿童肾脏常见的肿瘤是
A. 肾盂乳头状瘤
B. 血管平滑肌脂肪瘤
C. 肾母细胞瘤
D. 肾细胞癌
E. 肾腺瘤

116. 肾母细胞瘤最常见的临床症状是
A. 腹部肿块
B. 血尿
C. 高血压
D. 水肿
E. 蛋白尿

117. 肾母细胞瘤的肉眼特点是
A. 鱼肉状，边界清楚
B. 淡黄色，质软，边界清楚
C. 灰白，质硬，边界不清
D. 鱼肉状，边界不清
E. 淡黄色，质软，边界不清

118. 不符合肾母细胞瘤的组织学改变
A. 原始肾小管结构
B. 原始肾小球结构
C. 横纹肌细胞
D. 软骨细胞
E. 角化珠

119. 不符合肾母细胞瘤的描述是
A. 儿童患者多见
B. 起源于肾小管上皮细胞
C. 肿瘤呈鱼肉状
D. 肿瘤细胞有多方向分化潜能
E. 可经血道和淋巴道转移

120. 符合肾母细胞瘤的描述是
A. 成人患者多见
B. 肿瘤呈灰白、质硬的结节
C. 引起肾病综合征
D. 肿瘤由癌细胞团和硬化的纤维间质构成
E. 最常见的临床症状是腹部肿块

121. 尿结石不会引起
A. 肾盂肾炎
B. 输尿管粘膜溃疡
C. 血尿
D. 粘膜慢性炎

E. 肾细胞癌

122. 最常见的尿结石类型是

A. 草酸盐结石

B. 磷酸钙结石

C. 磷酸铵酶结石

D. 尿酸盐结石

E. 胱氨酸结石

123. 尿结石中主要的成分是

A. 无机盐结晶

B. 粘蛋白

C. 胆固醇

D. 胆色素

E. 虫卵

124. 尿结石对机体的主要影响是

A. 癌变

B. 纤维化

C. 血尿

D. 慢性炎

E. 阻塞和损伤

125. 不符合尿结石形成的原因是

A. 尿内盐类浓度增高

B. 蛋白尿

C. 尿潴留

D. 尿内异物

E. 尿液 pH 的改变

126. 引起尿结石的最常见的因素是

A. 高尿酸尿

B. 高钙尿

C. 高草酸尿

D. 高胱氨酸尿

E. 高嘌呤尿

127. 最常见的引起高尿钙的原因是

A. 长期卧床引起骨钙吸收过多

B. 肾近曲管钙重吸收减少

C. 甲状旁腺功能亢进

D. 经消化道钙吸收过多

E. 恶性肿瘤骨转移

128. 尿结石最常见的部位是

A. 肾实质

B. 肾盏

C. 肾盂

D. 输尿管

E. 膀胱

129. 不符合继发性肾小球肾炎的描述是

A. 肾小球病变是全身疾病的一部分

B. 其他脏器疾病累及肾小球

C. 临床表现与原发性肾小球肾炎不同

D. 肾小球病变与原发性肾小球肾炎不易区分

E. 预后取决于原发性疾病和肾脏疾病

130. 不引起继发性肾小球肾炎的疾病有

A. 糖尿病

B. Wegener 肉芽肿

C. 系膜性红斑狼疮

D. 结核病

E. 乙型肝炎

131. 预后较好的狼疮性肾小球肾炎是

A. 轻微病变性肾小球肾炎

B. 膜性肾小球肾炎

C. 膜增生性肾小球肾炎

D. 新月体性肾小球肾炎

E. 系膜增生性肾小球肾炎

132. 亚急性细菌性心内膜炎常引起

A. 轻微病变性肾小球肾炎

B. 膜性肾小球肾炎
C. 膜增生性肾小球肾炎
D. 局灶性肾小球肾炎
E. 系膜增生性肾小球肾炎

133. 过敏性紫癜性肾小球肾炎最常见的是
A. 轻微病变性肾小球肾炎
B. 膜性肾小球肾炎
C. 膜增生性肾小球肾炎
D. 新月体性肾小球肾炎
E. 系膜增生性肾小球肾炎

134. Goodpasture 综合征是
A. 循环免疫复合物性肾小球肾炎
B. 植入性非肾性抗原引起的肾小球肾炎
C. 抗基底膜抗体引起的肺出血肾炎综合征
D. IgA 肾病
E. 致密物沉积病

135. Wegener 肉芽肿主要引起
A. 轻微病变性肾小球肾炎
B. 膜性肾小球肾炎
C. 膜增生性肾小球肾炎
D. 局灶性节段性坏死性肾小球肾炎
E. 系膜增生性肾小球肾炎

136. 糖尿病主要引起
A. 肾小球硬化
B. 膜性肾小球肾炎
C. 膜增生性肾小球肾炎
D. 新月体性肾小球肾炎
E. 系膜增生性肾小球肾炎

137. 最常见的肾囊肿性疾病是
A. 婴儿型多囊肾
B. 成人型多囊肾
C. 囊性肾结构不良
D. 单纯性肾囊肿
E. 髓质海绵肾

138. 常染色体显性遗传引起的肾囊肿性疾病是
A. 婴儿型多囊肾
B. 成人型多囊肾
C. 囊性肾结构不良
D. 单纯性肾囊肿
E. 髓质海绵肾

139. 常染色体隐性遗传引起的肾囊肿性疾病是
A. 婴儿型多囊肾
B. 成人型多囊肾
C. 囊性肾结构不良
D. 单纯性肾囊肿
E. 髓质海绵肾

140. 不符合成人型多囊肾的描述是
A. 常在 50 岁左右出现肾功能衰竭
B. 多累及单侧肾脏
C. 肾脏体积增大
D. 肾脏由大小不等的囊肿构成
E. 囊肿来自肾小管和集合管

141. 不符合急性肾功能衰竭的描述是
A. 急性少尿或无尿
B. 氮质血症
C. 水电解质和酸碱平衡紊乱
D. 肾小管急性损伤是发病的中心环节
E. 蛋白尿、血尿、水肿和少尿

142. 不属于急性肾小管坏死的是
A. 挤压综合征

B. 肝肾综合征
C. Goodpasture 综合征
D. 休克肾
E. 中毒性肾病

143. 急性肾功能衰竭的主要病变是
A. 蛋白管型
B. 肾小球损伤
C. 集合管扩张
D. 肾小管坏死
E. 肾细动脉纤维素样坏死

144. 引起急性肾小管坏死的主要因素是
A. 缺血和中毒
B. 水电解质和酸碱平衡紊乱
C. 免疫复合物沉积
D. 变态反应
E. 细菌感染

145. 不引起肾病综合征的肾小球肾炎的是
A. 轻微病变性肾小球肾炎
B. 膜性肾小球肾炎
C. 膜增生性肾小球肾炎
D. 新月体性肾小球肾炎
E. 系膜增生性肾小球肾炎

146. 男性，34 岁，肾病综合征十多年。近年来出现高血压、多尿、夜尿。近来出现贫血、视力减退、心率紊乱、全身出现尿味、身体虚弱等症状。数月来，呕吐、抽搐、昏迷而死亡。肾脏的病理表现可能为
A. 蚤咬肾
B. 大白肾
C. 大红肾
D. 颗粒性萎缩肾
E. 肾梗死

147. 女性，45 岁，数年前曾出现发烧、尿频、尿急、尿疼等尿路刺激症，未彻底治疗。近年来上述症状时轻时重。符合该患者疾病的病理变化是
A. 肾入球小动脉玻璃样变
B. 肾小球新月体形成
C. 肾小管坏死
D. 尿内可见大量脓细胞，可查到大肠杆菌
E. 肾小球广泛硬化

148. 男性，10 岁，2 周前患扁桃腺炎，经治疗好转。近 2 天来出现眼睑水肿，少尿、血尿，尿蛋白（+++），血压升高（150/90mmHg）。患者的临床表现符合
A. 急性肾炎综合征
B. 慢性肾炎综合征
C. 急进性肾炎综合征
D. 隐匿性肾炎综合征
E. 肾病综合征

149. 肾病综合征不包括
A. 蛋白尿
B. 血尿
C. 严重水肿
D. 高脂血症
E. 低蛋白血症

150. 急进性肾炎综合征是指
A. 蛋白尿、脓尿、菌尿
B. 血尿、蛋白尿、少尿，快速进入肾功能衰竭
C. 无症状的蛋白尿、血尿
D. 大量蛋白尿、低蛋白血症、高度水

肿和高胆固醇血症

E. 突起性血尿，伴有水肿、蛋白尿、少尿

151. 隐匿性肾炎综合征可见于

A. 慢性硬化性肾小球肾炎

B. 轻微病变性肾小球肾炎

C. 新月体性肾小球肾炎

D. 局灶性节段性肾小球肾炎

E. 膜性肾小球肾炎

152. 小儿肾病综合征主要见于

A. 慢性硬化性肾小球肾炎

B. 轻微病变性肾小球肾炎

C. 新月体性肾小球肾炎

D. 局灶性节段性肾小球肾炎

E. 膜性肾小球肾炎

153. glomerulonephritis

A. 肾盂肾炎

B. 肾母细胞瘤

C. 肾小球肾炎

D. 肾病

E. 尿石症

154. nephroblastoma

A. 肾盂肾炎

B. 肾母细胞瘤

C. 肾小球肾炎

D. 肾病

E. 尿石症

155. pyelonephritis

A. 肾盂肾炎

B. 肾母细胞瘤

C. 肾小球肾炎

D. 肾病

E. 尿石症

156. urolithiasis

A. 肾盂肾炎

B. 肾母细胞瘤

C. 肾小球肾炎

D. 肾病

E. 尿石症

157. transitional cell carcinoma

A. 移行细胞癌

B. 鳞状细胞癌

C. 肾细胞癌

D. 肾母细胞瘤

E. 移行细胞乳头状瘤

158. uremia

A. 肾盂肾炎

B. 肾母细胞瘤

C. 肾小球肾炎

D. 尿毒症

E. 尿石症

159. crescentic glomerulonephritis

A. 膜性肾小球肾炎

B. 轻微病变性肾小球肾炎

C. 新月体性肾小球肾炎

D. 系膜增生性肾小球肾炎

E. 膜性增生性肾小球肾炎

160. mesangioproliferative glomerulonephritis

A. 膜性肾小球肾炎

B. 轻微病变性肾小球肾炎

C. 新月体性肾小球肾炎

D. 系膜增生性肾小球肾炎

E. 膜性增生性肾小球肾炎

英文题

1. What is the most common cause of glomerulonephritis in the world?
   A. Minimal change disease
   B. Membranous glomerulonephritis
   C. IgA nephropathy
   D. Acute poststreptococcal glomerulonephritis
   E. Membranoproliferative glomerulonephritis
2. What is the leading cause of end stage renal disease?
   A. Membranous glomerulonephritis
   B. Diabetic nephropathy
   C. IgA nephropathy
   D. Minimal change disease
   E. Membranoproliferative glomerulonephritis
3. A 65-year-old man recently retired after many years in a job that involved exposure to aniline dyes, including β-naphthylamine. One month ago, he had an episode of hematuria that was not accompanied by abdominal pain. On physical examination, there are no abnormal findings. Urinalysis shows 4+hematuria and no ketones, glucose, or protein. Microscopic examination of the urine shows RBCs that are too numerous to count, 5 to 10 WBCs per high-power field, and no crystals or casts. The result of a urine culture is negative. Which of the following is the most likely diagnosis?
   A. Renal cell carcinoma
   B. Hemorrhagic cystitis
   C. Urothelial carcinoma
   D. Tubercular cystitis
   E. Squamous cell carcinoma of the urethra
4. A 49-year-old man had increased swelling in the extremities for 2 months. Physical examination showed generalized edema. A 24-hour urine collection yielded 4.1 g of protein as well as albumin and globulins. Extensive testing showed no evidence of any systemic disease, such as diabetes mellitus or systemic lupus erythematosus. He did not respond to a course of corticosteroid therapy. A renal biopsy was performed, and microscopic examination showed diffuse thickening of the basement membrane. Immunofluorescence staining with antibody to the C3 component of complement was positive in a granular pattern in the glomerular capillary loops. Two years later, he experiences increasing malaise, and laboratory studies show serum creatinine level of 45 mg/dL and urea nitrogen level of 44 mg/dL. Which of the following immunologic mechanisms was most likely responsible for the glomerular changes observed in the biopsy specimen?

A. Antibodies that react with basement membrane collagen
B. Antibodies against streptococci that cross-react with the basement membrane
C. Release of cytokines by inflammatory cells
D. Cytotoxic T cells directed against renal antigens
E. Deposition of immune complexes on the basement membrane

5. One week after a mild flu like illness, a 9-year-old boy has an episode of hematuria that subsides within 2 days. One month later, he tells his parents that his urine is red again. On physical examination, there are no significant findings. Urinalysis shows 1+ proteinuria, 1+ hematuria, and no ketones, glucose, or urobilinogen. The serum urea nitrogen level is 36 mg/dL and the creatinine level is 32 mg/dL. A renal biopsy specimen shows diffuse mesangial proliferation and electron-dense deposits in the mesangium. Which of the following mechanisms is most likely to produce these findings?
A. Deposition of immune complexes containing IgA
B. Formation of antibodies against type IV collagen
C. Virus-mediated injury to the glomeruli
D. Cytokine-mediated injury to the capillaries
E. Congenital defects in the structure of glomerular basement membranes

6. A 3-year-old girl has complained of abdominal pain for the past month. On physical examination, she is febrile, and palpation of the abdomen shows a tender mass on the right. Laboratory studies show hematuria without proteinuria. An abdominal CT scan shows a 12-cm circumscribed, solid mass in the right kidney. A right nephrectomy is performed, and the gross appearance of the mass is shown above. Which of the following is the most likely diagnosis?
A. Angiomyolipoma
B. Interstitial cell tumor
C. Renal cell carcinoma
D. Wilms tumor
E. Transitional cell carcinoma

7. A 40-year-old man has experienced increasing lower extremity edema for the past 3 weeks. On physical examination, he has pitting edema up to the knees. The 24-hour urine protein level is 7g. Urinalysis shows 4+ protein but no blood or glucose. Serologic test results for ANA, anti-streptolysin O (ASO), HIV, and ANCA are all negative. A renal biopsy is performed. Light microscopy shows glomeruli which appear normal except for diffuse, uniform thickening of

the glomerular capillary basement membranes. Immunofluorescence shows granular peripheral staining of the basement membrane for IgG and C3. Given the clinical and histopathologic findings, electron microscopy of this biopsy specimen is most likely to show which of the following findings?

A. Subepithelial electron-dense deposits
B. Subendothelial electron-dense deposits with duplication of the basement membrane
C. Mesangial electron-dense deposits only
D. No deposits but marked podocyte foot process fusion
E. No abnormality

8. A 26-year-old man is involved in a motor vehicle accident and sustains acute blood loss. He is hypotensive for several hours before paramedical personnel arrive. They stabilize the bleeding and transport him to a hospital, where he receives a transfusion of 3 units of packed RBCs. Over the next week, the serum urea nitrogen level increases to 48md/dL, the serum creatinine level increases to 5mg/dL, and the urine output decreases. He undergoes hemodialysis for the next 2 weeks and then develops marked polyuria, with urine output of 2 to 3L per day. His recovery is complicated by bronchopneumonia, but renal function gradually returns to normal. Which of the following histological features best characterizes the patient's transient renal disease?


A. Glomerular crescents in Bowman's space
B. Interstitial lymphocytic infiltrates
C. Arteriolar fibrinoid necrosis
D. Nodular glomerulosclerosis
E. Rupture of the tubular basement membrane

9. A 50-year-old woman has had flank pain for the past 2 days. Physical examination shows bilateral costovertebral angle tenderness. Urinalysis shows 2+ hematuria, and no protein, glucose, or ketones. An abdominal ultrasound scan shows bilaterally enlarged kidneys that have been replaced completely by 0.5- to 0.3-cm cysts. Which of the following complication is most likely to develop in this patient?

A. Hypertension
B. Renal cell carcinoma
C. Nephrotic syndrome
D. Microangiopathic hemolytic anemia
E. Renal infarction

10. A 77-year-old man has had increasing difficulties with urination for the past 2 years. He has difficulty starting and stopping the urine stream. On physical examination, his temperature is 37.0 ℃ and blood pressure is 130/85 mmHg.

The representative gross appearance of the bladder is shown. Which of the following laboratory findings is most likely to be reported in this patient?



A. Positive ANA test result
B. Urine culture positive for Mycobacterium tuberculosis
C. Hemoglobin concentration 225g /dL
D. Schistosoma haematobium eggs in urine
E. Prostate-specific antigen level of 5ng/mL


11. A 25-year-old woman experiences a sudden onset of fever, malaise, and nausea. Her temperature is 38.2 ℃ , pulse 85/min, respirations 18/min, and blood pressure 140/90mmHg. A routine urinalysis shows 1+ proteinuria, 4+ hematuria, and no ketones or glucose. RBC casts are seen on microscopic examination of the urine. A renal biopsy is performed, and light microscopic examination shows marked glomerular hypercellularity with neutrophils in the glomerular capillary loops. Immunofluorescence microscopy shows granular deposition of IgG and C3 in glomerular capillary basement membranes. Electron microscopy shows electron-dense subepithelial “humps”. Which of the following is the most likely diagnosis?
A. Goodpasture syndrome
B. Systemic amyloidosis
C. Postinfectious glomerulonephritis
D. Diabetes mellitus
E. Membranous glomerulonephritis

12. The parents of a 6-year-old girl notice that she has become increasingly lethargic over the past 2 weeks. On examination by the physician, she has puffiness around the eyes. Laboratory findings show a serum creatinine level of 0.7 mg/dL and a urea nitrogen level of 12 md/dL. Urinalysis shows 4+ proteinuria, and no blood or glucose. The 24-hour urine protein level is 3.8g . The child’s condition improves after a course of glucocorticoid therapy. Which of the following electron microscopic findings is most likely to characterize this disease process?
A. Subepithelial electron-dense humps
B. Effacement of podocyte foot processes
C. Areas of thickened and thinned basement membrane
D. Increased mesangial matrix
E. Reduplication of glomeruhr basement membrane

13. Several days after a mild upper respiratory infection, a 29-year-old man begins passing dark red-brown urine. The dark urine persists for the next 3 days and then becomes clear and yellow, only to become red-brown again 1 month later. There are no remarkable findings on physical examination. Urinalysis shows 3+ hematuria, 1+ proteinuria, and no glucose or ketones. Microscopic examination of the urine shows RBCs and no WBCs, casts, or crystals. The 24-hour urine protein level is 200mg. A renal biopsy for evaluation of the glomeruli in this patient is most likely to show which of the following alterations?
    A. Thrombosis in the glomerular capillaries
    B. Diffuse proliferation and basement membrane thickening
    C. Mesangial lgA staining by immunofluorescence
    D. Granular staining of the basement membrane by anti-IgG antibodies
    E. Subepithelial electron-dense deposits
14. A 72-year-old woman has noticed a lowly enlarging mass on the urethra. The mass causes local pain and irritation and is now bleeding. Physical examination shows a 2.5 cm warty, ulcerated mass protruding from the external urethral meatus. There are no lesions on the labia or in the vagina. A biopsy specimen of the lesion is most likely to identity which of the following?
    A. Papilloma
    B. Leiomyoma
    C. Embryonal rhabdomyosarcoma
    D. Squamous cell carcinoma
    E. Syphilitic chancre
15. A 49-year-old man who is a long-term smoker presents with frequency and hematuria. Histologic examination of sections taken from an exophytic lesion of the urinary bladder reveals groups of atypical cells with frequent mitoses forming finger-like projections that have thin, fibrovascular cores. These groups of atypical cells do not extend into the lamina propria and muscularis. No glands or keratin production are found. Which of the following is the most likely diagnosis?
    A. Adenocarcinoma, noninvasive
    B. Inverted papilloma, noninvasive
    C. Transitional cell carcinoma in situ
    D. Squamous cell carcinoma in situ
    E. Papillary transitional cell carcinoma, noninvasive
16. Which of the following histologic changes is most likely to be seen when examining a mucosal biopsy of the urinary bladder from an individ-ual with acute cystitis due to infection with Escherichia coli?
    A. An infiltrate of lymphocytes and plasma cells

B. An infiltrate of neutrophils
C. Inflammation with eosinophils
D. Noncaseating granulomas
E. Sheets of macrophages with granular cytoplasm

17. An 8-month-old infant boy presents with an enlarging abdominal mass. Laboratory evaluation finds normal urinary levels of vanillylman-delic acid (VMA). The mass is removed surgically and microscopic sections reveal undifferentiated mesenchymal cells, immature tubules, and abortive glomerular formation. This tumor is most closely associated with abnor-malities involving which one of the following genes?
A. MET gene
B. PRCC gene
C. p16INK4a gene
D. VHLgene
E. WT-1 gene




18. A 58-year-old man presents with left-sided costovertebral pain and gross hematuria. A large mass is found in the upper pole of one of his kidneys, as seen in the picture. Which of the following histologic changes is most likely to be seen when examining microscopic sections from this mass?
A. Groups and sheets of transitional epithelial cells
B. Immature tubules and abortive glomerular formation
C. Malignant undifferentiated mesenchymal cells
D. Large cells with prominent eosinophilic cytoplasm containing numerous mitochondria
E. Uniform cells with clear cytoplasm containing glycogen and lipid

19. A 56-year-old man presents with severe headaches, nausea, and vomiting. He also relates seeing spots before his eyes and is found to have a diastolic blood pressure of 160 mmHg. Microscopic examination of a renal biopsy demonstrates hyperplastic arteriolitis. Gross examination of his kidneys is most likely to reveal which one of the following changes?
A. A finely granular appearance on the surface
B. Multiple small petechial hemorrhages on the surface
C. Diffuse, irregular cortical scars overlying dilated calyces
D. Cortical scars overlying dilated calyces in the renal poles
E. Depressed cortical areas overlying

necrotic papillae of varying stages

20. A 67-year-old woman presents with signs of slowly progressive renal failure. Physical examination reveals mild hypertension, while laboratory tests find increased serum creatinine and BUN with mild proteinuria. A few scat-tered neutrophils and rare bacteria are seen microscopically in her urine. Grossly both of her kidneys are small and irregular with dilation of the renal pelvis and clubbing of the calyces, and these changes are similar in appearance to those seen in the picture below. Histologic sections from her kidneys reveal chronic inflammation of the interstitium. Which one of the following additional microscopic findings should be present in these histologic sections?



A. The renal papillae should have numerous activated macrophages
B. The tubules should be dilated and filled with colloid casts
C. The afferent and efferent arterioles should demonstrate fibroelastic hyperplasia
D. The interstitium should have needle-shaped crystals
E. The glomeruli should have multiple fibrin microthrombi

21. Histologic sections of a kidney reveal patchy necrosis of epithelial cells of both the proximal and distal tubules with flattening of the epithelial cells, rupture of the basement membrane (tubulorrhexis), and marked interstitial edema. Acute inflammatory cells are not seen. Which of the following is the most likely diagnosis?
A. Acute pyelonephritis
B. Acute tubular necrosis
C. Diffuse cortical necrosis
D. Chronic pyelonephritis
E. Chronic glomerulonephritis

22. A 50-year-old man presents with increasing peripheral edema and dark, tea-colored urine. Laboratory examination finds decreased serum albumin, while examination of a 24h urine specimen reveals marked proteinuria. Microscopic examination of this patient's urine reveals numerous red cells along with rare red cell casts. Electron microscopic examination of a renal biopsy from this patient reveals dense, ribbon-like deposits in the lamina densa of the glomerular basement mem-

brane. Which of the following is the most likely diagnosis?

A. Acute glomerulonephritis
B. IgA nephropathy
C. Lipoid nephrosis
D. Membranoproliferative glomerulonephritis
E. Membranous glomerulopathy

23. A 3-year-old boy is being evaluated for the development of progressive peripheral edema. He is afebrile, and his blood pressure is within normal limits. Laboratory examination finds decreased serum albumin, increased serum cholesterol, and normal BUN and creatinine levels. Examination of his urine shows massive proteinuria and lipiduria, but no red blood cells are seen. A histologic section from a renal biopsy examined with a routine H&E stain is unremarkable, but electron microscopic examination finds flattening and fusion of the foot processes of the podocytes. The basement membrane is not fragmented and electron dense deposits are not found. What is the best diagnosis?

A. Diffuse proliferative glomerulonephritis (DPGN)
B. Heymann glomerulonephritis (HGN)
C. Minimal change disease (MCD)
D. Membranous glomerulopathy (MGN)
E. Membranoproliferative glomerulonephritis (MPGN)

24. Which of the following is the most likely cause of the clinical combination of generalized edema, hypoalbuminemia, and hypercholesterolemia in an adult whose urinalysis demonstrates marked proteinuria, with fatty casts and oval fat bodies?

A. Nephritic syndrome
B. Nephrotic syndrome
C. Acute renal failure
D. Renal tubular defect
E. Urinary tract infection

25. A prostatectomy specimen was examined pathologically and several sites with adenocarcinoma are mapped within the prostate. In which of the following regions of the prostate are these adenocarcinomas most likely to arise?

A. Anterior fibromuscular stroma
B. Central zone
C. Peripheral zone
D. Transitional zone
E. Periurethral zone

26. A 75-year-old man gets up several times during a football match to go to the restroom to urinate, even though he has had only one beer. This is a problem that has plagued him for severa1 years. When he visits his physician for a checkup, on physical examination he has an enlarged, nodular prostate palpated on digital rectal examination. Laboratory studies show his serum prostate specific antigen is 5

ng/mL. Which of the following pathologic findings is most likely to be present on prostate biopsy in this man?

A. Adenocarcinoma
B. Acute inflammation
C. Multiple infarctions
D. Nodular hyperplasia
E. Granulomatous inflammation

27. A 75-year-old healthy man goes to his physician for a routine check-up. On physical examination there is a firm nodule palpable in the prostate via digital rectal examination. Prostate biopsies are performed and on microscopic examination show small, crowded glands containing cells with prominent nucleoli within the nuclei. Which of the following is the most likely diagnosis?

A. Chronic prostatitis
B. Benign prostatic hyperplasia
C. Adenocarcinoma of the prostate
D. Metastatic urothelial carcinoma
E. Recent infarction of the prostate

28. A 4-year-old child has been noted by his mother to be lethargic for 2 weeks. He has periorbital edema. He is afebrile. Dipstick urinalysis reveals no glucose, ketones, or blood, but he has 4+ proteinuria. Microscopic urinalysis reveals no casts, but oval fat bodies are seen. He is treated with corticosteroid therapy and his condition improves. Which of the following renal electron micrographic findings is most characteristic of this child's disease?

A. Fusion of foot processes
B. Mesangial cell proliferation
C. Irregular thickening of the glomerular basement membranes
D. Duplication of glomerular capillary basement membranes
E. Subepithelial electron dense deposits

29. A 49-year-old woman has been hospitalized for the past 10 days for treatment of bronchopneumonia. She has developed chills and fever over the past 2 days. Her temperature is 38.8 C and she has a diffuse erythematous skin rash. Laboratory studies show a serum creatinine of 2.2 mg/dL and glucose 73 mg/dL. A peripheral blood smear reveals eosinophilia. On urinalysis she has 2+ proteinuria but no blood, glucose, or ketones. Which of the following is the most likely diagnosis?

A. Acute tubular necrosis
B. Drug-induced interstitial nephritis
C. IgA nephropathy
D. Post-streptococcal glomerulonephritis
E. Acute serum sickness

30. A 65-year-old woman is admitted with sudden onset of chest pain and is diagnosed with an acute myocardial infarction. There is difficulty maintaining adequate blood pressure and tissue perfusion for 3 days. Her serum lactate becomes el-

evated. Her serum urea nitrogen and creatinine are found to be increasing. Granular and hyaline casts are present on microscopic urinalysis. Which of the following renal lesions is most likely to be present in this situation?

A. Chronic pyelonephritis
B. Renal vein thrombosis
C. Nodular glomerulosclerosis
D. Acute tubular necrosis
E. Minimal change disease

31. A 2-year-old child has become more irritable over the past two months and does not want to eat much at meals. On physical examination the pediatrician notes an enlarged abdomen and a mass is palpable on the right. An abdominal CT scan reveals a 10 cm solid mass involving the right kidney. The resected mass has a microscopic appearance with sheets of small blue cells along with primitive tubular structures. The child receives chemotherapy and radiation therapy, and there is no recurrence. Which of the following neoplasms is this child most likely to have had?

A. Angiomyolipoma
B. Renal cell carcinoma
C. Urothelial carcinoma
D. Wilm's tumor
E. Medullary fibroma

32. A clinical study is performed with pediatric subjects who have minimal change disease. These patients are observed to have prominent periorbital edema. Laboratory test findings from serum and urine tests are analyzed. Which of the following laboratory test findings is most likely to be consistently present in these subjects?

A. Nitrite positive urinalysis specimen
B. Proteinuria >3.5 gm/24 hours
C. Hematuria with >10 RBC/hpf
D. Renal tubular epithelial cells and casts
E. Lipiduria in association with hypercholesterolemia

33. A 55-year-old man has noted passing darker urine for the past week. A urinalysis shows pH 5.5, specific gravity 1.013, 2+ blood, no protein, and no glucose. A urine cytology is performed and atypical uroepithelial cells are found. A urologist performs a cystoscopy, but no mucosal lesions are noted. He has a 60 pack year history of smoking cigarettes. Which of the following is the most likely diagnosis?

A. Adenocarcinoma of prostate
B. Acute interstitial nephritis
C. Urothelial carcinoma of renal pelvis
D. Nodular glomerulosclerosis
E. Squamous cell carcinoma of penis

34. Which of the following glomerulonephropathies is the most common cause of nephrotic syndrome in adults? It is characterized by diffuse thickening of the glomerulocapillary wall with elec-

tron dense deposits on the subepithelial side of the epithelial membrane.

A. Minimal change disease
B. Focal Segmental GN
C. Membranous GN
D. Membranoproliferative GN
E. IgA nephropathy

35. This glomerular disease is associated with low serum complement levels, the presence of granular immune deposits within the glomeruli, and subepithelial hump-like electron-dense deposits. The best diagnosis is:

A. Poststreptococcal glomerulonephritis
B. Goodpasture syndrome
C. Membranous glomerulonephropathy
D. Focal segmental glomerulosclerosis
E. IgA nephropathy

36. Nephrotic syndrome is defined by which of the following:

A. > 0.5 grams of proteinuria in 24 hours
B. > 1.5 grams of proteinuria in 24 hours
C. > 15 grams of proteinuria in 24 hours
D. > 5 grams of proteinuria in 24 hours
E. > 3.5 grams of proteinuria in 24 hours

37. A 6-year-old presents with 12 grams proteinuria/24 hour and normal renal function. He was treated with corticosteroids and had a complete remission in proteinuria. If a renal biopsy had been performed, what would likely be the sole abnormality identified?

A. Subepithelial deposits
B. Subendothelial deposits
C. Glomerular basement membrane necrosis
D. Diffuse hypercellularity
E. Diffuse loss of podocyte foot process

38. Which finding is fairly specific for chronic renal failure

A. Anemia
B. Hyaline casts
C. Hypocalcemia
D. Proteinuria
E. Broad casts see in urinalysis

39. Cellular crescents form in Bowman's space in all forms of rapidly progressive glomerulonephritis. What is the abnormality that is responsible for crescent formation?

A. Glomerular basement membrane thickening
B. Glomerular basement membrane spikes
C. Glomerular basement membrane duplication
D. Glomerular basement membrane necrosis
E. Glomerular basement membrane collapse

40. A 13-year-old girl presents with hematuria, red blood cell casts, hypertension and low complement. What form of glomerulonephritis would explain these

findings and be associated with a favorable prognosis without therapy?

A. Acute poststreptococcal glomerulonephritis
B. Membranoproliferative glomerulonephritis
C. Minimal change disease
D. Membranous glomerulonephritis
E. IgA nephropathy


41. A high fractional excretion of sodium is typically found in which of the following?

A. Hepatorenal syndrome
B. Acute glomerulonephritis
C. Acute tubular necrosis
D. Urinary tract obstruction
E. Heart failure

42. A 55-year-old male who is a long-term smoker presents with urinary frequency and hematuria. Histologic examination of sections taken from an exophytic lesion of the urinary bladder reveal groups of atypical cells with frequent mitoses forming finger-like projections that have thin fibrovascular cores. These groups of atypical cells do not extend into the lamina propria and muscularis. No glands or keratin production are found. Which of the following is the most accurate diagnosis for this bladder tumor?

A. Transitional cell carcinoma in situ
B. Inverted papilloma
C. Adenocarcinoma, noninvasive
D. Papillary transitional cell carcinoma (TCC), noninvasive
E. Squamous cell carcinoma in situ

43. The kidney shown in the photomicrograph below exhibits a tumor that originated in the upper pole. Histologic sections from this lesion would most likely reveal which of the following carcinomas?



A. Squamous cell carcinoma
B. Small cell carcinoma
C. Signet cell carcinoma
D. Clear cell carcinoma
E. Transitional cell carcinoma

44. What is the most likely cause of the combination of generalized edema, hypoalbuminemia, hypercholesterolemia, marked proteinuria, with fatty casts and oval fat bodies in the urine?

A. Acute renal failure
B. Nephrotic syndrome
C. Nephritic syndrome
D. Renal tubular defect
E. Urinary tract infection

45. A 55-year-old male is found to have a 3.0-cm “coin lesion” in the upper lobe of his left lung. The lesion is removed surgi-

cally, and histologic sections reveal sheets of malignant cells with clear cytoplasm (clear cell carcinoma). This lung lesion is most likely a metastasis from which of the following sites?

A. Breast
B. Appendix
C. Stomach
D. Pancreas
E. Kidney

46. A 69-year-old male presents with urinary frequency, nocturia, dribbling, and difficulty in starting and stopping urination. Rectal examination reveals the prostate is enlarged, firm, and rubbery. A needle biopsy reveals increased numbers of glandular elements and stromal tissue. The glands are found to have a double layer of epithelial cells. Prominent nuclei and back-to-back glands are not seen. Which of the following is the most likely diagnosis?

A. Granulomatous prostatitis
B. Chronic bacterial prostatitis
C. Acute prostatitis
D. Prostatic adenocarcinoma
E. Benign prostatic hyperplasia

47. A 49-year-old male presents with a sudden onset of fever, chills, and dysuria. During review of symptoms you discover that he has no history of recurrent urinary tract infections. Upon rectal examination, the prostate gland is very sensitive and examination is painful. Which of the following is the most likely diagnosis?

A. Chronic non-bacterial prostatitis
B. Chronic bacterial prostatitis
C. Acute prostatitis
D. Benign prostatic hyperplasia
E. Granulomatous prostatitis


48. A 5-year-old boy presents with bilateral swelling around his eyes which has been present over the past several weeks, and his urine has become cocoa-colored. The boy is afebrile and his blood pressure is slightly elevated. A urinary dipstick reveals mild proteinuria, while microscopic examination of the boy's urine reveals hematuria with red blood cell casts. Laboratory tests reveal increased ASO titers. A throat swab for streptococcus is negative. A microscopic section from the kidney reveals increased numbers of cells within the glomeruli. An electron microscopic section of the kidney reveals large electron-dense deposits in the glomeruli that are located between the basement membrane and the podocytes. The foot processes of the podocytes are otherwise unremarkable. Which one of the following renal diseases most likely produced these abnormalities in this young boy?

A. Acute post-streptococcal glomerulonephritis

B. Focal segmental glomerulonephritis
C. Minimal change disease
D. Membranous glomerulonephritis
E. Focal segmental glomerulosclerosis

49. A 37-year-old female recovering from hepatitis B develops hematuria, proteinuria, and red cell casts in the urine. Which of the following would best describe the changes within the kidney in this patient?
A. Diffuse thickening of the glomerular basement membrane by subepithelial immune deposits
B. IgG linear fluorescence along the glomerular basement membrane
C. Nodular hyaline glomerulosclerosis
D. Plasma cell interstitial nephritis
E. Granular deposits of antibodies in the glomerular basement membrane

50. The gross appearance of the kidney shown below is most compatible with which of the following conditions?



A. Chronic glomerulonephritis
B. Acute glomerulonephritis
C. Chronic pyelonephritis
D. Acute pyelonephritis
E. Cystic renal dysplasia

51. Histologic sections of a kidney reveal patchy necrosis of epithelial cells of both the proximal and distal tubules with flattening of the epithelial cells, rupture of the basement membrane (tubulorrhexis), and marked interstitial edema. Acute inflammatory cells are not seen. Which of the following is the most likely diagnosis?
A. Acute pyelonephritis
B. Acute tubular necrosis
C. Chronic pyelonephritis
D. Chronic glomerulonephritis
E. Diffuse cortical necrosis

52. An 8-year-old girl develops a severe sore throat, which resolves after a few days. 10 days later, her mother notices dark stains on the child's underwear, and takes her to the pediatrician. A urine sample is smoky in color, and red cell casts are noted in the urinary sediment. If a renal biopsy were obtained from this child it would probably show
A. Linear deposition of IgG
B. Fusion of podocyte foot processes
C. Dense deposits
D. Mesangial IgA deposits
E. Subepithelial humps

53. A 2-year-old boy has an abdominal mass

and undergoes diagnostic biopsy. The tumor reveals a variety of cellular patterns including dense immature islands of epithelial cells, ribbons of spindled fibroblast-like stromal cells, and poorly formed tubular structures. This triphasic histology is most suggestive of which of the following childhood neoplasms?

A. Embryonal rhabdomyosarcoma
B. Wilms' tumor
C. Neuroblastoma
D. Hodgkin's disease
E. Ewing's sarcoma

54. Renal biopsy demonstrates concentric, laminated thickening of arteriolar walls due to proliferation of smooth muscle cells. This process is best described by which of the following terms?

A. Polyarteritis nodosa
B. Mönckeberg's arteriosclerosis
C. Hyperplastic arteriolosclerosis
D. Hyaline arteriolosclerosis
E. Atherosclerosis

55. A 3 year-old child develops a large erythematous rash around the site of a mosquito bite. One month later, she is taken to a pediatrician because she has a puffy face and swollen ankles. The scanty urine sample has a reddish-brown hue, and contains both red blood cells and protein. Which of the following distinctive features would be most likely to be seen on renal biopsy?

A. Linear IgG deposits
B. Subepithelial electron dense humps
C. Fusion of podocyte foot processes
D. Onion-skinning of renal arterioles
E. IgA in the mesangium

56. A 60-year-old man develops peripheral edema over a period of weeks. Urinalysis reveals hematuria and proteinuria; examination of the urinary sediment reveals red cell casts. Radiologic and ultrasound studies fail to demonstrate an obstructive lesion. Renal biopsy shows many glomerular crescents. This presentation is most suggestive of which of the following conditions?

A. Anti-glomerular basement membrane disease
B. Hypertensive nephropathy
C. Diabetic nephropathy
D. Minimal change disease
E. Lupus nephritis

57. A 3-year-old girl is brought to the clinic by her parents who are very concerned about her. Physical examination is remarkable for extremity edema. Urinalysis shows significant proteinuria, but no red cells or casts. Electron microscopy performed on a kidney biopsy would be most likely to show which of the following findings?

A. Subepithelial spikes
B. Subendothelial deposits
C. Mesangial deposits

D. Fusion of epithelial foot processes
E. Dense deposits

58. A pediatrician discovers a large mass in the abdomen of a 2-year-old child. Ultrasound examination demonstrates that the mass appears to arise from the right kidney. Which of the following tumors is most likely present?
A. Renal cell carcinoma
B. Oncocytoma
C. Nephroblastoma
D. Hemangioma
E. Cortical adenoma

59. A 60-year-old man develops low back pain, and radiologic investigation reveals multiple osteoblastic lesions of the lumbar vertebrae. A hard, irregular nodule on the posterior surface of the prostate is found on rectal examination. Plasma levels of prostatic specific antigen (PSA) are markedly increased. Which of the following is the most likely diagnosis?
A. Prostatic hyperplasia
B. Multiple myeloma
C. Prostatic intraepithelial neoplasia (PIN)
D. Chronic prostatitis
E. Prostatic adenocarcinoma

60. A concerned couple bring their 1.5-year-old boy with gonadal dysgenesis to a clinic for a follow-up visit. The physician notices a large abdominal mass during his physical examination. Which of the following disorders does the patient most likely have?
A. Renal hamartoma
B. Renal cell carcinoma
C. Squamous cell carcinoma of the bladder
D. Wilms' tumor
E. Transitional cell carcinoma of the bladder

61. A 3-year-old child has a large abdominal mass. CT studies show that the mass arises in the kidney. Which of the following is most likely a distinctive feature of this lesion?
A. Hamartomatous blood vessels
B. Embryonic glomerular structures
C. Eosinophilic cells packed with mitochondria
D. Abundant clear cells
E. Necrotic bacterial forms in macrophages

62. A 5-year-old boy has developed a "puffy face." On examination, the child has marked periorbital edema, and pitting edema of both the hands and feet. His cardiac examination is unremarkable and he has no splenomegaly or signs of liver disease. Laboratory values are notable for decreased serum albumin and increased total cholesterol. A urinalysis reveals 4+ proteinuria but no red blood cells, white

blood cells, or casts. The most likely cause of this child's disease is

A. Wegener's granulomatosis
B. Systemic lupus erythematosus
C. Poststreptococcal glomerulonephritis
D. Minimal change nephropathy
E. Goodpasture's syndrome

63. A 60-year-old man develops increased frequency of urination, nocturia, dysuria, and difficulty starting and stopping the flow of urine. Serum prostate-specific antigen (PSA) is within normal limits, and prostate biopsy demonstrates benign tissue. In which of the following sites is this patient's lesion most likely located?

A. Periurethral prostate
B. Penile urethra
C. Bladder neck
D. Posterior prostate
E. Ureters

64. A pathologist examining electron micrographs of glomeruli sees prominent deposits between the podocytes and the basement membrane of the glomerular capillaries. These findings are most likely related to prior infection with which of the following bacteria?

A. Neisseria
B. Pseudomonas
C. Escherichia
D. Klebsiella
E. Streptococcus

65. A 58-year-old woman is noted to have a ruddy complexion. Her hematocrit is 52%. Her lungs are clear and she does not smoke. Serum erythropoietin levels are elevated. Cancer of which of the following organs is most likely the cause of her increased hematocrit?

A. Breast
B. Kidney
C. Colon
D. Thyroid
E. Stomach

66. The main histopathologic feature of membranous glomerulonephritis is:

A. Glomerular mesangial cell and endothelial proliferation
B. Diffuse thickening of the GBM with spikes
C. Glomerular mesangial cell proliferation with GBM thickening
D. Glomerular focal sclerosis
E. Loss of the podocyte foot processes

67. Which of the following is the characteristic feature of diffuse endocapillary proliferative glomerulonephritis by electron microscopy?

A. Electron-dense subendothelial deposits
B. Electron-dense subepithelial hump deposits
C. GBM damaged seriously
D. Diffuse fusion and loss of the

podocyte foot processes

E. Diffuse thickening of the GBM with spikes

68. The glomerulonephritis which is curable in the majority of cases is:

A. Acute diffuse endocapillary proliferative glomerulonephritis

B. Membranoproliferative glomerulonephritis

C. Crescentic glomerulonephritis

D. Membranous glomerulonephritis

E. Chronic pyelonephritis

69. Which of the following is NOT true concerning glomerulonephritis?

A. Is caused directly by bacterial infection

B. Is caused by immune complexes in situ

C. Is caused by implanted antigen

D. Is caused by intrinsic antigen

E. Is caused directly by fungus infection

70. Which of the following is NOT associated with diffuse endocapillary proliferative glomerulonephritis?

A. Diffuse involvement of bilateral kidneys

B. Electron-dense subepithelial hump deposits

C. Poststreptococcal infection

D. Very poor prognosis

E. TypeIII hypersensitivity reactions

71. Which of the following is characteristic of mesangial proliferative glomerulonephritis?

A. Electron-dense deposits in the mesangium and subendothelium

B. Electron-dense deposits in the mesangium

C. Electron-dense subepithelial humps and intramembranous deposits

D. Electron-dense subepithelial deposits and spikelike protrusions of GBM matrix

E. Electron-dense deposits in the subendothelium

72. Which of the following is characteristic of membranoproliferative glomerulonephritis?

A. Electron-dense deposits in the mesangium and subendothelium

B. Electron-dense deposits in the mesangium

C. Electron-dense subepithelial humps and intramembranous deposits

D. Electron-dense subepithelial deposits and spikelike protrusions of GBM matrix

E. Electron-dense deposits in the subendothelium

73. Which of the following is characteristic of diffuse endocapillary proliferative glomerulonephritis?

A. Electron-dense deposits in the mesangium and subendothelium

B. Electron-dense deposits in the mesangium
C. Electron-dense subepithelial humps and intramembranous deposits
D. Electron-dense subepithelial deposits and spikelike protrusions of GBM matrix
E. Electron-dense deposits in the subendothelium

74. Which of the following is characteristic of membranous glomerulonephritis?
A. Electron-dense deposits in the mesangium and subendothelium
B. Electron-dense deposits in the mesangium
C. Electron-dense subepithelial humps and intramembranous deposits
D. Electron-dense subepithelial deposits and spikelike protrusions of GBM matrix
E. Mesangial deposition of IgA

75. Glomerulonephritis is:
A. Type II hypersensitivity reaction
B. Type II and typeIIIhypersensitivity reaction
C. Type IV hypersensitivity reaction
D. Type III hypersensitivity reaction
E. Type I hypersensitivity reactions

76. The most common cause of acute renal failure is:
A. Diffuse crescentic glomerulonephritis
B. Diffuse mesangial proliferative glomerulonephritis
C. Membranous glomerulonephritis
D. Focal glomerulonephritis
E. Minimal change disease

77. The cells in crescents of crescentic glomerulonephritis include:
A. Parietal epithelial cells of the renal glomerulus
B. Parietal epithelial cells of the renal glomerulus and monocytes
C. Visceral epithelial cells of the renal glomerulus and monocytes
D. Mesangial cells of the glomerulus
E. Monocytes

78. Which of the following is a key feature of membranous nephropathy by light microscopy?
A. Diffuse thickening of the GBM
B. Endothelial proliferation
C. Thickening of the GBM and proliferation of mesangial cells and matrix
D. Diffuse fusion and loss of the podocyte foot processes
E. Podocytes proliferation

79. The proliferative element(s) in diffuse membranoproliferative glomerulonephritis is/are:
A. Visceral epithelial cells of the renal glomerulus
B. Parietal epithelial cells of the renal glomeruli
C. Mesangial cells and endothelium

D. Mesangial cells and matrix
E. Podocytes

80. IgA belongs to which of the following?
A. Endocapillary proliferative glomerulonephritis
B. Secondary glomerulonephritis
C. Diffuse mesangial proliferative glomerulonephritis
D. Diffuse membranoproliferative glomerulonephritis
E. Primary glomerulonephritis

81. Which of the following is most characteristic of IgA nephropathy?
A. Glomerular basement membrane(GBM) deposition of C3
B. Mesangial deposition of IgG and C3
C. Mesangial deposition of IgA
D. GBM and mesangial deposition of IgG and C3
E. Mesangial deposition of IgM

82. Which of the following symptoms is most characteristic of focal segmental glomerulosclerosis?
A. Acute nephritic syndrome
B. Nephrotic syndrome
C. Chronic nephritic syndrome
D. Asymptomatic nephritic syndrome
E. Rapidly progressive nephritic syndrome

83. Which of the following types of glomerulonephritis is NOT associated with immune complexes?
A. Crescentic glomerulonephritis
B. Membranous glomerulopathy
C. Minimal change disease
D. Endocapillary proliferation glomerulonephritis
E. IgA nephropathy

84. Which of the following is NOT generally a feature of mesangial proliferation glomerulonephritis?
A. Glomerular focal sclerosis
B. Increased mesangial matrix
C. Immunofluorescence positive deposition of IgG and C3 in mesangial regions
D. Deposition of electron-dense material in mesangial regions
E. Focal fusion of the foot processes of the podocytes

85. The basic lesion of acute pyelonephritis is:
A. Suppurative inflammation
B. Fibrinous inflammation
C. Hemorrhagic inflammation
D. Serous inflammation
E. Granulomatous inflammation

86. pyelonephritis primarily involves:
A. Renal tubules, renal interstitium and renal pelvis
B. Renal tubules
C. Renal pelvis
D. Glomeruli
E. Renal interstitium

87. Acute pyelonephritis caused by ascending

infection first involves:

A. Mucosal surfaces of the renal pelvis
B. Renal tubules
C. Collecting ducts
D. Renal papillae
E. Renal glomeruli

88. The most common type of kidney tumor found in children is:

A. Angiomyolipoma
B. Papilloma of renal pelvis
C. Nephroblastoma
D. Renal cell carcinoma
E. Transitional cell carcinoma

89. The most common histopathologic type of bladder carcinoma is:

A. Squamous cell carcinoma
B. Adenocarcinoma
C. Transitional cell carcinoma
D. Undifferentiated carcinoma
E. Small cell carcinoma

90. Nephroblastoma is derived from:

A. Transitional cells
B. Renal tubule
C. Primitive renal tissue
D. Parietal cells of renal glomeruli
E. Squamous epithelium

91. Nephroblastoma is:

A. Wilm's tumor
B. Ewing tumor
C. Borderline tumor
D. Krukenberg tumor
E. PNET

92. Renal cell carcinoma is derived from:

A. Renal tubular epithelium
B. Squamous epithelium
C. Renal collecting duct epithelium
D. Parietal cells of renal glomeruli
E. Transitional epithelium

93. The most common histopathologic type of renal cell carcinoma is:

A. Papillary renal cell carcinoma
B. Clear cell renal cell carcinoma
C. Chromophobe renal cell carcinoma
D. Collecting duct carcinoma
E. Undifferential carcinoma

94. The most common type of kidney tumor is:

A. Angiomyolipoma
B. Papilloma of renal pelvis
C. Nephroblastoma
D. Renal cell carcinoma
E. Transitional cell carcinoma

95. Which of the following genes is associated with renal cell carcinoma?

A. RB gene
B. APC gene
C. WT-1 gene
D. VHL gene
E. BRCA gene

96. The most common location of transitional cell carcinoma is:

A. Ureter
B. Anterior wall of bladder
C. Renal pelvis
D. Ttrigone and lateral wall of bladder
E. Dome of bladder

答案

中文题

1.E	2.E	3.C	4.A	5.E	6.C	7.C	8.A	9.A	10.C
11.C	12.E	13.B	14.E	15.C	16.B	17.B	18.E	19.E	20.A
21.E	22.A	23.C	24.D	25.B	26.B	27.D	28.E	29.B	30.C
31.E	32.A	33.D	34.B	35.D	36.E	37.D	38.E	39.B	40.A
41.C	42.E	43.E	44.A	45.D	46.D	47.D	48.A	49.E	50.E
51.B	52.E	53.C	54.E	55.B	56.A	57.E	58.D	59.C	60.C
61.D	62.E	63.E	64.A	65.D	66.D	67.C	68.D	69.B	70.E
71.D	72.C	73.B	74.D	75.D	76.C	77.D	78.A	79.D	80.D
81.A	82.E	83.C	84.B	85.E	86.C	87.D	88.E	89.D	90.C
91.D	92.C	93.A	94.E	95.E	96.D	97.B	98.C	99.C	100.A
101.A	102.E	103.A	104.A	105.D	106.C	107.D	108.D	109.C	110.A
111.D	112.A	113.A	114.C	115.C	116.A	117.A	118.E	119.B	120.E
121.E	122.A	123.A	124.E	125.B	126.B	127.D	128.C	129.C	130.D
131.A	132.D	133.E	134.C	135.D	136.A	137.D	138.B	139.A	140.B
141.E	142.C	143.D	144.A	145.D	146.D	147.D	148.A	149.B	150.B
151.D	152.B	153.C	154.B	155.A	156.E	157.A	158.D	159.C	160.D

英文题

1.C	2.B	3.C	4.E	5.A	6.D	7.A	8.E	9.A	10.E
11.C	12.B	13.C	14.D	15.E	16.B	17.E	18.C	19.B	20.B
21.B	22.D	23.C	24.B	25.C	26.D	27.C	28.A	29.D	30.D
31.D	32.B	33.C	34.C	35.A	36.E	37.E	38.E	39.D	40.A
41.C	42.D	43.D	44.B	45.D	46.E	47.C	48.A	49.E	50.C
51.B	52.E	53.B	54.C	55.B	56.A	57.D	58.C	59.E	60.D
61.B	62.D	63.A	64.D	65.B	66.B	67.B	68.A	69.A	70.D
71.B	72.A	73.C	74.D	75.D	76.A	77.B	78.A	79.D	80.B
81.C	82.B	83.C	84.A	85.A	86.E	87.A	88.C	89.C	90.C
91.A	92.A	93.B	94.D	95.D	96.D				

9 生殖系统疾病

中文题

1. 女性生殖系统结核最常见于
   A. 子宫内膜
   B. 输卵管
   C. 卵巢
   D. 子宫颈
   E. 外阴
2. 不符合女性生殖系统感染的描述是
   A. 可引起下生殖道炎症
   B. 可引起盆腔炎
   C. 可引起败血症
   D. 可引起不孕症
   E. 不导致胎儿宫内感染
3. 不属于性传播的女性生殖系统炎症性疾病是
   A. 淋病
   B. 滴虫性阴道炎
   C. 结核性盆腔炎
   D. 单纯疱疹
   E. 衣原体性盆腔炎
4. 不符合盆腔炎症性疾病的描述是
   A. 可引起下腹痛和发热
   B. 多由肠道杆菌、淋球菌和衣原体引起
   C. 可引起急性化脓性输卵管炎
   D. 可引起输卵管卵巢囊肿
   E. 流产和产后盆腔炎多为单一细菌感染
5. 属于性传播性疾病的阴道炎是
   A. 念珠菌阴道炎
   B. 细菌性阴道炎
   C. 老年性阴道炎
   D. 滴虫性阴道炎
   E. 幼女性阴道炎
6. 不符合单纯疱疹引起的女性下生殖道感染的描述是
   A. 多由 HSV-1 病毒引起
   B. 可累及外阴、阴道和子宫颈
   C. 可引起粘膜疱疹和溃疡形成
   D. 多于 1～3 周内愈合
   E. 分娩时可引起胎儿感染
7. 不符合霉菌性阴道炎的描述是
   A. 多为性传播引起
   B. 糖尿病患者易发病
   C. 孕妇易发病
   D. 阴道粘膜可见白斑
   E. 病变处可见孢子和菌丝
8. 急性盆腔炎多由
   A. 血道感染引起
   B. 腹腔脏器感染直接蔓延引起
   C. 慢性盆腔炎急性发作引起
   D. 产后、流产后、宫腔内手术后感染引起
   E. 性传播引起
9. 尖锐湿疣的病因是
   A. HPV 感染
   B. HSV 感染
   C. CMV 感染
   D. 衣原体感染

E. 细菌感染

10. 外阴尖锐湿疣属于
A. 炎症病变
B. 癌前病变
C. 原位癌
D. 早期侵润性癌
E. 良性肿瘤

11. 不符合外阴尖锐湿疣的描述是
A. 好发于外阴、肛周、阴道粘膜
B. 不累及子宫颈粘膜
C. 呈淡红色、乳头状、斑块状病变
D. 鳞状上皮乳头状增生伴角化过度
E. 棘细胞层增生并形成空泡状细胞

12. 外阴尖锐湿疣较具特征的病变是
A. 棘细胞层增生
B. 空泡状细胞
C. 角化过度
D. 乳头状增生
E. 基底细胞增生

13. 最常见的女性外阴恶性肿瘤是
A. 黑色素瘤
B. 基底细胞癌
C. Paget 病
D. 鳞状细胞癌
E. 前庭大腺癌

14. 不符合女性外阴癌的描述是
A. 与 HPV16、18 感染有关
B. 多经鳞状上皮非典型增生演化而来
C. 鳞状上皮非典型增生呈多灶性
D. 与萎缩硬化性苔癣有关
E. 多经血道转移

15. 不符合女性外阴癌的描述是
A. 多发生于老年女性
B. 可呈白斑状
C. 可呈菜花状
D. 多为鳞状细胞癌
E. 多为腺癌

16. 决定女性外阴癌的预后的主要因素是
A. 肿瘤的体积
B. 肿瘤的分化
C. 肿瘤的浸润深度
D. 肿瘤的淋巴结转移
E. 肿瘤的分期

17. 不符合有癌变倾向的外阴白色病变的描述是
A. 硬化性苔癣
B. 白癜风
C. 鳞状上皮增生
D. 鳞状上皮增生伴上皮内肿瘤
E. 增生性营养不良

18. 可以引起外阴白色病变的恶性肿瘤是
A. 硬化性苔癣
B. 白癜风
C. 鳞状上皮增生
D. 鳞状上皮增生伴上皮内肿瘤
E. Paget 病

19. 不符合外阴硬化性苔癣的描述是
A. 外阴部粘膜呈白色
B. 属癌前病变
C. 表皮变薄
D. 真皮浅层水肿
E. 真皮深层慢性炎细胞带状浸润

20. 不符合外阴鳞状上皮增生的描述是
A. 表皮增厚

B. 属癌前病变
C. 棘细胞增生
D. 真皮慢性炎细胞浸润
E. 角化过度

21. 外阴上皮内肿瘤的特点是
A. 表皮增厚
B. 棘细胞增生
C. 鳞状上皮细胞大小形态不一、核深染
D. 真皮慢性炎细胞浸润
E. 角化过度

22. 不符合外阴上皮内肿瘤的描述是
A. 常为多中心发生
B. 常合并 HPV16、18 感染
C. 鳞状上皮细胞大小形态不一、核深染
D. 可进展为鳞状细胞癌
E. 青年妇女多见

23. 外阴上皮内肿瘤常合并感染的病毒是
A. HPV
B. HSV
C. CMV
D. HIV
E. HBV

24. 符合外阴上皮内肿瘤的描述是
A. 多由硬化性苔癣发展而来
B. 常合并 CMV 感染
C. 鳞状上皮萎缩
D. 可进展为鳞状细胞癌
E. 青年妇女多见

25. 子宫颈癌开始最常发生于
A. 子宫颈外口
B. 子宫颈内口
C. 子宫颈前唇
D. 子宫颈后唇
E. 子宫颈管

26. 子宫颈癌常见的组织学类型是
A. 鳞状细胞癌
B. 原位癌
C. 早期浸润癌
D. 高分化腺癌
E. 透明细胞癌

27. 不符合子宫颈癌的描述是
A. 鳞状上皮不典型增生是癌前病变
B. 原位癌是可治愈的癌
C. 原位癌累及腺体是早期浸润癌
D. 高分化腺癌可引起粘液分泌增多
E. 淋巴道转移多见

28. 不符合子宫颈癌的描述是
A. 与慢性细菌感染有关
B. 与 HPV 感染有关
C. 与多胎生产有关
D. 与早婚有关
E. 与性生活紊乱有关

29. 女性，55 岁，停经 4 年后，近 2 月来阴道不规则少量出血。活体病理诊断为鳞状细胞癌。子宫颈癌根治标本病理诊断为早期浸润癌。符合该患者疾病的描述是
A. 癌细胞占据上皮全层
B. 癌细胞累及腺体
C. 癌细胞侵及子宫颈 2mm 的组织
D. 癌细胞侵及子宫颈深层
E. 癌细胞侵及阴道上段

30. 子宫颈早期浸润癌是指
    A. 癌浸润深度不超过 1mm
    B. 癌浸润深度不超过 2mm
    C. 癌浸润深度不超过 5mm
    D. 癌浸润深度不超过 8mm
    E. 癌浸润深度为 8mm～10mm
31. 子宫颈原位癌累及腺体是指
    A. 子宫颈腺上皮癌变
    B. 原位癌阻塞腺体开口
    C. 子宫颈腺体被鳞状上皮取代
    D. 子宫颈鳞癌细胞突破基底膜侵及腺体
    E. 子宫颈原位癌取代腺体上皮细胞，腺体基底膜完整
32. 影响子宫颈癌预后的主要因素是
    A. 肿瘤的肉眼类型
    B. 肿瘤角化珠的多少
    C. 肿瘤的腺体排列
    D. 肿瘤的临床病理分期
    E. 肿瘤细胞的分化程度
33. 不符合子宫颈癌扩散途径的描述是
    A. 以淋巴道转移为主
    B. 可直接侵犯直肠
    C. 晚期可发生血道转移
    D. 可直接侵犯膀胱
    E. 首先转移到髂总淋巴结
34. 符合子宫颈鳞状上皮异型增生的描述是
    A. 鳞状上皮细胞大小、形态不一，核深染
    B. 鳞状上皮细胞变成癌细胞，但基底膜完整
    C. 鳞状上皮增厚，钉突延长
    D. 鳞状上皮细胞不典型增生 I 级最易癌变
    E. 子宫颈柱状上皮被鳞状上皮代替
35. 子宫颈上皮内新生物是指
    A. 鳞状上皮化生
    B. 鳞状上皮不典型增生
    C. 原位癌
    D. 鳞状上皮增生
    E. 腺上皮增生
36. 子宫颈癌最常见的肉眼类型是
    A. 溃疡型
    B. 外生型
    C. 糜烂型
    D. 内生浸润型
    E. 颈管型
37. 子宫颈癌通过淋巴道首先转移到
    A. 髂总淋巴结
    B. 骶前淋巴结
    C. 子宫颈旁、闭孔、髂内、髂外淋巴结
    D. 深腹股沟淋巴结
    E. 锁骨上淋巴结
38. 与子宫颈癌发生关系密切的病毒是
    A. EBV
    B. HPV
    C. HBV
    D. CMV
    E. HSV
39. 不符合慢性子宫颈炎的描述是
    A. 孕龄妇女的常见病
    B. 子宫颈粘膜充血糜烂

C. 子宫颈慢性炎细胞浸润
D. 子宫颈息肉形成
E. 子宫颈上皮内肿瘤

40. 子宫颈真性糜烂是指
A. 粘膜上皮坏死脱落
B. 柱状上皮取代鳞状上皮
C. 重度不典型增生
D. 囊肿形成
E. 鳞状上皮化生

41. 子宫腺肌病是指子宫内膜异位于
A. 子宫肌层
B. 输卵管
C. 卵巢
D. 腹壁
E. 盆腔组织

42. 子宫外子宫内膜异位症最常见于
A. 子宫圆韧带
B. 卵巢
C. 腹壁
D. 阴道壁
E. 外阴

43. 卵巢巧克力囊肿是指
A. 输卵管卵巢囊肿
B. 卵巢子宫内膜异位囊肿
C. 卵巢粘液性囊腺瘤
D. 卵巢囊性畸胎瘤
E. 卵巢黄体血肿

44. 不符合子宫内膜增生症的描述是
A. 多发生于更年期和青春期
B. 表现为月经不规则、经期延长、月经量过多
C. 与卵巢雌激素分泌过多有关
D. 阴道壁多由卵巢滤泡不排卵引起
E. 子宫内膜单纯增生易发生癌变

45. 不符合子宫内膜增生症的病理变化是
A. 子宫内膜增厚
B. 子宫内膜腺体增生
C. 子宫内膜腺体囊性扩张
D. 子宫内膜腺上皮异型增生
E. 子宫内膜腺体背靠背、细胞异型性显著、核分裂多见

46. 功能性子宫出血是指
A. 绝经后子宫出血
B. 阴道接触出血
C. 炎症性疾病引起的子宫出血
D. 神经内分泌失常引起的子宫出血
E. 子宫肿瘤引起的子宫出血

47. 子宫内膜腺瘤样增生的特点是
A. 腺体和间质增生
B. 增生的腺体呈囊性扩张
C. 增生的腺体密集排列
D. 增生的腺体密集排列，伴腺上皮异型性
E. 腺体密集伴腺上皮异型性显著，间质浸润

48. 子宫内膜不典型增生的特点是
A. 腺体和间质增生
B. 增生的腺体呈囊性扩张
C. 增生的腺体密集排列
D. 增生的腺体密集排列，伴腺上皮异型性
E. 腺体密集伴腺上皮异型性显著，间质浸润

49. 与雌激素水平过高有密切关系的

肿瘤为

A. 子宫颈鳞癌
B. 子宫颈腺癌
C. 绒毛膜癌
D. 子宫内膜癌
E. 卵巢囊腺癌

50. 不符合高分化子宫内膜腺癌的描述是
A. 腺体密集、大小形态不规则形
B. 核分裂多见
C. 腺上皮呈复层、排列紊乱
D. 有间质浸润
E. 腺体间为子宫内膜间质

51. 诊断子宫平滑肌肉瘤的主要依据是
A. 肿瘤细胞密集
B. 可见奇异核瘤细胞
C. 肿瘤粘液变性
D. 肿瘤组织坏死、核分裂象多见（每10个高倍视野10个核分裂）
E. 肿瘤组织发生囊性变

52. 不符合子宫内膜癌的描述是
A. 与雌激素长期持续作用有关
B. 生长较缓慢
C. 主要是淋巴道转移
D. 多发生于青年妇女
E. 常引起阴道出血

53. 子宫体最常见的肿瘤是
A. 平滑肌瘤
B. 平滑肌肉瘤
C. 子宫内膜腺癌
D. 子宫内膜间质肿瘤
E. 鳞癌

54. 不符合子宫内膜腺癌的描述是
A. 肿瘤生长迅速
B. 可沿子宫内膜蔓延
C. 可侵犯子宫壁
D. 可沿淋巴道转移
E. 可沿血道转移

55. 不符合子宫内膜腺癌的描述是
A. 可呈结节状
B. 可呈息肉状
C. 可呈弥漫型
D. 多为低分化腺癌
E. 可为腺鳞癌

56. 最常见的子宫肉瘤为
A. 子宫内膜间质肉瘤
B. 平滑肌肉瘤
C. 葡萄状肉瘤
D. 癌肉瘤
E. 纤维肉瘤

57. 不符合葡萄胎的病理变化是
A. 胎盘绒毛高度水肿
B. 绒毛间质血管消失
C. 滋养层细胞增生
D. 子宫增大超出正常妊娠月份
E. 绒毛常侵犯子宫壁深肌层

58. 恶性葡萄胎的特征性病理变化是
A. 胎盘绒毛水肿
B. 滋养层细胞增生，并有异型性
C. 绒毛消失
D. 绒毛侵犯子宫壁深肌层
E. 间质血管消失

59. 恶性葡萄胎与绒毛膜癌的主要区别是
A. 浸润子宫深肌层
B. 细胞明显增生和具异型性

C. 血道转移
D. 有绒毛结构
E. 出血坏死

60. 不符合葡萄胎临床表现的是
A. HCG 增高
B. 子宫显著增大
C. 无胎心和胎动
D. 双卵巢黄素囊肿形成
E. 无恶变倾向

61. 符合葡萄胎病理变化的是
A. 浸润子宫深肌层
B. 由分化不良的两种滋养层细胞构成
C. 血道转移
D. 有绒毛结构
E. 出血坏死

62. 符合葡萄胎的病因的描述是
A. 与病毒感染有关
B. 与雌激素水平过高有关
C. 与染色体异常有关
D. 与胚胎早期死亡有关
E. 与营养缺乏有关

63. 不符合完全型葡萄胎的描述是
A. 染色体核型为二倍体
B. 染色体核型多为 46XX
C. 两组染色体均来自父方
D. 可由单精子与空卵受精引起
E. 可由双精子与正常卵子受精引起

64. 不符合部分性葡萄胎的描述是
A. 染色体核型为三倍体
B. 染色体核型多为 69XXY 或 69XXX
C. 三组染色体均来自父方
D. 也可由正常的单倍体卵子与二倍体精子受精引起
E. 可由双精子与正常卵子受精引起

65. 绒毛膜癌最常转移到
A. 阴道
B. 肝
C. 肺
D. 肠
E. 脑

66. 绒毛膜癌可以产生
A. 雌激素
B. 雄激素
C. 绒毛膜促性腺激素
D. 催乳素
E. 生长激素

67. 青年女性，半年前刮宫诊断为葡萄胎，近年出现咳嗽咯血，胸透发现两肺有团块状阴影，血清 HCG 升高，穿刺活检可见异型性显著的两种滋养层细胞，无间质、无绒毛结构。该患者应诊断为
A. 肺鳞状细胞癌
B. 肺小细胞癌
C. 绒毛膜癌肺转移
D. 恶性葡萄胎肺转移
E. 肺腺癌

68. 绒毛膜癌的病理特点是
A. 有纤维组织性癌间质
B. 有丰富的毛细血管
C. 有丰富的淋巴细胞
D. 形成绒毛样结构
E. 肿瘤细胞为合体细胞样和细胞滋养层细胞样

69. 不符合绒毛膜癌的描述是

A. 可由葡萄胎恶变而来
B. 可发生于流产后
C. 可发生于正常分娩后
D. 可发生于卵巢、睾丸
E. 可发生于脑组织

70. 不符合绒毛膜癌的描述是
A. 与不全流产无关
B. 与葡萄胎关系密切
C. 无胎盘绒毛样结构
D. 可产生绒毛膜促性腺激素
E. 转移癌可比原发癌明显

71. 女性，40 岁，半年前曾患葡萄胎，经刮宫治疗。近 1 个月来，阴道出血，刮宫送病理检查，大量凝血坏死组织内可见异型性显著的合体滋养层细胞和细胞滋养层细胞。符合该患者的病理诊断是
A. 子宫内膜腺癌
B. 绒毛膜癌
C. 葡萄胎
D. 子宫平滑肌瘤
E. 子宫内膜腺鳞癌

72. 青年女性，停经 3 个月，近来发现阴道结节，镜检找到变性绒毛，最大可能性是
A. 宫外孕
B. 葡萄胎
C. 绒毛膜癌
D. 恶性葡萄胎
E. 葡萄状肉瘤

73. 女性生殖器官结核的原发部位是
A. 卵巢
B. 输卵管
C. 子宫
D. 子宫颈
E. 阴道

74. 急性输卵管炎的病原体是
A. 结核杆菌
B. 衣原体
C. 化脓菌
D. 支原体
E. 病毒

75. 输卵管卵巢囊肿是
A. 真性肿瘤
B. 错构瘤
C. 子宫内膜异位
D. 慢性炎症病变
E. 囊腺瘤

76. 不符合输卵管卵巢脓肿的描述是
A. 多发生于分娩后、流产后或刮宫术后
B. 多由化脓菌感染引起
C. 多由血道感染引起
D. 可形成输卵管卵巢囊肿
E. 输卵管卵巢粘连

77. 异位妊娠最常见的部位是
A. 卵巢
B. 输卵管
C. 子宫颈
D. 盆腔
E. 腹腔

78. 输卵管妊娠的最常见原因是
A. 输卵管发育异常
B. 慢性输卵管炎

C. 输卵管瘘管
D. 受精卵发育异常
E. 输卵管肿瘤

79. 输卵管妊娠最常见的合并症是
A. 腹腔出血
B. 阴道出血
C. 胚胎钙化
D. 腹腔妊娠
E. 肠粘连

80. 输卵管妊娠的好发部位是
A. 伞端
B. 壶腹部
C. 狭部
D. 间质部
E. 以上都不是

81. 不符合卵巢浆液性囊腺癌的描述是
A. 乳头多
B. 单层上皮
C. 细胞有异型性
D. 可见核分裂像
E. 包膜和间质有浸润

82. 引起卵巢 Krukenberg 瘤的是
A. 宫颈癌
B. 乳腺癌
C. 胃肠道癌
D. 肺癌
E. 肝癌

83. 卵巢最常见的肿瘤是
A. Krukenberg 瘤
B. 上皮性肿瘤
C. 性腺间质肿瘤
D. 生殖细胞肿瘤
E. 纤维瘤

84. 低度恶性的上皮性肿瘤是
A. 浆液性囊腺瘤
B. 粘液性囊腺瘤
C. 交界性浆液性囊腺瘤
D. 浆液性囊腺癌
E. 粘液性囊腺癌

85. 不符合浆液性囊腺瘤的是
A. 圆形囊肿
B. 清亮浆液
C. 乳头形成
D. 被覆上皮呈复层结构
E. 单房或多房

86. 不符合浆液性囊腺癌的是
A. 多囊性
B. 囊内实性菜花状突起
C. 上皮细胞复层排列
D. 上皮细胞异型性明显
E. 多无包膜和间质浸润

87. 不符合交界性浆液性囊腺瘤的是
A. 圆形囊肿
B. 清亮浆液
C. 多量乳头形成
D. 被覆上皮呈 2～3 层
E. 多有包膜和间质浸润

88. 不符合交界性粘液性囊腺瘤的是
A. 囊性肿物
B. 浓稠粘液
C. 多量乳头形成
D. 被覆上皮呈多层
E. 多无包膜和间质浸润

89. 最常见的良性卵巢性腺间质肿瘤是

A. 颗粒细胞瘤
B. 泡膜细胞瘤
C. 卵巢纤维瘤
D. 两性母细胞瘤
E. 支持细胞-间质细胞瘤

90. 最常见的低度恶性卵巢性腺间质肿瘤是
A. 颗粒细胞瘤
B. 泡膜细胞瘤
C. 卵巢纤维瘤
D. 两性母细胞瘤
E. 支持细胞-间质细胞瘤

91. Call-Exner 小体多见于
A. 卵巢颗粒细胞瘤
B. 卵巢无性细胞瘤
C. 卵巢纤维瘤
D. 内胚窦瘤
E. 卵巢门细胞瘤

92. 不符合颗粒细胞瘤的描述是
A. 常产生雌激素
B. 可引起月经紊乱
C. 多为实性
D. 可见 Call-Exner 小体
E. 高度恶性

93. 不符合泡膜细胞瘤的描述是
A. 常产生雌激素
B. 低度恶性
C. 多为实性
D. 肿瘤细胞与泡膜细胞相似
E. 绝经期妇女多见

94. 最常见的卵巢良性生殖细胞肿瘤是
A. 精原细胞瘤
B. 内胚窦瘤
C. 良性畸胎瘤
D. 恶性畸胎瘤
E. 胚胎性癌

95. 不符合良性畸胎瘤的描述是
A. 多为囊性
B. 囊内含毛发油脂
C. 囊壁可见骨、软骨
D. 囊壁可见神经组织
E. 肿瘤来源于性腺间质

96. 不符合恶性畸胎瘤的描述是
A. 多为囊性
B. 常有出血坏死
C. 囊壁可见骨、软骨
D. 囊壁可见胚胎性组织
E. 肿瘤内可见未成熟的神经组织

97. 生殖细胞向胚外组织分化而形成的肿瘤是
A. 颗粒细胞瘤
B. 内胚窦瘤
C. 粘液性囊腺瘤
D. Brenner 瘤
E. 浆液性囊腺瘤

98. 未成熟畸胎瘤的恶性成分是
A. 皮肤及其附属器
B. 软骨组织
C. 脑组织
D. 原始神经上皮
E. 平滑肌

99. 卵巢无性细胞瘤相当于
A. 精原细胞瘤
B. 内胚窦瘤

C. 两性母细胞瘤
D. 颗粒细胞瘤
E. 胚胎性癌

100. 不符合内胚窦瘤的描述是
A. 由生殖细胞发生
B. 低度恶性
C. 产生甲胎蛋白
D. 形成内胚窦样结构
E. 对放射治疗不敏感

101. 不属于生殖细胞起源的肿瘤是
A. 无性细胞瘤
B. 内胚窦瘤
C. 绒毛膜癌
D. 畸胎瘤
E. 颗粒细胞瘤

102. 不符合卵巢绒毛膜癌的描述是
A. 由生殖细胞发生
B. 低度恶性
C. 产生 HCG
D. 常合并其他生殖细胞肿瘤
E. 由异型性明显的滋养层细胞构成

103. 甲胎蛋白阳性的肿瘤是
A. 卵巢颗粒层细胞瘤
B. 卵巢无性细胞瘤
C. 卵巢纤维瘤
D. 内胚窦瘤
E. 卵巢门细胞瘤

104. 不符合卵巢无性细胞瘤的描述是
A. 由生殖细胞发生
B. 对放射治疗不敏感
C. 肿瘤呈实性、常有出血坏死
D. 常伴淋巴细胞浸润
E. 由体积大而圆的肿瘤细胞构成

105. 乳腺癌最常发生于乳房的
A. 外上象限
B. 外下象限
C. 内上象限
D. 内下象限
E. 中央部

106. 一般不出现乳房内包块的乳腺癌是
A. 导管内癌
B. 浸润性导管癌
C. 小叶原位癌
D. 浸润小叶癌
E. 髓样癌

107. 与长期雌激素刺激有关的肿瘤是
A. 子宫颈鳞状上皮癌
B. 子宫颈腺癌
C. 绒毛膜癌
D. 乳腺癌
E. 卵巢乳头状囊腺癌

108. 外上象限乳腺癌最常转移到
A. 同侧腋窝淋巴结
B. 同侧锁骨上淋巴结
C. 同侧锁骨下淋巴结
D. 同侧内乳淋巴结
E. 同侧肋间淋巴结

109. 不符合乳腺癌的描述是
A. 肿瘤灰白、质硬
B. 呈浸润性生长
C. 容易发生血道转移
D. 皮肤可呈桔皮样改变
E. 可出现乳头回缩现象

110. 女性，52 岁，右乳外上象限肿块 2 年

余，肿块质硬、边界不清。局部皮肤破溃，患侧乳头内陷。符合该患者疾病的诊断是

A. 乳腺增生症
B. 乳腺纤维腺瘤
C. 乳腺癌
D. 乳腺导管内乳头状瘤
E. 乳腺脓肿

111. 乳腺硬癌的特点是

A. 癌细胞局限于小叶腺泡内
B. 癌细胞局限于导管内
C. 癌细胞丰富、间质少
D. 纤维间质丰富，癌细胞少
E. 癌细胞与间质大致相等

112. 乳腺髓样癌的特点是

A. 癌细胞局限于小叶腺泡内
B. 癌细胞局限于导管内
C. 癌细胞丰富、间质少
D. 纤维间质丰富，癌细胞少
E. 癌细胞与间质大致相等

113. 乳腺实性癌的特点是

A. 癌细胞局限于小叶腺泡内
B. 癌细胞局限于导管内
C. 癌细胞丰富、间质少
D. 纤维间质丰富，癌细胞少
E. 癌细胞与间质大致相等

114. 乳腺小叶原位癌的特点是

A. 癌细胞局限于小叶腺泡内
B. 癌细胞局限于导管内
C. 癌细胞丰富、间质少
D. 纤维间质丰富，癌细胞少
E. 癌细胞与间质大致相等

115. 乳腺导管内癌的特点是

A. 癌细胞局限于小叶腺泡内
B. 癌细胞局限于导管内
C. 癌细胞丰富、间质少
D. 纤维间质丰富，癌细胞少
E. 癌细胞与间质大致相等

116. 妇女最常见的乳腺良性肿瘤是

A. 乳腺纤维腺瘤
B. 乳腺腺瘤
C. 叶状瘤
D. 导管内乳头状瘤
E. 脂肪瘤

117. 与遗传性乳腺癌密切相关的基因是

A. APC 基因
B. BRCA1 和 BRC2 基因
C. NF-1 基因
D. RB 基因
E. WT-1 基因

118. 不符合乳腺癌的描述是

A. 多来源于导管上皮细胞
B. 多发生于外上象限
C. 以鳞癌多见
D. 多数为实性癌
E. 以淋巴道转移多见

119. 金黄色葡萄球菌引起

A. 导管周围乳腺炎
B. 乳腺导管扩张症
C. 脂肪坏死
D. 肉芽肿性乳腺炎
E. 急性乳腺炎

120. 不符合乳腺纤维囊性病的描述是

A. 与雌激素分泌过多有关

B. 囊肿形成
C. 纤维增生
D. 腺泡和导管增生
E. 导管乳头状瘤

121. 符合前列腺癌的描述是
A. 对放射线敏感
B. 鳞癌多见
C. 容易转移到脊椎
D. 青年患者多见
E. 产生碱性磷酸酶

122. 前列腺癌的肿瘤标记物是
A. 碱性磷酸酶
B. 酸性磷酸酶
C. ATP 酶
D. 淀粉酶
E. 琥珀酸脱氢酶

123. 前列腺癌最常见的好发部位是
A. 外区
B. 后叶
C. 中叶
D. 前叶
E. 两叶以上

124. 前列腺癌最常见的组织学类型是
A. 移行细胞癌
B. 腺鳞癌
C. 鳞状细胞癌
D. 腺癌
E. 透明细胞癌

125. 前列腺癌最常见的组织学类型是
A. 移行细胞癌
B. 腺鳞癌
C. 鳞状细胞癌
D. 高分化腺癌
E. 低分化腺癌

126. 前列腺癌的肿瘤标记物是
A. 癌胚抗原
B. 甲胎蛋白
C. 碱性磷酸酶
D. 淀粉酶
E. 前列腺特异性抗原

127. 不符合前列腺癌的描述是
A. 老年患者多见
B. 多发生于前列腺外区
C. 多为高分化腺癌
D. 可引起排尿困难
E. 多继发于前列腺增生

128. 符合前列腺癌扩散的描述是
A. 可直接侵犯周围器官
B. 可转移到髂内、髂外淋巴结
C. 可转移到腹股沟淋巴结
D. 可转移到肺
E. 骨转移少见

129. 前列腺增生最常见于
A. 内区
B. 后叶
C. 前叶
D. 两侧叶
E. 两叶以上

130. 不符合前列腺增生症的病理变化是
A. 前列腺腺上皮异型性显著
B. 前列腺腺体增生
C. 前列腺间质纤维组织增生
D. 前列腺平滑肌增生
E. 前列腺腺体肥大

131. 在前列腺增生症发生中，最重要的激素是
A. 双氢睾丸酮
B. 促性腺激素
C. 生长激素
D. 雌激素
E. 泌乳素

132. 前列腺增生症最常见的临床症状是
A. 血尿
B. 蛋白尿
C. 尿频、尿急、尿痛
D. 排尿困难
E. 脓尿

133. 不符合前列腺增生症的描述是
A. 老年患者多见
B. 多发生于前列腺内区
C. 可继发尿路感染
D. 常引起排尿困难
E. 多引起前列腺癌发生

134. 不符合前列腺增生症的描述是
A. 中青年患者多见
B. 多发生于前列腺内区
C. 可继发尿路感染
D. 常引起排尿困难
E. 属良性增生性病变

135. 不符合前列腺增生症的描述是
A. 老年患者多见
B. 多发生于前列腺内区
C. 属癌前病变
D. 常引起排尿困难
E. 可继发尿路感染

136. 不符合前列腺增生症的病理变化是
A. 多累及前列腺外区
B. 前列腺腺体增生
C. 前列腺间质纤维组织增生
D. 前列腺平滑肌增生
E. 前列腺腺体肥大

137. 男性，30 岁，左侧睾丸逐渐肿大 2 个月，近来常咳嗽、咯血，体检时发现左侧睾丸比正常大 3 倍，胸部拍片见左肺有 3cm×4cm 阴影，肺叶切除见肿瘤由两种滋养细胞组成。睾丸肿瘤可能是
A. 精原细胞瘤
B. 胚胎性癌
C. 内胚窦瘤
D. 绒毛膜癌
E. 恶性畸胎瘤

138. 睾丸肿瘤中对放射治疗最敏感的是
A. 精原细胞瘤
B. 胚胎性癌
C. 内胚窦瘤
D. 绒毛膜癌
E. 恶性畸胎瘤

139. 睾丸的性索间质性肿瘤是
A. 精原细胞瘤
B. 间质细胞瘤
C. 内胚窦瘤
D. 绒毛膜癌
E. 恶性畸胎瘤

140. 与隐睾关系密切的肿瘤是
A. 精原细胞瘤
B. 胚胎性癌
C. 内胚窦瘤
D. 绒毛膜癌

E. 恶性畸胎瘤

141. 不符合睾丸精原细胞瘤的描述是

A. 隐睾患者发病率高

B. 睾丸肿大

C. 肿瘤实性、鱼肉状

D. 肿瘤细胞体积大、形态较一致

E. 常转移到腹股沟淋巴结

142. 引起 HCG 升高的睾丸肿瘤是

A. 内胚窦瘤

B. 胚胎性癌

C. 间质细胞瘤

D. 精原细胞瘤

E. 恶性畸胎瘤

143. 引起 AFP 升高的睾丸肿瘤是

A. 间质细胞瘤

B. 胚胎性癌

C. 绒毛膜癌

D. 精原细胞瘤

E. 恶性畸胎瘤

144. 不符合睾丸胚胎性癌的描述是

A. 对放射线敏感

B. 恶性度高

C. 癌组织结构多样

D. 转移到髂内、髂总淋巴结

E. 转移到肝、肺等器官

145. 阴茎龟头的 Queyrat 增殖性红斑属于

A. 瘤样病变

B. 癌前病变

C. 原位癌

D. 早期侵润性癌

E. 良性肿瘤

146. 引起睾丸树胶样肿的疾病是

A. 肉芽肿性睾丸炎

B. 睾丸结核

C. 睾丸梅毒

D. 腮腺炎性睾丸炎

E. 隐睾症

147. 阴茎癌常可早期转移到

A. 腹股沟淋巴结

B. 腹膜后淋巴结

C. 盆腔淋巴结

D. 髂外淋巴结

E. 主动脉旁淋巴结

148. 梅毒可引起

A. 软性下疳

B. 硬性下疳

C. 精子肉芽肿

D. 肉芽肿性睾丸炎

E. 糜烂性龟头炎

149. 男性生殖系统结核最常累及

A. 睾丸

B. 附睾

C. 输精管

D. 精囊

E. 前列腺

150. 引起阴茎癌的病毒是

A. HIV

B. CMV

C. HBV

D. HPV

E. HCV

151. 符合阴茎癌的描述是

A. 多为低分化鳞癌

B. 多见于包皮过长患者

C. 早期发生血道转移
D. 与 HPV 感染无关
E. 与吸烟有关

152. 不符合阴茎癌的描述是
A. 多为低分化鳞癌
B. 多见于包皮过长患者
C. 以淋巴道转移为主
D. 与 HPV 感染有关
E. 可侵犯阴茎海绵体

153. condyloma acuminatum
A. 阴茎癌
B. 胚胎性癌
C. 尖锐湿疣
D. 硬化性苔癣
E. 子宫内膜异位症

154. endometriosis
A. 阴茎癌
B. 胚胎性癌
C. 尖锐湿疣
D. 硬化性苔癣
E. 子宫内膜异位症

155. hydatidiform mole
A. 葡萄胎
B. 胚胎性癌
C. 尖锐湿疣
D. 硬化性苔癣
E. 子宫内膜异位

156. borderline serous cystadenoma
A. 交界性浆液性囊腺瘤
B. 交界性粘液性囊腺瘤
C. 浆液性囊腺瘤
D. 粘液性囊腺瘤
E. 粘液性囊腺癌

157. granulosa cell tumor
A. 颗粒细胞瘤
B. 泡膜细胞瘤
C. 支持细胞瘤
D. 间质细胞瘤
E. 良性母细胞瘤


158. cervicitis
A. 子宫内膜炎
B. 输卵管炎
C. 阴道炎
D. 子宫颈炎
E. 子宫内膜异位症

159. seminoma
A. 颗粒细胞瘤
B. 泡膜细胞瘤
C. 支持细胞瘤
D. 间质细胞瘤
E. 精原细胞瘤

160. hyperplasia of prostate
A. 子宫内膜增生
B. 前列腺增生
C. 乳腺增生
D. 鳞状上皮异型增生
E. 滋养层细胞增生

英文题

1. A 45-year-old woman has had a small amount of vaginal bleeding and a brownish, foul-smelling discharge for the past month. On pelvic examination, there is a 3-cm lesion on the ectocervix. Microscopic findings of this lesion are shown in the figure. Which of the following is the most likely diagnosis?



A. Sertoli-Leydig cell tumor
B. Mature cystic teratoma
C. Adenocarcinoma
D. Squamous cell carcinoma
E. Granulosa cell tumor

2. A 35-year-old woman has had increasing abdominal enlargement for the past 6 months. On physical examination, there is abdominal distention with a fluid wave. A pelvic ultrasound scan shows bilateral cystic ovarian masses, 10 cm on the right and 7 cm on the left. The masses are surgically removed. On gross examination, the excised masses are unilocular cysts filled with clear fluid, and papillary projections extend into the central lumen of these cysts. Microscopic examination shows that the papillae are covered with atypical cuboidal cells that invade underlying stroma. Psammoma bodies are present. Which of the following is the most likely diagnosis?

A. Sertoli-Leydig cell tumor
B. Mature cystic teratoma
C. Cystadenocarcinoma
D. Dysgerminoma
E. Granulosa cell tumor

3. A l4-year-old girl began menstruation 1 year ago. She now has abnormal uterine bleeding. On physical examination, there are no remarkable findings. A pelvic ultrasound scan shows no abnormalities. Which of the following is most likely to produce these findings?

A. Endometrial polyp
B. Anovulatory cycles
C. Ectopic pregnancy
D. Uterine leiomyomata
E. Endometrial carcinoma

4. A 20-year-old woman notices a bloody, brownish vaginal discharge. On physical examination, a 3-cm red-brown mass is seen on the lateral wall of the vagina. A chest radiograph shows numerous 2-to 5-cm nodules in both lungs. A biopsy specimen of the vaginal mass shows malignant cells resembling syncytiotro-

phoblasts. Which the following proteins is most likely to be elevated in the serum?

A. Human chorionic gonadotropin
B. α-Fetoprotein
C. Estradiol
D. Testosterone
E. Careinoembryonic antigen

5. A 42-year-old woman has a Pap smear as part of a routine health maintenance examination. A cervical biopsy is performed, and microscopic examination shows dysplastic cells that occupy the full thickness of the epithelium above the basement membrane (CIN III) . Which of the following is the best rational for proceeding with cervical conization in this patient?

A. She has a high risk of invasive carcinoma
B. Human papinomavirus infection cannot be treated
C. She is perimenopausal
D. She has chronic cervicitis
E. Her reproductive years are over

6. A 21-year-old woman who is sexually active has noticed multiple warty lesions on the perineum over the past 3 years. A scraping from one of the lesions yields cells with cytologic features consistent with koilocytotic atypia. Which of the following infectious agents is most likely to produce these findings?

A. Neisseria gonorrhoeae
B. Chlamydia trachomatis
C. Human papillomavirus
D. Trichomonas vaginalis
E. Herpes simplex virus type 2

7. A 32-year-old woman has cyclic abdominal pain that coincides with her menses. Attempts to become pregnant have failed over the past 5 years. There are no abnormal findings on physical examination. Laparoscopic examination shows numerous hemorrhagic 0.2-to 0.5-cm lesions over the peritoneal surfaces of the uterus and ovaries. Which of the following ovarian lesions is most likely to be visualized during the laparoscopic procedure?

A. Fibroma
B. Brenner tumor
C. Endometriotic cyst
D. Krukenberg tumor
E. Mature cystic teratoma

8. An 18-year-old woman has had pelvic discomfort for several months. On pelvic examination, there is a 10-cm right adnexal mass. Upon surgical resection, the mass is solid and white, with small areas of necrosis. Microscopically, it contains mostly primitive mesenchymal cells along with some cartilage, muscle, and foci of neuroepithelial differentiation. Which of the following is the most likely diagnosis?

A. Brenner tumor

B. Dysgerminoma
C. Granulosa cell tumor
D. Immature teratoma
E. Leiomyosarcoma

9. A 43-year-old woman has had postcoital bleeding for 6 months. A Pap smear shows high-grade cervical intraepithelial neoplasia (CIN III). In situ hybridization performed on cells from the cervix shows the presence of human papillomavirus type 16. If the cervical lesion is not treated, which of the following malignancies is she at greatest risk for developing?
A. Clear ceil carcinoma
B. Squamous cell carcinoma
C. Krukenberg tumor
D. Leiomyosarcoma
E. Papillary serous cystadenocarcinoma

10. A 58-year-old woman has had dull pain in the lower abdomen for the past 6 months and minimal vaginal bleeding on three occasions. Pelvic examination shows a right adnexal mass, and the uterus appears normal in size. A total abdominal hysterectomy is performed, and the mass is diagnosed as an ovarian granulosa-theca cell tumor. Which of the following additional lesions is most likely to be found in the excised specimen?
A. Condyloma acuminata of the cervix
B. Endometrial hyperplasia
C. Metastases to the uterine serosa
D. Bilateral chronic salpingitis
E. Partial mole of the uterus

11. A 50-year-old woman has noticed vaginal bleeding for the past month. Her last menstrual period was 5 years ago. On physical examination, there are no abnormal findings. An endometrial biopsy shows the histological findings above. Which of the following is the most likely diagnosis?
A. Adenoma
B. Leiomyoma
C. Choriocarcinoma
D. Adenocarcinoma
E. Adenomyosis

12. A 28-year-old sexually active woman comes to her physician for a routine health maintenance examination. A Pap smear shows a moderate dysplasia, or cervical intraepithelial neoplasia (CIN) II. Which of the following is the significance of this finding?
A. A cervicitis needs to be treated
B. The patient has an increased risk of cervical carcinoma
C. Condyloma acuminata are probably present
D. An endocervical polyp needs to be excised
E. The patient should not go to her physician at regular intervals

13. A 42-year-old woman has had menometr-

orrhagia for the past 2 months. On physical examination, there are no vaginal or cervical lesions, and the uterus appears to be of normal size, but there is a right adnexal mass. An abdominal ultrasound scan shows the presence of a 7-cm solid right adnexal mass. Endometrial biopsy shows hyperplastic endometrium but no cellular atypia. Which of the following is the most likely diagnosis?

A. Mature cystic teratoma
B. Endometrioma
C. Corpus luteum cyst
D. Metastasis
E. Granulosa-theca cell tumor

14. The most common histologic type of cancer of cervix is

A. Squamous carcinoma
B. Mucinous carcinoma
C. Papillary carcinoma
D. Lymphoma
E. Leiomyosarcoma

15. The most common histologic type of tumor of the uterus is

A. Stromal tumor
B. Leiomyoma
C. Mesothelioma
D. Angioma
E. Adenoma

16. The most common histologic type of germ cell tumor occurring in the ovary is

A. Embyonal carcinoma
B. Yolk sac tumor
C. Granulosa cell tumor
D. Clear cell tumor
E. Teratoma

17. A 23-year-old woman, G3, P2, has a spontaneous abortion at 15 weeks gestation. The male fetus is small for gestational age. The placenta is also small and shows 0.5-cm grapelike villi scattered among morphologically normal villi. Chromosomal analysis of placental tissue is most likely to show which of the following karyotypes?

A. 69, XXY
B. 46, XX
C. 23, Y
D. 45, X
E. 47, XXY

18. A 26-year-old woman in her fifteenth week of pregnancy presents with uterine bleeding and passage of a small amount of watery fluid and tissue. She is found to have a uterus that is much larger than estimated by her gestational dates. Her uterus is found to be filled with cystic, avascular, grapelike structures that do not penetrate the uterine wall. No fetal parts are found. Which of the following is the best diagnosis?

A. Partial hydatidiform mole
B. Complete hydatidiform mole
C. Invasive mole

D. Adenocarcinoma
E. Choriocarcinoma

19. A 27-year-old woman experiences a sudden onset of severe lower abdominal pain. On pelvic examination there are no lesions of the cervix or vagina. An abdominal ultrasound shows a 4-cm focal enlargement of the proximal right fallopian tube. A dilation and curettage procedure shows only decidua in the endometrial cavity. Which of the following laboratory findings is most likely to be reported for this patient?
A. Cervical culture positive for Neisseria gonorrhoeae
B. 69, XXY karyotype on decidual tissue
C. Positive result of serum pregnancy test
D. Positive result of serologic testing for syphilis
E. Pap smear showing Candida

20. A 27-year-old woman acutely develops lower abdominal pain and vaginal bleeding. While in the bathroom she passes a cast of tissue com-posed of clot material and then collapses. She is brought to the hospital, where a physical examination reveals a soft, tender mass in the right adnexa and pouch of Douglas. Histologic examination of the tissue passed in the bathroom reveals blood clot and decidualized tissue. No chorionic villi or trophoblastic tissue are present. Which of the following conditions is most likely present in this individual?
A. Aborted intrauterine pregnancy
B. Complete hydatidiform mole
C. Ectopic pregnancy
D. Endometrial hyperplasia
E. Partial hydatidiform mole

21. An 28-year-old woman presents with amenorrhea and is found to have normal secondary sex characteristics. She states that her last menstrual period was 8 weeks prior to this visit. A urine test for hCG is positive. Which of the fol-lowing is the most likely diagnosis?
A. leiomyoma
B. Intrauterine pregnancy
C. Stromal tumor
D. Turner's syndrome
E. Endometrial carcinoma

22. A 39-year-old woman presents with increasing abdominal distention and pelvic pain. A CT scan shows a 3.5 cm tumor of the right ovary. The tumor is resected and histologic sections reveal the tumor to be composed of spindle-shaped cells with mild atypia. Which of the following types of ovarian tumor is most likely to produce this constella-tion of findings?
A. Epithelial tumor
B. Stromal tumor
C. Germ cell tumor
D. Surface cell tumor

E. Metastatic carcinoma


23. A 29-year-old woman presents with intermittent lower abdominal pain. A pelvic examination reveals an ovarian mass involving her right ovary. This ovarian mass is resected and gross examination reveals a cystic tumor that measures about 4 cm in diameter. A histologic section from this tumor reveals a mixture of elements as seen in the picture. Neither immature nor neural tis-sue is found. Which of the following is the most likely diagnosis?

A. Brenner tumor
B. Chronic salpingitis
C. Dermoid cyst
D. Ectopic pregnancy
E. Mucinous cystadenoma

24. A 32-year-old woman presents with a recent onset of oligomenor-rhea followed by amenorrhea. Physical examination reveals loss of female secondary characteristics and slight abdominal distention. She is found to have a 4.0 cm mass within her left ovary. This mass is resected surgically and a histologic section reveals the tumor cells focally form Call-Exner bodies. What is the correct diagnosis?

A. Serous cystadenocarcinoma
B. Borderline mucinous tumor
C. Granulosa cell tumor
D. Pseudomyxoma peritonei
E. Sertoli cell tumor

25. A 40-year-old woman has noticed progressive enlargement of the abdomen over the past 5 months. Physical examination shows no palpable masses, but a fluid wave is present. An abdominal ultrasound scan shows a 15-cm multilobular mass that involves the right adnexal region. The mass is surgically removed, and the gross features of the excised mass are shown above. Which of the following is the most likely diagnosis?



A. Immature teratoma
B. Serous cystadenocarcinoma
C. Granulosa cell tumor
D. Choriocarcinoma
E. Dysgerminoma

26. A 54-year-old woman has had weight loss accompanied by abdominal enlargement for the past 6 months. She is concerned because of a family history of ovarian carcinoma. On physical examination, there are no lesions of the cervix, and the uterus is normal in size, but there is a left adnexal mass. An abdominal ultrasound scan shows a l0-cm cystic mass in the left adnexal region, with scattered 1-cm peri-

toneal nodules. Cytologic studies of peritoneal fluid show malignant cells consistent with a cystadenocarcinoma. Which of the following mutated genes is most likely a factor in the development of this neoplasm?

A. RAS
B. BRCA1
C. ERBB2 (HER2)
D. MYC
E. RBl

27. Over the past 9 months, a 30-year-old man has noticed a feeling of heaviness with enlargement of the scrotum. On physical examination, there is an enlarged, firm left testis. A left orchiectomy is performed. Microscopic examination of the mass shows areas of mature cartilage, keratinizing squamous epithelium, and colonic glandular epithelium. Laboratory findings include elevated levels of serum human chorionic gonadortropin (hCG) and α-fetoprotein (AFP). Despite the appearance of the cells in the tumor, the surgeon tells the patient that he probably ha s a malignant testicular tumor. The surgeon's conclusion is most likely to be based on which of the following factors?

A. Size of the tumor
B. Age of the patient
C. Presence of colonic glandular epithelium
D. Elevation of hCG and AFP levels
E. Location of the mass in the left testis

28. A 28-year-old woman presents with pelvic pain and is found to have an ovarian mass of the left ovary that measures 3 cm in diameter. Grossly, the mass consists of multiple cystic spaces. Histologically, these cysts are lined by tall columnar epithelium, and some of the cells are ciliated. Which of the following is the correct diagnosis for this ovarian tumor?

A. Brenner tumor
B. Mucinous tumor
C. Endometriosis
D. Clear cell tumor
E. Serous tumor

29. An otherwise healthy 72-year-old man has had increasing difficulty with urination for the past 10 years. On physical examination, the prostate is enlarged to twice its normal size but is not tender to palpation. The serum prostate-specific antigen (PSA) level is normal. Which of the following findings is most likely to be seen microscopically in a biopsy specimen of the prostate gland?

A. Hyperplastic nodules of stroma and glands lined by two layers of epithelium
B. Poorly differentiated glands lined by a single layer of epithelium and packed back to back

C. Foci of chronic inflammatory cells in the stroma and in normal-appearing glands
D. Areas of liquefactive necrosis filled with neutrophils
E. Multiple areas of caseating granulomatous inflammation

30. Prolonged unopposed estrogen stimulation in an adult woman increases the risk of development of endometrial hyperplasia and subse-quent carcinoma. In such cases which of the following is the most common histologic type of cancer?
A. Adenocarcinoma
B. Clear cell carcinoma
C. Small cell carcinoma
D. Squamous cell carcinoma
E. Transitional cell carcinoma

31. A 33-year-old man has noted asymmetric enlargement of the scrotum over the past 4 months. On physical examination, the right testis is twice its normal size and there it increased tenderness to palpation. The right testis is removed. The epididymis and the upper aspect of the right testis have extensive granulomatous inflammation with epithelioid cells, Langhans giant cells, and caseous necrosis. Which of the following is the most likely cause of these findings?
A. Mumps
B. Syphilis
C. Tuberculosis
D. Gonorrhea
E. Sarcoidosis

32. A healthy 30-year-old woman comes to the physician for a routine physical examination. No abnormalities are found on review of systems or physical examination, but a screening Pap smear shows cells consistent with cervical intraepithelial neoplasia (CIN). Which of the following is most likely related to the Pap smear findings?
A. Intake of oral contraceptives
B. Diethylstilbestrol (DES) exposure
C. Vitamin B12 (cobalamin) deficiency
D. Treated squamous cell carcinoma
E. HPV infection


33. The mother of a 2-year-old child notices that he has had increasing asymmetric enlargement of the scrotum over the past 6 months. On physical examination, there is a well-circumscribed 2.5-cm mass in the left testis. Histologic examination of this mass shows sheets of cells and ill-defined glands composed of cuboidal cells, some of which contain eosinophilic hyaline globules. Microcysts and primitive glomeruloid structures also are seen. Immunohistochemical staining shows α-fetoprotein in the cytoplasm of the neoplastic cells. Which of the following is the most likely diagnosis?

A. Choriocarcinoma
B. Seminoma
C. Yolk sac tumor
D. Teratoma
E. Leydig cell tumor

34. A 65-year-old postmenopausal woman presents with a new onset of uterine bleeding. An endometrial biopsy is diagnosed as atypical hyper-plasia. Which of the following histologic changes is most characteristic of this abnormality?
A. Crowding of endometrial glands with budding and epithelial atypia
B. Lymphatic invasion by interlacing bundles of atypical spindle-shaped cells
C. Menstrual-type endometrial glands with focal cystic dilatation
D. Endometrial tissue was present within the wall of uterus
E. Stromal invasion by malignant glands with focal areas of atypical squamous differentiation

35. The most common histologic type of malignant tumor of the breast is
A. Angiosarcoma
B. Invasive ductal carcinoma
C. Phyllodes tumor
D. Lymphoma
E. Papillary carcinoma

36. A 70-year-old man comes to his physician for a routine health examination. On palpation, his prostate is normal in size. However, laboratory studies show an elevated level of serum prostate-specific antigen (PSA). Which of the following histologic findings in a subsequent biopsy of the prostate is most likely to account for the patient's elevated PSA?
A. Hyperplastic nodules of the stroma and glands lined by two layers of epithelium
B. Poorly differentiated glands lined by a single layer of epithelium and packed back to back
C. Foci of chronic inflammatory cells in the stroma and in normal-appearing glands
D. Areas of liquefactive necrosis filled with neutrophils
E. Multiple caseating granulomas



37. A healthy 52-year-old woman has had a feeling of pelvic heaviness for the past 11 months. There is no history of abnormal bleeding. Her physician palpates an enlarged nodular uterus on pelvic examination. Pelvic CT imaging shows multiple solid, uterine masses. A total abdomi-

nal hysterectomy is performed. Based on the gross appearance of the mass shown above, which of the following is the most likely diagnosis?

A. Metastases
B. Endometriosis
C. Infiltrative leiomyosarcoma
D. Multiple leiomyomas
E. Adenomyosis


38. The most common histologic type of benign tumor at breast is

A. Papilloma
B. Adenoma
C. Fibroadenoma
D. Angioma
E. Lipoma

39. A 32-year-old man has noticed an increased feeling of heaviness in his scrotum for the past 10 months. On physical examination, the left testis is three times the size of the right testis and is firm on palpation. An ultrasound scan shows a 6-cm solid mass within the body of the left testis. Laboratory studies include an elevated serum α-fetoprotein level. Which of the following is the most likely diagnosis?

A. Yolk sac tumor
B. Leydig cell tumor
C. Seminoma
D. Lymphoma
E. Embryonal carcinoma cells

40. A 31-year-old woman has had dull, constant, abdominal pain for 6 months. On physical examination, the only finding is a right adnexal mass. The right fallopian tube and ovary are surgically excised. The gross and histologic appearance of the ovary are shown in the figure above. Which of the following is the most likely diagnosis?



A. Mucinous cystadenoma
B. Choriocarcinoma
C. Dysgerminoma
D. Serous cystadenoma
E. Mature cystic teratoma

41. A 56-year-old woman presents with a slowly enlarging 2.5-cm firm, irregular mass in the upper outer quadrant of her left breast. A biopsy from this mass is interpreted by the pathologist as an infiltrating lobular carcinoma of the breast. Which of the following histologic features is most characteristic of this tumor?

A. Expansion of lobules by monotonous proliferation of epithelial cells
B. Large cells with clear cytoplasm within the epidermis
C. Large syncytium-like sheets of pleo-

morphic cells surrounded by aggregates of lymphocytes

D. Small individual malignant cells dispersed within extracellular pools of mucin

E. Small tumor cells with little cytoplasm infiltrating in a single-file pattern

42. A 30-year-old man comes to his physician because he has noticed increasing enlargement and a feeling of heaviness in his scrotum for the past year. On physical examination, the right testis is twice the normal size, and it is firm and slightly tender. An ultrasound examination shows a 3.5-cm solid mass in the right testis. An abdominal CT scan shows enlargement of the para-aortic lymph nodes. Laboratory findings include markedly increased serum levels of chorionic gonadotropin and α-fetoprotein. Which of the following testicular neoplasms does this patient most have?

A. Leydig cell tumor

B. Mixed germ cell tumor

C. Pure spermatocytic seminoma

D. Choriocarcinoma

E. Metastatic adenocarcinoma of the prostate gland

43. A 40-year-old woman presents with urinary frequency and abnormal uterine bleeding. A pelvic examination finds several masses in the wall of the uterus, and this is confirmed by ultrasonography. Which of the following is the most likely diagnosis?

A. Endometriosis

B. Rhabdomyosarcoma

C. Ectopic pregnancy

D. Adenoarcinoma

E. Leiomyomas

44. A 48-year-old woman presents with a 1.5-cm firm mass in the upper outer quadrant of her left breast. A biopsy from this mass reveals many of the ducts are filled with atypical cells. In the center of these ducts there is extensive necrosis. No invasion into the surrounding fibrous tissue is seen. Which of the following is the most likely diagnosis?

A. Colloid carcinoma

B. Intraductal carcinoma

C. Infiltrating ductal carcinoma

D. Infiltrating lobular carcinoma

E. Lobular carcinoma in situ

45. A 67-year-old man is found on rectal examination to have a single, hard, irregular nodule within his prostate. A biopsy of this lesion reveals the presence of small glands lined by a single layer of cells with enlarged, prominent nucleoli. Which of the following sub-stances is most likely to be increased in this patient's serum?

A. Acid phosphatase

B. Alpha-fetoprotein (AFP)

C. Alkaline phosphatase
D. Human chorionic gonadotropin (hCG)
E. Prostate-specific antigen (PSA)

46. A 39-year-old woman presents with severe menorrhagia and colicky dysmenorrhea. A hysterectomy including resection of the fallopian tubes and ovaries is performed and the histologic section shows endometrial tissue present within the wall of uterus. Which of the following is the most likely diagnosis?
A. Chronic endometritis
B. Adenocarcinoma
C. Ectopic pregnancy
D. Adenomyosis
E. Endometriosis

47. A 48-year-old woman presents with a painless mass located in her left breast. Physical examination confirms the presence of a firm, non-tender, 3-cm mass in the upper outer quadrant of her left breast. The mass is resected and histologic sections reveal an inva-sive ductal carcinoma. Biopsy of her axillary lymph nodes reveals the presence of metastatic disease in 4 of 18 axillary lymph nodes. The expression of which one of the following by the tumor would best predict a response to hormonal therapy?
A. BRCAI
B. Estrogen receptors
C. HER2/neu
D. AFP
E. CEA


48. A 78-year-old man is found on rectal examination to have a single, hard, irregular nodule within his prostate. A biopsy of this lesion reveals the presence of small glands lined by a single layer of cells with enlarged, prominent nucleoli. Which of the following is the most likely diagnosis?
A. Atrophic prostatitis
B. Atypical small acinar proliferation
C. High-grade prostatic intraepithelial neoplasia
D. Benign prostatic hyperplasia
E. Prostatic adenocarcinoma

49. A 35-year-old woman presents with severe menorrhagia and colicky dysmenorrhea. A hysterectomy including resection of the fallopian tubes and ovaries is performed and the pathologist who examines the specimen makes the diagnosis of "adenomyosis." Which one of the listed abnormali-ties was present in this resected specimen?
A. Ectocervical tissue was present within the wall of the fallopian tube
B. Endocervical tissue was present on the surface of the ovary
C. Endometrial tissue was present within the hilum of the ovary
D. Endometrial tissue was present within the wall of uterus

E. Ovarian tissue was present on the surface of the uterus

50. A 39-year-old woman presents with a new onset of a bloody dis-charge from her right nipple. Physical examination reveals a l-cm freely movable mass that is located directly beneath the nipple. Sections from this mass reveal multiple fibrovascular cores lined by several layers of epithelial cells. Atypia is minimal. The lesion is completely contained within the duct and no invasion into underlying tissue is seen. Which of the following is the most likely diagnosis?
A. Benign phyllodes tumor
B. Intraductal papilloma
C. Intraductal carcinoma
D. Paget's disease
E. Papillary carcinoma

51. A 72-year-old man is found on rectal examination to have a single, hard, irregular nodule within his prostate. A biopsy of this lesion reveals the presence of small glands lined by a single layer of cells with enlarged, prominent nucleoli. From what portion of the prostate did this lesion most likely originate?
A. Anterior zone
B. Central zone
C. Peripheral zone
D. Periurethral glands
E. Transitional zone



52. A 45-year-old woman has had menometrorrhagia for the past 3 months. On physical examination, there are no remarkable findings. The microscopic appearance of an endometrial biopsy specimen is shown in the photomicrograph above. Which of the following conditions is most likely to produce these findings?
A. Ovarian mature cystic teratoma
B. Chronic endometritis
C. Failure of ovulation
D. Pregnancy
E. Use of oral contraceptives

53. A 23-year-old woman presents with a rubbery, freely movable 2-cm mass in the upper outer quadrant of the left breast. Which of the following histologic features is most likely to be seen when examining a biopsy spec-imen from this mass?
A. Large numbers of neutrophils
B. Large numbers of plasma cells
C. Duct ectasia with inspissation of breast secretions
D. Necrotic fat surrounded by lipid-laden macrophages
E. A mixture of fibrous tissue and ducts

54. A 66-year-old man presents with urinary frequency, nocturia, drib-bling, and difficulty in starting and stopping urination. Rectal examination reveals the prostate to be enlarged, firm, and rubbery. A needle biopsy reveals increased numbers of glandular elements and stromal tissue. The glands are found to have a double layer of epithelial cells. Prominent nuclei or back-to-back glands are not seen. From what portion of the prostate did this lesion most likely originate?
    A. Anterior zone
    B. Central zone
    C. Peripheral zone
    D. Periurethral glands
    E. Transitional zone
55. A 39-year-old woman presents with excessive bleeding at irregular intervals. An endometrial biopsy is obtained. Histologic sections reveal menstrual-type endometrial glands with focal cystic dilatation and the epithelial cells show little atypia. Which of the following is the most likely diagnosis?
    A. Endometrial hyperplasia
    B. Adenocarcinoma
    C. Chronic endometritis
    D. Adenomyosis
    E. Endometriosis
56. HER/neu staining is used to predict:
    A. Tumor grade
    B. Response to anti-HER2 therapy
    C. Response to hormone manipulation
    D. Lymph node status
    E. Increase risk for adenoma in the contralateral breast
57. A 69-year-old man presents with urinary frequency, nocturia, drib-bling, and difficulty in starting and stopping urination. A needle biopsy reveals increased numbers of glandular elements and stromal tissue. The glands are found to have a double layer of epithelial cells without atypia. Which of the following is the most likely diagnosis?
    A. Atrophic prostatitis
    B. Atypical small acinar proliferation
    C. High-grade prostatic intraepithelial neoplasia
    D. Benign prostatic hyperplasia
    E. Prostatic adenocarcinoma
58. A 40-year-old woman presents with a new onset of uterine bleeding. A careful medical history finds that her abnormal menstrual bleeding is characterized by excessive bleeding at irregular intervals. An endometrial biopsy is obtained. The pathology report from this specimen makes the diagnosis of endometriosis. Which of the following histologic changes is most characteristic of this abnormality?
    A. Crowding of endometrial glands with budding and epithelial atypia

B. Lymphatic invasion by interlacing bundles of atypical spindle-shaped cells
C. Menstrual-type endometrial glands with focal atypical cystic dilatation
D. Endometrial tissue was present within the wall of uterus
E. Stromal invasion by malignant glands with focal areas of atypical squamous differentiation

59. Which breast tumor has the best prognosis?
A. Mucinous carcinoma
B. Invasive lobular carcinoma
C. Fibroadenoma
D. Adenoid cystic carcinoma
E. Phyllodes tumor

60. A 47-year-old man presents with the sudden onset of fever, chills, and dysuria. Rectal examination finds the prostate gland to be edematous and very sensitive; examination is quite painful. Microscopic examination of prostatic secretions reveals the presence of numerous neutrophils. Which of the following is the most likely diagnosis?
A. Atrophic prostatitis
B. Atypical small acinar proliferation
C. Acute prostatitis
D. Benign prostatic hyperplasia
E. Prostatic adenocarcinoma

61. A 62-year-old obese nulliparous woman had an episode of vaginal bleeding. On pelvic examination, there appears to be no enlargement of the uterus, and the cervix appears normal. A Pap smear shows cells consistent with squamous cell carcinoma. Which of the following conditions is most likely to have contributed to the development of this malignancy?
A. Endometrial hyperplasia
B. Chronic endometritis
C. Use of oral contraceptives
D. Human papillomavirus infection
E. Adenomyosis

62. Moderate or florid ductal hyperplasia indicates
A. Slightly increased risk for subsequent invasive breast carcinoma
B. Subsequent development of invasive carcinoma in 50% of all patients
C. Has no predictive value
D. Slightly increase risk for phyllodes tumors
E. Slightly increase risk for tuberculosis

63. A 25-year-old man presents with a testicular mass, which is resected and diagnosed as a yolk sac tumor. Which of the following sub-stances is most likely to be increased in this patient's serum as a result of secretion from the cells of this tumor?
A. Acid phosphatase
B. a-fetoprotein (AFP)
C. Alkaline phosphatase

D. Human chorionic gonadotropin (hCG)
E. Prostate-specific antigen (PSA)

64. A 27-year-old woman presents to her gynecologist for a routine phys-ical examination that includes a Pap smear. Her sexual history includes many sexual partners beginning at an early age. Her cervix is painted with iodine and there is an area near the cervical os which does not stain with iodine. Several biopsies are obtained from this pale area, and a representative histologic section is seen in the picture below. This histologic section shows koilocytosis, which is most characteristic of infec-tion with which one of the following organisms?
A. Cytomegalovirus
B. Epstein-Barr virus
C. Herpes simplex virus
D. Human papillomavirus
E. Parvovirus B19

65. Hormone receptors are b
A. Most useful in chosing patients for polyagent chemotherapy
B. Most useful in chosing patients for hormonal therapy
C. Most useful as a prognostic test
D. Provide the therapeutic basis for chemotherapy
E. Provide the therapeutic basis for radiotherapy

66. The most common histologic type of malignant tumor occurring in the testis is
A. Seminoma
B. Adenomatoid tumor
C. Leukemia
D. Metastatic carcinoma
E. Leydig cell tumor?

67. Which one of the following numbered sequences best illustrates the postulated sequence of events that precedes the formation of an infiltrating squamous cell carcinoma of the cervix? (1=Carcinoma in situ；2=Invasive carcinoma；3=Mild dysplasia；4=Moderate dysplasia；5=Severe dysplasia；6=Squamous metaplasia)
A. 3, then 4, then 5, then 1, then 6, then 2
B. 3, then 4, then 5, then 6, then 1, then 2
C. 5, then 4, then 3, then 1, then 6, then 2
D. 6, then 3, then 4, then 5, then 1, then 2
E. 6, then 4, then 3, then 5, then 2, then 1

68. HER2/neu is a test for
A. A membrane receptor
B. A nuclear antigen
C. A cytoplasmic receptor
D. A histone protein
E. A tumor suppressor gene

69. A 26-year-old man presents with a testicular mass and is found to have high serum levels of alpha-fetoprotein (AFP). Microscopic examination of a biopsy from this mass reveals sheets of predominately undifferentiated cells showing focal primitive glandular differentiation. The tumor

cells have large and hyperchromatic nuclei. Which of the following is the most likely diagnosis?

A. Choriocarcinoma
B. Embryonal carcino-ma
C. Mature teratoma
D. Lymphoma
E. Mesothelioma

70. A 28-year-old woman in the second month of pregnancy pre-sents with lower abdominal pain and vaginal bleeding. Histologic examina-tion of products from her spontaneous abortion reveals decidualized tissue and edematous chorionic villi. Chromosomal analysis of the villi show a diploid karyotype. Which of the following is the most likely diagnosis?

A. Yolk sac tumor
B. Papillary adenocarcinoma
C. Hemangioma
D. Teratoma
E. Complete hydatidiform mole

71. Which tumor has the best prognosis in breast carcinoma patients?

A. Invasive ductal carcinoma
B. Invasive lobular carcinoma
C. Intraductal carcinoma
D. Invasive mixed mucinous and ductal carcinoma

72. A 58-year-old man presents with painless enlargement of one testicle. Physical examination reveals a single testicular mass that does not trans-illuminate. The mass is resected. Microscopic examination shows numerous lymphocytes in fibrous stroma between groups of tumor cells having distinct cell membranes and clear cytoplasm. Which of the following is the most likely diagnosis?

A. Seminoma
B. Adenomatoid tumor
C. Teratoma
D. Tuberculosis
E. Leydig cell tumor

73. A 25-year-old woman in the second month of pregnancy pre-sents with lower abdominal pain and vaginal bleeding. Histologic examina-tion of products from her spontaneous abortion reveals decidualized tissue and edematous chorionic villi. Chromosomal analysis of the villi show a triploid karyotype. Which of the following is the most likely diagnosis?

A. Choriocarcinoma
B. Adenocarcinoma
C. Hemangioma
D. Partial hydatidiform mole
E. Teratoma

74. Which has not been found to be predictive of outcome in breast carcinoma patients?

A. Perineural invasion
B. Lymph node status
C. Tumor size
D. Tumor grade

E. Axillary lymph node status

75. A 44-year-old man presents with painless enlargement of one testicle. Physical examination reveals a solitary testicular mass that does not trans-illuminate. The mass is resected, examined histologically, and radiation ther-apy is subsequently given based on the pathologist's diagnosis. Which of the following best describes the expected microscopic appearance of this tumor?
    A. A mixture of malignant cytotrophoblasts and syncytiotrophoblasts
    B. Abnormal tissue derived from all three germ levels with scattered immature neural elements
    C. Large tumor cells with abundant eosinophilic, granular cytoplasm, and rare intracytoplasmic rhomboid crystals
    D. Numerous lymphocytes in the fibrous stroma between groups of tumor cells having distinct cell membranes and clear cytoplasm
    E. Sheets of undifferentiated tumor cells having focal glandular differentiation
76. A 54-year-old woman presents with intermittent pelvic pain. Physical examination reveals a 3-cm mass in the area of her right ovary. Histologic sections from this ovarian mass reveal a papillary tumor with scattered small, round, laminated calcifications. Which of the following is the most likely diagnosis?
    A. Carcinoid
    B. Teratoma
    C. Yolk sac tumor
    D. Serous adenocarcinoma
    E. Dysgerminoma
77. The most common form of invasive breast carcinoma is:
    A. Lobular carcinoma
    B. Mucinous carcinoma
    C. Medullary carcinoma
    D. Ductal carcinoma (or carcinoma of no special type)
    E. Papillary carcinoma
78. The photomicrograph is of a section from a testis removed from the inguinal region of a 28-year-old man. Which of the following statements about this condition is false?
    A. It is unilateral in the majority of cases
    B. Seminoma is the most common malignancy to arise
    C. The condition illustrated is cryptorchidism
    D. There is increased risk of malignancy in the contralateral testis
    E. Both Leydig and Sertoli cells are reduced in number
79. A 45-year-old woman presents with intermittent pelvic pain. Physical examination reveals a 3-cm mass in the area of

her right ovary. Histologic sections from this ovarian mass reveal a papillary tumor with scattered small, round, laminated calcifications. Which of the following is the most likely diagnosis?

A. Granulosa cell tumor
B. Teratoma
C. Serous adenocarcinoma
D. Seminoma
E. Yolk sac tumor

80. Which is the best prognostic indicator in breast carcinoma patients?

A. Tumor size
B. Axillary lymph node status
C. HER2/neu staining of tumor cells
D. Grading of tumor cells
E. Age of the patient

81. This photomicrograph is of a section from a testis removed from the inguinal region of a 25-year-old man. Which of the following statements about this condition is true?

A. It is bilateral in the majority of cases
B. Teratoma is the most common malignancy to arise
C. Risk of associated malignancy is reduced by orchiopexy
D. There is increased risk of malignancy in this testis
E. Both Leydig and Sertoli cells are reduced in number

答案

中文题

1.B	2.E	3.C	4.E	5.D	6.A	7.A	8.D	9.A	10.E
11.B	12.B	13.D	14.E	15.E	16.E	17.D	18.E	19.B	20.B
21.C	22.E	23.A	24.D	25.A	26.A	27.C	28.A	29.C	30.C
31.E	32.D	33.E	34.A	35.B	36.D	37.C	38.B	39.E	40.A
41.A	42.B	43.B	44.E	45.E	46.D	47.C	48.D	49.D	50.E
51.D	52.D	53.A	54.A	55.D	56.B	57.E	58.D	59.D	60.E
61.D	62.C	63.E	64.C	65.C	66.C	67.C	68.E	69.E	70.A
71.B	72.D	73.B	74.C	75.D	76.C	77.B	78.B	79.A	80.B
81.B	82.C	83.B	84.C	85.D	86.E	87.E	88.D	89.B	90.A
91.A	92.E	93.B	94.C	95.E	96.A	97.B	98.D	99.A	100.B
101.E	102.B	103.D	104.B	105.A	106.C	107.D	108.A	109.C	110.C
111.D	112.C	113.E	114.A	115.B	116.A	117.B	118.C	119.E	120.E
121.C	122.B	123.A	124.D	125.D	126.E	127.E	128.E	129.A	130.A
131.A	132.D	133.E	134.A	135.C	136.A	137.D	138.A	139.B	140.A
141.E	142.B	143.B	144.A	145.B	146.C	147.A	148.B	149.B	150.D
151.B	152.A	153.C	154.E	155.A	156.A	157.A	158.D	159.E	160.B

英文题

1.C	2.C	3.B	4.A	5.A	6.C	7.C	8.D	9.B	10.B
11.D	12.B	13.E	14.A	15.B	16.E	17.A	18.B	19.C	20.C
21.B	22.B	23.B	24.C	25.B	26.B	27.D	28.E	29.A	30.A
31.C	32.E	33.D	34.A	35.B	36.B	37.D	38.C	39.A	40.E
41.E	42.B	43.E	44.B	45.E	46.D	47.B	48.E	49.D	50.B
51.C	52.C	53.E	54.E	55.A	56.B	57.D	58.C	59.C	60.C
61.D	62.A	63.B	64.D	65.B	66.A	67.D	68.A	69.B	70.E
71.C	72.A	73.D	74.A	75.D	76.D	77.D	78.E	79.C	80.B
81.D									

10 淋巴造血系统疾病

中文题

1. 霍奇金淋巴瘤最常发生的部位是
   A. 腹膜后淋巴结
   B. 腹股沟淋巴结
   C. 颈部淋巴结
   D. 纵隔淋巴结
   E. 腋下淋巴结
2. III 期霍奇金淋巴瘤的病变范围为
   A. 限于一组淋巴结
   B. 病变限于隔的一侧
   C. 隔肌两侧的淋巴结、脾及邻近器官均累及
   D. 限于一个结外器官
   E. 扩散到淋巴结外，累及一个或多个结外器官或组织
3. 经典霍奇金淋巴瘤预后最好的组织学类型是
   A. 淋巴细胞为主型
   B. 结节硬化型
   C. 混合细胞型
   D. 淋巴细胞削减型
   E. 结节型淋巴细胞为主型
4. 霍奇金淋巴瘤伴有明显纤维条索增生的组织学类型是
   A. 淋巴细胞为主型
   B. 结节硬化型
   C. 混合细胞型
   D. 淋巴细胞削减型
   E. 结节型淋巴细胞为主型
5. 霍奇金淋巴瘤伴有大量淋巴细胞弥漫增生的组织学类型是
   A. 淋巴细胞为主型
   B. 结节硬化型
   C. 混合细胞型
   D. 淋巴细胞削减型
   E. 结节型淋巴细胞为主型
6. 淋巴细胞显著减少的霍奇金淋巴瘤的组织学类型是
   A. 淋巴细胞为主型
   B. 结节硬化型
   C. 混合细胞型
   D. 淋巴细胞削减型
   E. 结节型淋巴细胞为主型
7. 霍奇金淋巴瘤伴有多种细胞混合增生、多数典型 R-S 细胞的组织学类型是
   A. 淋巴细胞为主型
   B. 结节硬化型
   C. 混合细胞型
   D. 淋巴细胞削减型
   E. 结节型淋巴细胞为主型
8. 霍奇金淋巴瘤伴有结节性淋巴细胞增生，肿瘤细胞为爆米花细胞的组织学类型是
   A. 淋巴细胞为主型
   B. 结节硬化型
   C. 混合细胞型
   D. 淋巴细胞削减型
   E. 结节型淋巴细胞为主型
9. 可演化为 B 细胞淋巴瘤的霍奇金淋巴

瘤组织学类型是

A. 淋巴细胞为主型
B. 结节硬化型
C. 混合细胞型
D. 淋巴细胞削减型
E. 结节型淋巴细胞为主型

10. 符合霍奇金淋巴瘤预后好坏的排列顺序为

A. 混合细胞型–淋巴细胞消减型–结节硬化型–淋巴细胞为主型
B. 淋巴细胞消减型–混合细胞型–结节硬化型–淋巴细胞为主型
C. 淋巴细胞为主型–结节硬化型–淋巴细胞消减型–混合细胞型
D. 淋巴细胞为主型–结节硬化型–混合细胞型–淋巴细胞消减型
E. 淋巴细胞消减型–淋巴细胞为主型–结节硬化型–混合细胞型

11. 预后最差的霍奇金病是

A. 结节硬化型
B. 淋巴细胞为主型
C. 淋巴细胞消减型
D. 混合细胞型
E. 结节型淋巴细胞为主型

12. 霍奇金淋巴瘤的具有诊断意义的细胞是

A. 多形性细胞
B. 霍奇金细胞
C. R-S 细胞
D. 陷窝细胞
E. 爆米花细胞

13. 霍奇金淋巴瘤中陷窝细胞多出现在

A. 淋巴细胞为主型
B. 结节硬化型
C. 混合细胞型
D. 淋巴细胞消减型
E. 结节型淋巴细胞为主型

14. 霍奇金淋巴瘤中的非肿瘤细胞成分是

A. 霍奇金细胞
B. 组织细胞
C. 陷窝细胞
D. R-S 细胞
E. 多形性细胞

15. R-S 细胞的组织起源是

A. B 细胞
B. T 细胞
C. 单核细胞
D. NK 细胞
E. 来源未确定

16. II 期霍奇金淋巴瘤的病变范围是

A. 限于一个淋巴结
B. 病变限于膈的一侧
C. 膈肌两侧的淋巴结、脾及邻近器官均累及
D. 限于结外一个器官
E. 弥漫扩散到淋巴结外，累及一个或多个结外器官或组织

17. I 期霍奇金淋巴瘤的病变范围是

A. 限于一个淋巴结或限于结外一个器官
B. 病变限于膈的一侧
C. 膈肌两侧的淋巴结、脾及邻近器官均累及
D. 弥漫扩散到淋巴结外，累及一个或

多个结外器官或组织

E. 以上都不是

18. IV 期霍奇金淋巴瘤的病变范围是

A. 限于一个淋巴结或限于结外一个器官

B. 病变限于膈的一侧

C. 膈肌两侧的淋巴结、脾及邻近器官均累及

D. 弥漫扩散到淋巴结外，累及一个或多个结外器官或组织

E. 以上都不是

19. 符合霍奇金淋巴瘤的描述是

A. 淋巴细胞为主型预后最差

B. 可能与 EB 病毒感染有关

C. I 期肿瘤预后差

D. 淋巴细胞消减型预后最好

E. 陷窝细胞具有诊断意义

20. 非霍奇金淋巴瘤多发生于

A. 纵隔淋巴结

B. 肠系膜淋巴结

C. 腹膜后淋巴结

D. 颈部淋巴结

E. 腋下淋巴结

21. 非霍奇金淋巴瘤工作分类中，高度恶性的是

A. 裂细胞性

B. 小无裂细胞性

C. 大细胞性

D. 小淋巴细胞性

E. 小裂和大裂细胞

22. Lukes-CoIIins 的淋巴瘤分类依据是

A. 瘤细胞形态

B. 肿瘤累及范围

C. 肿瘤细胞的免疫功能

D. 肿瘤细胞对化疗的敏感性

E. 肿瘤细胞对放疗的敏感性

23. 非霍奇金淋巴瘤中最常见的是

A. T 细胞来源

B. B 细胞来源

C. NK 细胞来源

D. 组织细胞来源

E. Langerhans 细胞来源

24. 符合非霍奇金淋巴瘤的描述是

A. 大多数为 T 细胞来源

B. 不引起免疫功能紊乱

C. 多为多克隆起源

D. 确诊时常有播散

E. 多需活体组织检查确诊

25. 符合非霍奇金淋巴瘤的描述是

A. 大多数为 T 细胞来源

B. 不引起免疫功能紊乱

C. 多为单克隆起源

D. 确诊时常有播散

E. 均为高度恶性

26. B 细胞及其肿瘤的表型特征是

A. CD2、CD3、CD4、CD7、CD8

B. CD10、CD19、CD20

C. CD16、CD56

D. CD13、CD14、CD15

E. CD34

27. T 细胞及其肿瘤的表型特征是

A. CD2、CD3、CD4、CD7、CD8

B. CD10、CD19、CD20

C. CD16、CD56

D. CD13、CD14、CD15
E. CD34

28. 低度恶性的非霍奇金淋巴瘤是
A. 滤泡性淋巴瘤
B. 弥漫性大细胞性淋巴瘤
C. 淋巴母细胞性淋巴瘤
D. Burkitt 淋巴瘤
E. 免疫母细胞性淋巴瘤

29. 低度恶性的非霍奇金淋巴瘤是
A. 小细胞性淋巴瘤
B. 弥漫性大细胞性淋巴瘤
C. 淋巴母细胞性淋巴瘤
D. Burkitt 淋巴瘤
E. 免疫母细胞性淋巴瘤

30. 不符合滤泡性淋巴瘤特征的是
A. 肿瘤细胞形成滤泡样结构
B. 由中心细胞构成
C. 由中心母细胞构成
D. 有 t(14;18) 引起 bcl-2 过表达
E. 由 T 细胞起源

31. 符合小淋巴细胞性淋巴瘤的描述是
A. 大部分肿瘤细胞为较成熟的淋巴细胞
B. 高度恶性
C. 形成滤泡结构
D. 与慢性淋巴细胞性白血病无重叠
E. 不引起肝脾肿大

32. 符合弥漫型大细胞性 B 细胞淋巴瘤的描述是
A. 肿瘤细胞呈中心母细胞样或免疫母细胞样
B. 低度恶性
C. 多表达 CD2、CD3
D. 多见于青年患者
E. 对综合化疗不敏感

33. 不符合 Burkitt 淋巴瘤的描述是
A. 儿童多见
B. 多累及颈部淋巴结
C. B 细胞来源
D. 与 EB 病毒感染有关
E. 肿瘤细胞核分裂多见、伴有星天图像

34. 不符合蕈样霉菌病的描述是
A. 原发于皮肤
B. T 细胞来源
C. 形成红斑丘疹
D. 肿瘤细胞核高度扭曲
E. 不累及内脏

35. 儿童多见的非霍奇金淋巴瘤是
A. 小细胞性淋巴瘤
B. 弥漫性大性 B 细胞淋巴瘤
C. 滤泡型淋巴瘤
D. Burkitt 淋巴瘤
E. T 免疫母细胞性淋巴瘤

36. T 细胞淋巴瘤多见于
A. 颌骨
B. 颈部淋巴结
C. 胃肠道
D. 皮肤、鼻咽部
E. 腹膜后淋巴结

37. NK/T 细胞淋巴瘤的特点是
A. 与 EB 病毒感染无关
B. 多累及颈部淋巴结
C. 多形性明显、核不规则的肿瘤细胞

侵犯血管壁，使血管呈洋葱皮状

D. 放射治疗效果差

E. 坏死不明显

38. 不符合白血病的描述是

A. 造血干细胞的恶性肿瘤

B. 患者易发生贫血

C. 急性白血病增生的细胞是较成熟的白细胞

D. 患者易发生出血

E. 患者易发生感染

39. 白血病是

A. 造血干细胞的恶性肿瘤

B. B 细胞的恶性肿瘤

C. T 细胞的恶性肿瘤

D. 组织细胞的恶性肿瘤

E. NK 细胞的恶性肿瘤

40. 白血病在儿童恶性肿瘤中占据

A. 第一位

B. 第二位

C. 第三位

D. 第四位

E. 第五位

41. 不符合白血病的描述是

A. 肝肿大

B. 脾肿大

C. 淋巴结肿大

D. 外周血白细胞均增多

E. 贫血

42. 与白血病的发生无关的因素是

A. 人类 T 细胞病毒

B. 电离辐射

C. 氯霉素、保泰松

D. 苯

E. 3,4-苯并芘

43. 与白血病的发生无关的疾病是

A. Down 综合征

B. Fanconi 综合征

C. Bloom 综合征

D. Bruton 综合征

E. 慢性肉芽肿病

44. 与白血病发生无关的是

A. 放射治疗

B. 烷化剂类药物治疗

C. 细胞毒药物治疗

D. 介入化疗

E. 冷冻治疗

45. 不符合白血病的描述是

A. 慢性白血病急性变以淋巴细胞性白血病多见

B. 髓性白血病比淋巴细胞性白血病多见

C. 急性白血病比慢性白血病多见

D. 白细胞增多性白血病多见

E. 急性白血病儿童和青少年患者多见

46. 急性白血病的特征病变是

A. 外周血白细胞数目增高或减少

B. 淋巴结肿大

C. 脾肿大

D. 骨髓内原始和幼稚细胞大量堆积

E. 肝肿大

47. 不符合急性白血病的描述是

A. 起病急

B. 可转变为慢性白血病

C. 骨髓中原始细胞超过 30%

D. 可由慢性白血病急变而来
E. 外周血中白细胞可低于正常

48. 急性粒细胞性白血病时白血病细胞在肝内的浸润部位是
A. 肝被膜下
B. 集中在小叶中央
C. 集中在汇管区
D. 散在于肝小叶内
E. 围绕在小叶下静脉周围

49. 急性粒细胞性白血病时，瘤细胞在骨髓外浸润，聚集成肿块，称为
A. 棕色瘤
B. 黄色瘤
C. 绿色瘤
D. 黑色素瘤
E. 淋巴瘤

50. 不符合急性白血病的骨髓组织的描述的是
A. 主要为原始细胞
B. 巨核细胞增多
C. 红细胞系生成受抑制
D. 成熟的白细胞数量减少
E. 骨髓增生活跃

51. 不符合急性白血病的淋巴结变化的是
A. 急性淋巴细胞性白血病淋巴结肿大明显
B. 急性粒细胞性白血病淋巴结肿大较轻
C. 淋巴结结构破坏
D. 淋巴结内可见大量原始和幼稚细胞
E. 被膜及周围脂肪无肿瘤细胞浸润

52. 不符合急性淋巴细胞性白血病的描述是
A. 多见于儿童
B. T 细胞性多见
C. 根据 FAB 分类分为 L1、L2、L3 三种类型
D. L1 型分化最差
E. 无标记型少见

53. 不符合急性粒细胞性白血病的描述是
A. 又称为急性髓性白血病
B. 也称为急性非淋巴细胞性白血病
C. 根据 FAB 分类分为 M1～M7 七种类型
D. M1～M3 型均为粒细胞分化的白血病
E. M7 为单核细胞分化的白血病

54. 急性白血病的骨髓组织的描述中，哪一项是不正确的？
A. 主要为原始细胞
B. 巨核细胞增多
C. 红细胞系生成受抑制
D. 成熟的白细胞数量不多
E. 骨髓增生活跃

55. 淋巴细胞性白血病时白血病细胞在肝内的浸润部位是
A. 肝被膜下
B. 集中在小叶中央
C. 集中在汇管区
D. 散在于肝小叶内
E. 围绕在小叶下静脉周围

56. 脾脏在哪种白血病时肿大得最显著？
A. 急性粒细胞性白血病
B. 慢性粒细胞性白血病

C. 急性淋巴细胞性白血病
D. 慢性淋巴细胞性白血病
E. 毛细胞性白血病

57. 下列哪一项符合白血病?
A. 慢性白血病多见
B. 急性白血病可转变为慢性白血病
C. 急性白血病骨髓中原始细胞超过30%
D. 慢性淋巴细胞性白血病容易发生急变
E. 急性白血病外周血中白细胞通常低于正常

58. 不符合慢性白血病的描述是
A. 起病缓慢
B. 病程长
C. 脾肿大
D. 骨髓内原始造血细胞增生
E. 淋巴结肿大

59. 与慢性淋巴细胞白血病重叠的淋巴瘤是
A. 滤泡性型淋巴瘤
B. 弥漫型小细胞性淋巴瘤
C. 弥漫性大细胞性 B 细胞淋巴瘤
D. 蕈样霉菌病
E. Burkitt 淋巴瘤

60. 不符合慢性淋巴细胞性白血病的淋巴结的病理变化的是
A. 瘤细胞浸润包膜
B. 淋巴结结构破坏
C. 淋巴结内原始淋巴细胞大量增生
D. 淋巴结切面呈鱼肉状
E. 肿大淋巴结常互相融合

61. Ph1 染色体出现在
A. 急性淋巴细胞性白血病
B. 慢性淋巴细胞性白血病
C. 急性粒细胞性白血病
D. 慢性粒细胞性白血病
E. 毛细胞性白血病

62. 绝大多数慢性淋巴细胞性白血病来源于
A. B 细胞
B. NK 细胞
C. T4 细胞
D. T8 细胞
E. 无标记细胞

63. 慢性淋巴细胞性白血病外周血增生的细胞是
A. 成熟的小淋巴细胞
B. 中、晚幼粒细胞
C. 原始和幼稚粒细胞
D. 原始和幼稚淋巴细胞
E. 未分化的原巨核细胞

64. 慢性粒细胞性白血病外周血增生的细胞是
A. 成熟的小淋巴细胞
B. 中、晚幼粒细胞
C. 原始和幼稚粒细胞
D. 原始和幼稚淋巴细胞
E. 未分化的原巨核细胞

65. 急性淋巴细胞性白血病外周血增生的细胞是
A. 成熟的小淋巴细胞
B. 中、晚幼粒细胞
C. 原始和幼稚粒细胞

D. 原始和幼稚淋巴细胞
E. 未分化的原巨核细胞

66. 哪项不符合慢性淋巴细胞性白血病的淋巴结的病理变化？
A. 瘤细胞为原始幼稚淋巴细胞
B. 淋巴结结构破坏
C. 淋巴结肿大
D. 淋巴结切面呈鱼肉状
E. 淋巴结常互相融合

67. 慢性粒细胞性白血病与类白血病反应的区别是
A. 发热
B. 中性粒细胞碱性磷酸酶缺如或活性降低
C. 外周血白细胞数目增高
D. 出现幼稚粒细胞
E. 贫血

68. 不符合类白血病反应的描述是
A. 病因消除后血象可恢复正常
B. 常无明显贫血
C. 粒细胞出现毒性颗粒和空泡
D. 中性粒细胞碱性磷酸酶活性增高
E. 出现 Ph1 染色体

69. 慢性粒细胞性白血病的瘤细胞浸润于肝脏的
A. 肝小叶
B. 中央静脉周围
C. 包膜下
D. 汇管区
E. 以上都不是

70. 慢性淋巴细胞性白血病的瘤细胞浸润于肝脏的
A. 肝小叶
B. 中央静脉周围
C. 包膜下
D. 汇管区及其周围
E. 以上都不是

71. 慢性淋巴细胞性白血病主要累及脾的
A. 脾小动脉
B. 脾门
C. 脾包膜
D. 白髓
E. 髓索

72. 不符合慢性白血病的描述是
A. 发展缓慢，病程长
B. 开始化疗效果好
C. 慢性粒细胞白血病晚期易发生急性变
D. 慢性淋巴细胞性白血病较少发生急性变
E. 慢性粒细胞白血病发病时症状明显

73. 不符合慢性粒细胞性白血病急性变的描述是
A. 外周血中、晚幼粒细胞增多
B. 突然高热
C. 贫血加重
D. 出血加剧
E. 骨关节疼痛

74. 慢性淋巴细胞性白血病多见于
A. 婴幼儿
B. 儿童
C. 青年
D. 中年
E. 老年

75. 毛细胞性白血病的特点是
A. 具有 NK 细胞标记
B. 胞浆含溶菌酶阳性
C. 具有 T 细胞标记
D. 有树突状细胞特点
E. 胞浆形成细长突起
76. 毛细胞性白血病最常见的症状是
A. 贫血
B. 外周血白细胞增多
C. 发热
D. 脾肿大
E. 淋巴结肿大
77. 不符合毛细胞性白血病的描述是
A. 多见于男性老年患者
B. 起病缓慢
C. 贫血
D. 脾肿大
E. 常表现为外周血白细胞显著增高
78. 白血病患者贫血主要是由于
A. 红细胞生成减少
B. 缺铁
C. 药物中毒
D. 溶血
E. 出血
79. 白血病患者出血主要是由于
A. 血小板功能障碍
B. 血小板生成减少
C. 凝血因子异常
D. 抗凝血因子异常
E. 药物中毒
80. 白血病患者发热的主要原因是
A. 感染
B. 不明原因
C. 创伤
D. 坏死
E. 变态反应
81. 急性白血病最常见的死亡原因是
A. 感染
B. 出血、特别是颅内出血
C. 贫血
D. 免疫功能低下
E. 疼痛
82. 关于恶性组织细胞增生症，下列哪一项是错误的？
A. 主要侵犯淋巴结、肝、脾、骨髓
B. 一般至少有两个以上不同器官和组织同时受累
C. 脾内瘤细胞主要浸润于白髓
D. 淋巴结内瘤细胞浸润于淋巴窦
E. 肿瘤细胞有吞噬红细胞现象
83. 组织细胞增生症 X 时增生的细胞是
A. 淋巴细胞
B. 单核细胞
C. Langerhans 细胞
D. 巨噬细胞
E. 浆细胞
84. 不符合 Letteret-Siwe 病的描述是
A. 多见于 3 岁以下儿童
B. 发病急、进展快
C. 肝、脾、淋巴结肿大
D. 增生的 Langerhans 细胞分化差、异型性显著
E. 骨组织内形成局限性病灶
85. 不符合 Hand-Schüller-Christian 病的

描述是

A. 多见于幼儿

B. 病变广泛

C. 骨组织病变突出

D. 多累及颅骨和颌骨

E. 增生的 Langerhans 细胞分化差、异型性显著

86. 不符合嗜酸性肉芽肿的描述是

A. 多见于青少年

B. 病变局限于骨

C. 病灶多发

D. 一般不累及内脏

E. 病变内大量 Langerhans 细胞增生，伴嗜酸粒细胞浸润

87. 多发性骨髓瘤是

A. 淋巴细胞的恶性肿瘤

B. 单核细胞的恶性肿瘤

C. Langerhans 细胞的恶性肿瘤

D. 粒细胞的恶性肿瘤

E. 浆细胞的恶性肿瘤

88. 不符合多发性骨髓瘤的描述是

A. 多发性溶骨性破坏

B. 浆细胞的多克隆增生

C. 肿瘤细胞产生 M 蛋白

D. 患者血液中 M 蛋白增高

E. 多数患者尿中 Bence-Jones 蛋白阳性

89. 不符合多发性骨髓瘤的描述是

A. 多发性溶骨性破坏

B. 浆细胞呈单克隆增生

C. 肿瘤细胞产生 M 蛋白

D. 患者血液中 M 蛋白增高

E. 患者尿中 Bence-Jones 蛋白为重链

90. 不符合多发性骨髓瘤的描述是

A. 多发性溶骨性破坏

B. 病变内大量成熟的浆细胞增生

C. 半数患者引起骨髓瘤肾病

D. 可引起贫血、白细胞减少和出血倾向

E. 感染和肾功能衰竭是主要死亡原因

英文题

1. Which of the following diseases is derived from Langerhans cells?

A. Letterer-Siwe disease

B. Eosinophilic granuloma of bone

C. Kikuchi disease

D. A+B

E. A+C

2. Which of the following lymphomas is a mature T-cell lymphoma?

A. T lymphoblastic leukemia/lymphoma

B. Mycosis fungoides

C. Mantle cell lymphoma

D. Follicular lymphoma

E. Burkitt lymphoma

3. Which of the following in correct regarding peripheral T-cell lymphoma, NOS?

A. Association with EBV infection

B. Indolent clinical course

C. It is an immature T-cell lymphoma

D. TCR genes are clonally rearranged in

most cases

E. CD56 expression can be found in most cases

4. The microscopic appearance of a 21-year-old female with a neoplasm of the nasal cavity shows an angiodestructive growth pattern with polymorphic tumor cells. The tumor cells express CD3, CD56, Granzyme B and TIA-1 and EBER1/2 is positive. Which of the following is incorrect regarding this disease?


A. This lymphoma is more prevalent in Asians

B. Association with EBV infection

C. Usually shows an indolent clinical course

D. This is a mature T-cell lymphoma

E. This disease may disseminate to various sites



5. The histological findings shown here at high power microscopically were present in the nasal cavity. The patient is a 48-year-old male with low grade fever, night sweats, fatigue and weight loss. He has a medical history of chronic nasopharyngitis. Which of the following immunophenotypes is most likely?

A. CD3-, CD20+, CD2+

B. CD3+, CD56+, Granzyme B+, TIA-1+

C. CD3+, CD30+, CD20-

D. CD30+, CD15+, LMP-1-

E. CD3+, CD20+, CD68+

6. Which of the following diseases is a T-cell lymphoma?

A. Letterer-Siwe disease

B. Hand-Schuller-Christian disease

C. Histiocytic sarcoma

D. Mycosis fungoides

E. Eosinophilic granuloma

7. The microscopic appearance of a neoplasm the jaw bones of an 8-year-old boy shows a “starry sky” pattern. The tumor cells express CD19, CD20 and CD10. Genetic examination shows t(8;14)(q24;q32). Which of the following is the most likely diagnosis?

A. Burkitt lymphoma

B. Diffuse large B-cell lymphoma

C. Eosinophilic granuloma

D. Mycosis fungoides

E. Follicular lymphoma

8. A 30-year-old female has cervical lymphadenopathy and a low fever for three months. Biopsy of a cervical lymph node shows granuloma formation, composed of hyperplastic histocytes surrounding coag-

ulative necrosis. Ziehl-Neelson staining shows red bacilli. Which of the following is the most likely diagnosis?

A. Histiocytic necrotizing lymphadenitis
B. Castleman disease
C. Cat scratch lymphadenitis
D. Mycosis fungoides
E. Lymphatic tuberculosis

9. Which of the following is incorrect regarding Kikuchi disease?

A. Also known as histiocytic necrotizing lymphadenitis
B. Has a self-limited clinical course
C. C. Shows response to antibiotics
D. Necrosis occurs
E. Shows histocyte hyperplasia

10. Which of the following lymphomas is a primary cutaneous lymphoma characterized by pautrier microabscesses?

A. Extranodal NK/T-cell lymphoma
B. Mycosis fungoides
C. Subcutaneous panniculitis-like T-cell lymphoma
D. Primary cutaneous CD30 positive T-cell lymphoma
E. Primary cutaneous follicle centre lymphoma

11. Which of the following is incorrect regarding mycosis fungoides?

A. Is a primary cutaneous B-cell lymphoma
B. Usually shows an indolent clinical course
C. Can be classified into four clinical stages
D. Extracutaneous dissemination may occur in advanced stages
E. Pautrier microabscesses are a typical finding


12. Which of the following is correct regarding cat scratch lymphadenitis?

A. Associated with G- bacillus infection
B. Associated with a pet, especially a cat
C. Cutaneous changes may occur
D. Granulomas with microabscesses are a typical morphological finding
E. All of above

13. A 10-year-old boy has enlarged right cervical lymph nodes and mediastinal lymph nodes. Granulomas with microabscesses were observed on H&E slides of a right cervical lymph node biopsy. Which of the following is the most likely diagnosis?

A. Lymph node hyperplasia
B. Kikuchi disease
C. Tuberculosis of lymph node
D. Cat scratch lymphadenitis
E. Hodgkin lymphoma

14. This is the microscopic appearance of a neoplasm in the colon found on examination of a 40-year-old man who had abdominal distension for several months. This appearance is most suggestive of which diagnosis?




A. MALT lymphoma
B. Diffuse large B-cell lymphoma
C. Follicular lymphoma
D. Classical Hodgkin lymphoma
E. Mantle cell lymphoma

15. Which of the following regarding follicular lymphoma in incorrect?
A. Originates from marginal zone cells
B. Usually has a nodal pattern
C. It is an indolent neoplasm
D. Can transform into a diffuse large B-cell lymphoma
E. May occur primarily in the skin, GI tract, breast and testis

16. A 23-year-old male has left supraclavicular lymphadenopathy. The histological findings show scattered Hodgkin and Reed-Sternberg cells in a diffuse cellular background consisting of small lymphocytes and there are no neutrophils or eosinophils. The phenotype is CD30+, CD15+, CD45-, EMA-, CD20-. Which of the following is the most likely diagnosis?
A. Lymphocyte-rich classical Hodgkin lymphoma
B. Mixed cellularity classical Hodgkin lymphoma
C. Lymphocyte-depleted classical Hodgkin lymphoma
D. Nodular sclerosing classical Hodgkin lymphoma
E. Nodular lymphocyte predominant Hodgkin lymphoma

17. Diffuse large B-cell lymphoma will not occur in which of the following sites?
A. Cervical lymph nodes
B. Colon
C. C. Spleen
D. Bone marrow
E. None of above

18. A 35-year-old man who is a decorator, has had systematic lymphadenopathy for 6 months. He had a right cervical lymph node biopsy after there was no response to treatment with antibiotics. The histological findings are shown here. The cells express CD10, CD20 and BCL6. The most likely diagnosis is:



A. Follicular lymphoma
B. Burkitt lymphoma

C. Classical Hodgkin lymphoma
D. Diffuse Large B-cell lymphoma
E. Mantle cell lymphoma

19. Which of the following cells proliferates in eosinophilic granuloma?
A. lymphocytes
B. Macrophages
C. Pasmacytes
D. Langerhans cells
E. Monocytes

20. Which of the following factors causes anemia in patients with leukemia?
A. Drug poisoning
B. Decreased production of red blood cells
C. Hemolysis
D. Bleeding
E. Iron deficiency

21. Chronic lymphocytic leukemia mostly occurs in:
A. Infants
B. Kids
C. Youth
D. Midlife
E. The elderly

22. Acute lymphoblastic leukemia usually involves which of the following areas in liver?
A. Subcapsular areas
B. The center of hepatic lobules
C. Portal areas
D. Interspersed among hepatic lobules
E. Around sublobular veins

23. Acute myeloid leukemia usually involves which of the following in the liver?
A. Subcapsular areas
B. The center of hepatic lobules
C. Portal areas
D. Interspersed among hepatic lobules
E. Around sublobular veins

24. Which of the following changes can be found in most cases of acute leukemia?
A. WBCs increased in peripheral blood count
B. Immature and primary cell hyperplasia in bone marrow
C. Lymphadenopathy
D. Hepatomegaly
E. Splenomegaly

25. Which of the following factors is not associated with leukemia?
A. Radiation exposure
B. 3,4-Benzypyrene
C. Human T-cell leukemia virus (HTLV)
D. Chloramphenicol, phenylbutazone
E. Benzene

26. Which of the following phenotypes is found in T cells and T-cell lymphomas?
A. CD16+, CD56+
B. CD10+, CD19+, CD20+
C. CD34+
D. CD13+, CD14+, CD15+
E. CD2+, CD3+, CD4+, CD7+, CD8+

27. Which of the following phenotypes is

found in B cells and B-cell lymphomas?

A. CD16+, CD56+
B. CD13+, CD14+, CD15+
C. CD10+, CD19+, CD20+
D. CD2+, CD3+, CD4+, CD7+, CD8+
E. CD34+

28. Which of the following lymphomas is the most common non-Hodgkin lymphoma?

A. Mature T-cell lymphoma
B. B. Mature B-cell lymphoma
C. Mature NK-cell lymphoma
D. Histiocytic sarcoma
E. Langerhans cell sarcoma

29. Which of the following cells is characteristic for diagnosis of mixed cellularity classical Hodgkin lymphoma?

A. Polymorphical cells
B. Reed-Sternberg cells
C. Hodgkin cells
D. Lacunar cells
E. Popcorn cells

30. Hodgkin lymphoma, stage III by definition must involve which of the following areas?

A. A single lymph node region
B. Two or more lymph node regions on the same side of the diaphragm
C. Lymph node regions or structures on both sides of the diaphragm
D. A single lymphoid structure
E. Bone marrow

31. Classical Hodgkin lymphoma frequently involves which of the following sites?

A. Cervical lymph nodes
B. Mediastinal lymph nodes
C. Retroperitoneal lymph nodes
D. Inguinal lymph nodes
E. Axillary lymph nodes

32. Which of the following phenotypes is characteristic of B cells?

A. CD3
B. CD20
C. CD56
D. CD68
E. CD2

33. Which of the following lymphomas has the best prognosis?


A. Mantle cell lymphoma
B. Extranodal NK/T-cell lymphoma
C. MALT lymphoma
D. Burkitt lymphoma
E. Diffuse large B cell lymphoma

34. Which of the following is the correct order for the prognosis of different types of classical Hodgkin lymphoma?


A. NSCHL > LRCHL > MCCHL > LDCHL
B. LRCHL > NSCHL > MCCHL > LDCHL
C. LRCHL > NSCHL > LDCHL > MCCHL
D. LRCHL > MCCHL > NSCHL > LDCHL
E. MCCHL > LRCHL > NSCHL > LD-

CHL

35. The histological findings shown here at low power(left side) and high power (right side) microscopically were present in a cervical lymph node that was biopsied. The patient is a 33-year-old woman with low grade fever, night sweats, fatigue and non-tender cervical and supraclavicular lymphadenopathy. This appearance is most consistent with:



A. Chronic lymphocytic leukemia/small lymphocytic lymphoma
B. Nodular sclerosis classical Hodgkin lymphoma
C. Burkitt lymphoma
D. Metastatic carcinoma
E. Lymphoid reactive hyperplasis



36. This microscopic appearance of an enlarged, non-tender right cervical lymph node found on physical examination of a 38-year-old man which had been present for several months is most suggestive of a diagnosis of:

A. Chronic lymphocytic leukemia/small lymphocytic lymphoma
B. Metastatic carcinoma
C. Burkitt lymphoma
D. Mixed cellularity classical Hodgkin lymphoma
E. Nodular sclerosis classical Hodgkin lymphoma

37. Which of the following lymphomas may show a nodular growth pattern in lymph nodes?

A. Chronic lymphocytic leukemia/small lymphocytic lymphoma
B. Classical Hodgkin lymphoma
C. Mantle cell lymphoma
D. Follicular lymphoma
E. All of above

38. A diffuse lymphoma composed of atypical, polymorphic large cells and expressing CD20, CD45 (common leukocyte antigen) and CD79a should be classified as:

A. Anaplastic large cell lymphoma
B. Diffuse large B-cell lymphoma
C. Extranodal NK/T-cell lymphoma
D. Mantle cell lymphoma
E. None of the above

39. Which of the following is the typical phenotype of classical Hodgkin lymphoma?

A. CD3, CD30, EMA and ALK-1 positive

B. CD10, CD20 and CD43 positive
C. CD15 positive; CD20, CD30 and CD45 (common leukocyte antigen) negative
D. CD30 and CD15 positive; CD20 and CD45 (common leukocyte antigen) negative
E. CD30, CD20 and CD45 positive; CD15 negative

40. A 25-year-old male presents with an extensive destructive midfacial lesion. On biopsy, coagulative necrosis and an angiodestructive growth pattern can be observed. The immunophenotype is positive for CD3, CD56, Granzyme B, and TIA-1. Which of the following is most likely to be associated with this case?
A. H.pylori infection
B. EBV infection
C. Isolated inguinal lymphadenopathy
D. Sjogren's syndrome
E. Hemophagocytic syndrome

41. A 23-year-old female has cervical lymphadenectasis. CD10, CD19, CD20 and BCL2 immunostains are positive in the biopsy. Which of the following is the most likely diagnosis?
A. Hairy cell leukemia
B. Mantle cell lymphoma
C. Follicular lymphoma
D. Peripheral T-cell lymphoma, NOS
E. Mycosis fungoides

42. A 10-year-old female presents with a mediastinal mass and high white blood cell count. Which of the following is the most likely diagnosis?
A. Burkitt lymphoma
B. T lymphoblastic leukemia/lymphoma
C. Primary mediastinal (thymic) large B-cell lymphoma
D. Follicular lymphoma
E. Extranodal NK/T-cell lymphoma

43. In which of the following lymphomas is the cytogenetic/molecular abnormality t(14;18)(q32;q21) usually detectable?
A. T-cell prolymphocytic leukemia
B. Mantle cell lymphoma
C. Follicular lymphoma
D. Diffuse large B-cell lymphoma
E. Chronic lymphocytic leukaemia/small lymphocytic lymphoma

44. Which of the following lymphomas has the worst prognosis?
A. Mantle cell lymphoma
B. Nodular lymphocyte predominant Hodgkin lymphoma
C. MALT lymphoma
D. Follicular lymphoma
E. Chronic lymphocytic leukemia/small lymphocytic lymphoma

45. Which of the following regarding extranodal NK/T-cell lymphoma is incorrect?
A. Association with Epstein-Barr virus (EBV)

B. The upper aerodigestive tract is most commonly involved
C. Coagulative necrosis is a common finding
D. An angiocentric and angiodestructive growth pattern
E. Polymorphic tumor cells with a nodular growth pattern

46. Histological sections of an enlarged lymph node from a 9-year-old female reveal an increased number of reactive follicles containing germinal centers with proliferating B lymphocytes. Which of the following is the most likely diagnosis?
A. Follicular lymphoma
B. Follicular dysplasia
C. Follicular hyperplasia
D. Germinal center atrophy
E. Germinal center metaplasia

47. A non-tender, enlarged lymph node develops in the neck of a 17-year-old girl. On biopsy, the lymph node shows diffuse effacement of its architecture, with presence of fibrous bands creating a nodular pattern. Numerous lacunar cells and scattered Reed-Sternberg cells are present among a mixed population of lymphocytes, eosinophils, macrophages, and plasma cells. Which of the following is the most likely diagnosis?
A. Lymphocyte-depleted classical Hodgkin lymphoma
B. Lymphocyte-rich classical Hodgkin lymphoma
C. Mixed cellularity classical Hodgkin lymphoma
D. Nodular sclerosing classical Hodgkin lymphoma
E. Nodular lymphocyte predominant Hodgkin lymphoma

48. The presence of the Philadelphia chromosome is associated with a more favorable prognosis in patients with which of the following diseases?
A. Acute lymphoblastic leukemia
B. Acute myeloid leukemia
C. Chronic lymphocytic leukemia
D. Chronic myelogenous leukemia
E. Hairy cell leukemia

49. A patient complains of difficulty breathing through his nose and bone pain in his cheeks, near his nose. Physical examination and CT of the head reveal mass lesions involving the nose, pharynx, and sinuses. CT-guided biopsy demonstrates a non-keratinizing, squamous cell carcinoma. Which of the following disorders is associated with the same oncogenic virus that is the likely cause of this patient's cancer?
A. Adult T-cell leukemia
B. Hepatocellular carcinoma
C. Cervical carcinoma

D. Kaposi's sarcoma
E. Burkitt lymphoma

50. Molecular studies on an abdominal lymph node containing lymphoma demonstrate a t(8;14)(q24;q32) translocation. This is most compatible with which of the following diseases?
A. Multiple myeloma
B. Mantle cell lymphoma
C. Burkitt lymphoma
D. Follicular lymphoma
E. Anaplastic large cell lymphoma

51. Overexpression of BCL2 gene is most associated with which of the following conditions?
A. T lymphoblastic leukemia/lymphoma
B. Follicular lymphoma
C. Nodal marginal zone lymphoma
D. Extranodal NK/T-cell lymphoma, nasal type
E. Burkitt lymphoma

52. Which of following diseases is a T cell lymphoma?
A. Mantle cell lymphoma
B. Follicular lymphoma
C. Plasma cell myeloma
D. Anaplastic large cell lymphoma
E. Chronic lymphocytic leukemia/small lymphocytic lymphoma

答案

中文题

1.C	2.C	3.A	4.B	5.A	6.D	7.C	8.E	9.E	10.D
11.C	12.C	13.B	14.B	15.A	16.B	17.A	18.D	19.B	20.D
21.C	22.C	23.B	24.E	25.C	26.B	27.A	28.A	29.A	30.E
31.A	32.A	33.B	34.E	35.D	36.D	37.C	38.C	39.A	40.A
41.D	42.E	43.E	44.E	45.A	46.D	47.B	48.D	49.C	50.B
51.E	52.B	53.E	54.A	55.C	56.B	57.C	58.D	59.B	60.C
61.D	62.A	63.A	64.B	65.D	66.B	67.B	68.E	69.A	70.D
71.D	72.E	73.A	74.E	75.E	76.D	77.E	78.A	79.B	80.A
81.B	82.C	83.C	84.D	85.E	86.C	87.E	88.B	89.E	90.B

英文题

1.D	2.B	3.D	4.C	5.B	6.D	7.A	8.E	9.C	10.B
11.A	12.E	13.D	14.B	15.A	16.A	17.E	18.D	19.D	20.B
21.E	22.D	23.D	24.B	25.B	26.E	27.C	28.B	29.B	30.C
31.A	32.B	33.C	34.B	35.B	36.D	37.E	38.B	39.D	40.B
41.C	42.B	43.C	44.A	45.E	46.C	47.D	48.D	49.E	50.C
51.B	52.D								

11　内分泌系统疾病

中文题

1. 符合单纯性甲状腺肿的描述是
   A. 男性多于女性
   B. 老年患者多见
   C. 甲状腺多呈结节状肿大
   D. 一般不伴有甲状腺功能异常
   E. 属良性肿瘤
2. 不符合结节性甲状腺肿的描述是
   A. 结节具有完整的包膜
   B. 结节不压迫周围甲状腺组织
   C. 常有纤维组织增生
   D. 常有出血、坏死
   E. 一般不伴有甲状腺功能异常
3. 不符合地方性甲状腺肿的描述是
   A. 多见于内陆山区、半山区
   B. 女性患者多见
   C. 可引起克汀病
   D. 与缺碘有关
   E. 与甲状腺素合成障碍有关
4. 青春期甲状腺肿发生的原因是
   A. 缺碘
   B. 甲状腺素合成障碍
   C. 机体对碘和甲状腺素的需求增加
   D. 药物影响酪氨酸的转化
   E. 碘在肠道吸收障碍
5. 弥漫性胶样甲状腺肿的肉眼特点是
   A. 甲状腺弥漫肿大、光滑、质软
   B. 甲状腺弥漫肿大、多结节状
   C. 甲状腺轻度肿大、质硬
   D. 甲状腺内单一结节形成、质软、边界清楚
   E. 甲状腺内单一结节形成、质硬、边界不清
6. 结节性甲状腺肿的肉眼特点是
   A. 甲状腺弥漫肿大、光滑、质软
   B. 甲状腺弥漫肿大、多结节状
   C. 甲状腺轻度肿大、质硬
   D. 甲状腺内单一结节形成、质软、边界清楚
   E. 甲状腺内单一结节形成、质硬、边界不清
7. 弥漫性胶样甲状腺肿的镜下特点是
   A. 滤泡显著扩大，类胶质潴留
   B. 滤泡上皮增生与复旧混杂
   C. 滤泡上皮增生呈柱状
   D. 多级乳头形成
   E. 肉芽肿形成
8. 结节性甲状腺肿的镜下特点是
   A. 滤泡显著扩大，类胶质潴留
   B. 滤泡上皮增生与复旧混杂，多数结节形成
   C. 滤泡上皮增生呈柱状
   D. 多级乳头形成
   E. 肉芽肿形成
9. 最常见的甲状腺功能亢进的原因是
   A. Graves 病
   B. 毒性结节性甲状腺肿
   C. 毒性甲状腺腺瘤
   D. 甲状腺素摄入过多

E. 垂体腺瘤（分泌 TSH）

10. 不符合 Graves 病的描述是
    A. 又称 Basedow 病
    B. 又称毒性弥漫性甲状腺肿
    C. 是一种自身免疫性疾病
    D. T_3、T_4 分泌过高
    E. 继发于甲状腺炎
11. Graves 病中，引起甲状腺素分泌过高的抗体是
    A. 甲状腺生长免疫球蛋白
    B. 甲状腺生长抑制免疫球蛋白
    C. 甲状腺刺激免疫球蛋白和 TSH 结合抑制免疫球蛋白
    D. 抗甲状腺线粒体抗体
    E. 抗核抗体
12. Graves 病中，引起甲状腺腺上皮增生的抗体是
    A. 甲状腺生长免疫球蛋白和甲状腺生长抑制免疫球蛋白
    B. TSH 结合抑制免疫球蛋白
    C. 甲状腺刺激免疫球蛋白
    D. 抗甲状腺线粒体抗体
    E. 抗核抗体
13. 不符合 Graves 病的临床表现的是
    A. 男性多见
    B. 甲状腺肿大
    C. 代谢率增高
    D. 心悸、多食、多汗、消瘦、易激动
    E. 可伴有眼球突出
14. Graves 病的甲状腺肉眼特点是
    A. 甲状腺弥漫肿大、光滑、质软，伴有血管杂音
    B. 甲状腺弥漫肿大、多结节状
    C. 甲状腺轻度肿大、质硬
    D. 甲状腺内单一结节形成、质软、边界清楚
    E. 甲状腺内单一结节形成、质硬、边界不清
15. Graves 病甲状腺的镜下特点是
    A. 滤泡显著扩大，类胶质潴留
    B. 滤泡上皮增生与复旧混杂，多数结节形成
    C. 滤泡上皮增生呈柱状，伴乳头状突起，类胶质少
    D. 多级乳头形成
    E. 肉芽肿形成
16. Graves 病伴有眼球突出的原因是
    A. 球后神经纤维增生
    B. 球后淤血
    C. 球后毛细血管增生
    D. 球后脂肪纤维组织增生
    E. 球后淋巴水肿
17. 不符合地方性克汀病的描述是
    A. 发生于地方性甲状腺肿的流行区
    B. 患者身材矮小，智力低下
    C. 与近亲婚姻有关
    D. 甲状腺高度萎缩
    E. 表情痴呆
18. 克汀病的发病原因是
    A. 甲状腺生长抑制免疫球蛋白增高
    B. 缺碘
    C. 抗甲状腺抗体产生
    D. 病毒感染
    E. 甲状腺摘除

19. 与粘液水肿无关的甲状腺疾病是
    A. 垂体坏死
    B. 甲状腺全切
    C. 亚急性甲状腺炎
    D. 甲状腺腺瘤
    E. 慢性淋巴细胞性甲状腺炎
20. 符合粘液水肿的描述是
    A. 基础代谢率正常
    B. 组织间隙大量氨基多糖沉积
    C. 声音高亢
    D. 多食
    E. 易激动
21. 克汀病甲状腺病变特点是
    A. 滤泡显著扩大，类胶质潴留
    B. 滤泡上皮增生与复旧混杂，多数结节形成
    C. 滤泡萎缩或缺如
    D. 多级乳头形成
    E. 肉芽肿形成
22. 不符合克汀病的描述是
    A. 头颅较大
    B. 智力正常
    C. 粘液水肿
    D. 身材矮小
    E. 鼻根宽平
23. 不符合克汀病的描述是
    A. 头颅较大
    B. 散发病例也由缺碘引起
    C. 粘液水肿
    D. 身材矮小
    E. 鼻根宽平
24. 不符合粘液水肿的描述是
    A. 基础代谢率低
    B. 组织间隙大量粘液沉积
    C. 声音嘶哑
    D. 胃肠蠕动减慢
    E. 说话、动作、思维迟钝
25. 不符合亚急性甲状腺炎的是
    A. 女性多于男性
    B. 与病毒感染有关
    C. 易转变为慢性甲状腺炎
    D. 形成有巨细胞的肉芽肿
    E. 引起甲状腺肿大
26. 不符合慢性淋巴细胞性甲状腺炎的描述是
    A. 又称桥本甲状腺炎
    B. 为自身免疫性疾病
    C. 引起甲状腺肿大
    D. 大量淋巴细胞浸润
    E. 滤泡上皮增生
27. 不符合慢性纤维性甲状腺炎的描述是
    A. 又称 Riedel 甲状腺肿
    B. 甲状腺不对称肿大
    C. 病变质地硬如木样
    D. 纤维增生，伴玻璃样变
    E. 病变边界清楚
28. 慢性淋巴细胞性甲状腺炎病变的特点是
    A. 玻璃样变纤维增生，呈浸润性生长
    B. 大量淋巴细胞浸润伴滤泡萎缩
    C. 滤泡破坏，形成有巨细胞的肉芽肿
    D. 多发性脓肿形成
    E. 以上都不是
29. 亚急性甲状腺炎的主要病理变化为

A. 玻璃样变纤维增生，呈浸润性生长
B. 大量淋巴细胞浸润伴滤泡萎缩
C. 滤泡破坏，形成有巨细胞的肉芽肿
D. 多发性脓肿形成
E. 以上都不是

30. 慢性纤维性甲状腺炎的病变特点是
A. 玻璃样变纤维增生，呈浸润性生长
B. 大量淋巴细胞浸润伴滤泡萎缩
C. 滤泡破坏，形成有巨细胞的肉芽肿
D. 多发性脓肿形成
E. 以上都不是

31. 急性血源播散性化脓性甲状腺炎的病变特点是
A. 玻璃样变纤维增生，呈浸润性生长
B. 大量淋巴细胞浸润伴滤泡萎缩
C. 滤泡破坏，形成有巨细胞的肉芽肿
D. 多发性脓肿形成
E. 以上都不是

32. 可以恶变为B细胞淋巴瘤的甲状腺炎是
A. 亚急性甲状腺炎
B. 桥本甲状腺炎
C. 纤维性甲状腺炎
D. 急性甲状腺炎
E. 非特异性淋巴细胞性甲状腺炎

33. 不属于甲状腺腺瘤的描述是
A. 多为单发
B. 包膜不完整
C. 可发生囊性变
D. 可发生纤维化
E. 可发生钙化

34. 甲状腺胚胎性腺瘤的特点是
A. 瘤细胞小，排列为条索状
B. 由小滤泡构成
C. 由较正常的滤泡构成
D. 由嗜酸性细胞构成
E. 囊壁被覆单层细胞，并由乳头构成

35. 甲状腺胎儿性腺瘤的特点是
A. 瘤细胞小，排列为条索状
B. 由小滤泡构成
C. 由较正常的滤泡构成
D. 由嗜酸性细胞构成
E. 囊壁被覆单层细胞，并由乳头构成

36. 甲状腺单纯性腺瘤的特点是
A. 瘤细胞小，排列为条索状
B. 由小滤泡构成
C. 由较正常的滤泡构成
D. 由嗜酸性细胞构成
E. 囊壁被覆单层细胞，并由乳头构成

37. 甲状腺嗜酸性细胞腺瘤的特点是
A. 瘤细胞小，排列为条索状
B. 由小滤泡构成
C. 由较正常的滤泡构成
D. 由嗜酸性细胞构成
E. 囊壁被覆单层细胞，并由乳头构成

38. 甲状腺乳头状腺瘤的特点是
A. 瘤细胞小，排列为条索状
B. 由小滤泡构成
C. 由较正常的滤泡构成
D. 由嗜酸性细胞构成
E. 囊壁被覆单层细胞，并由乳头构成

39. 最常见的恶性甲状腺肿瘤是
A. 滤泡性癌
B. 乳头状癌

C. 髓样癌
D. 未分化癌
E. 鳞癌

40. 伴有砂砾体的甲状腺癌是
A. 滤泡性癌
B. 乳头状癌
C. 髓样癌
D. 未分化癌
E. 鳞状细胞癌

41. 分泌降钙素的甲状腺癌是
A. 滤泡性癌
B. 乳头状癌
C. 髓样癌
D. 未分化癌
E. 鳞状细胞癌

42. 恶性度最高的甲状腺癌是
A. 滤泡性癌
B. 乳头状癌
C. 髓样癌
D. 未分化癌
E. 鳞状细胞癌

43. 甲状腺乳头状腺癌的病理组织学特点是
A. 瘤细胞大小不一，可见奇异的巨核细胞
B. 乳头分支多，被覆上皮细胞核呈毛玻璃样
C. 滤泡结构，上皮具有不同程度的异型性，浸润被膜和血管
D. 瘤细胞圆形或多边形，排列成团，分泌降钙素
E. 瘤细胞呈梭形，异型性明显

44. 甲状腺滤泡腺癌的病理组织学特点是
A. 瘤细胞大小不一，可见奇异的巨核细胞
B. 乳头分支多，被覆上皮细胞核呈毛玻璃样
C. 滤泡结构，上皮具有不同程度的异型性，浸润被膜和血管
D. 瘤细胞圆形或多边形，排列成团，分泌降钙素
E. 瘤细胞呈梭形，异型性明显

45. 甲状腺髓样癌的病理组织学特点是
A. 瘤细胞大小不一，可见奇异的巨核细胞
B. 乳头分支多，被覆上皮细胞核呈毛玻璃样
C. 滤泡结构，上皮具有不同程度的异型性，浸润被膜和血管
D. 瘤细胞圆形或多边形，排列成团，分泌降钙素
E. 瘤细胞呈梭形，异型性明显

46. 甲状腺梭形细胞未分化癌的病理组织学特点是
A. 瘤细胞大小不一，可见奇异的巨核细胞
B. 乳头分支多，被覆上皮细胞核呈毛玻璃样
C. 滤泡结构，上皮具有不同程度的异型性，浸润被膜和血管
D. 瘤细胞圆形或多边形，排列成团，分泌降钙素
E. 瘤细胞呈梭形，异型性明显

47. 甲状腺巨细胞型未分化癌的病理组织

学特点是

A. 瘤细胞大小不一，可见奇异的巨核细胞

B. 乳头分支多，被覆上皮细胞核呈毛玻璃样

C. 滤泡结构，上皮具有不同程度的异型性，浸润被膜和血管

D. 瘤细胞圆形或多边形，排列成团，分泌降钙素

E. 瘤细胞呈梭形，异型性明显

48. 不符合甲状腺癌的描述是

A. 乳头状癌恶性度高，易发生血道转移

B. 滤泡腺癌恶性度较高，易发生血道转移

C. 未分化癌恶性度高，易发生血道转移

D. 髓样癌为 C 细胞发生的恶性肿瘤

E. 鳞癌罕见

49. 不符合胰岛素依赖型糖尿病的特点是

A. 患者多为青少年

B. 胰岛无明显病理变化

C. 血中胰岛素明显降低

D. 易合并酮症甚至昏迷

E. 治疗依赖胰岛素

50. 不符合非胰岛素依赖型糖尿病的特点是

A. 患者发病年龄多在 40 岁以上

B. 胰岛 B 细胞明显减少，胰岛数目亦减少

C. 血中胰岛素可以不下降

D. 无自家免疫反应的表现

E. 肥胖与本型发病有重要关系

51. I 型糖尿病的病变特点是

A. 胰岛无炎症改变

B. 胰岛进行性破坏，B 细胞减少

C. 胰岛体积正常

D. 胰岛数目正常

E. 胰岛数量正常

52. II 型糖尿病的病变特点是

A. 胰岛内淋巴细胞浸润

B. 胰岛进行性破坏，B 细胞减少

C. 胰岛细胞无明显减少

D. 胰岛纤维化

E. 胰岛变小

53. 不符合 I 型糖尿病的病因的描述是

A. 与病毒感染有关

B. 与自身免疫有关

C. 与遗传因素有关

D. 与肥胖有关

E. 以上都是

54. 不符合 II 型糖尿病的病因的描述是

A. 与胰岛素作用不足有关

B. 与自身免疫有关

C. 与遗传因素有关

D. 与肥胖有关

E. 与胰岛素抵抗性有关

55. 符合 II 型糖尿病的病因的描述是

A. 与感染有关

B. 与自身免疫有关

C. 与遗传因素有关

D. 与胰岛素分泌减少有关

E. 以上都不是

56. 符合 I 型糖尿病的病因的描述是

A. 与胰岛素作用不足有关
B. 与自身免疫有关
C. 与胰岛素抵抗性有关
D. 与肥胖有关
E. 以上都不是

57. 糖尿病引起的动脉病变是
A. 结节性动脉炎
B. 动脉粥样硬化及细动脉玻璃样变
C. 动脉中层钙化
D. 高安动脉炎
E. 血栓闭塞性脉管炎

58. 不符合糖尿病引起的肾脏病变是
A. 弥漫性肾小球硬化
B. 结节性肾小球硬化
C. 细动脉硬化
D. 脂性肾病
E. 肾盂肾炎

59. 糖尿病引起患者失明的主要原因是
A. 视网膜水肿
B. 视网膜出血
C. 视网膜纤维组织增生
D. 毛细血管基底膜增厚
E. 视网膜剥离

60. 不符合糖尿病引起的神经系统病变的描述是
A. 主要累及外周神经
B. 可引起肢体疼痛
C. 可引起肌肉麻痹
D. 可引起胃肠功能障碍
E. 脑细胞不受累

61. 糖尿病引起的动脉病变特点是
A. 引起动脉壁纤维素样坏死
B. 动脉粥样硬化发病年龄早，进展快
C. 引起动脉中层钙化
D. 累及大、中动脉
E. 引起心、脑血管病

62. 糖尿病引起的最常见的肾脏病变是
A. 肾小球硬化
B. 急性肾盂肾炎
C. 细动脉硬化
D. 脂性肾病
E. 慢性肾盂肾炎

63. 糖尿病引起的具有特异性的血管病变是
A. 血管壁纤维素样坏死
B. 粥瘤形成
C. 毛细血管基底膜增厚
D. 细动脉玻璃样变
E. 静脉血栓形成

64. 不符合糖尿病的描述是
A. 高血糖为主要指标
B. 多食、多饮、多尿、体重减少为主要症状
C. 易发生酮症酸中毒
D. 易发生心、脑血管病
E. 抗感染能力正常

65. APUD 细胞来源于
A. 外胚层
B. 内胚层
C. 间叶组织
D. 神经嵴
E. 以上都不是

66. 不符合 APUD 细胞的描述是
A. 具有摄取氨前提脱羧的功能

B. 分泌多肽激素
C. 胞浆内有神经内分泌颗粒
D. 存在于内分泌腺体内
E. 不存在于皮肤粘膜

67. 不属于 APUD 瘤的是
A. 肺小细胞癌
B. 胃肠道类癌
C. 胃泌素瘤
D. 胰岛细胞瘤
E. 神经纤维瘤

68. 属于 APUD 瘤的是
A. 肺鳞癌
B. 肺腺癌
C. 支气管类癌
D. 肺大细胞癌
E. 细支气管肺泡细胞癌

69. 不符合异位激素肿瘤的描述是
A. 肺癌可产生异位激素
B. 胰腺癌可产生异位激素
C. 肝癌可产生异位激素
D. 肾癌不产生异位激素
E. 肉瘤可产生异位激素

70. 异位激素肿瘤一般不产生
A. ACTH
B. 甲状旁腺素
C. 甲状腺素
D. 降钙素
E. 绒毛膜促性腺激素

71. 不符合多发性内分泌肿瘤的描述是
A. 累及两个以上的内分泌腺体
B. 多有内分泌障碍
C. 无家族遗传性
D. 可分为 MEN-I、II、III 型
E. 可发生肿瘤和增生病变

72. 不属于 APUD 瘤的是
A. 甲状腺未分化癌
B. 嗜铬细胞瘤
C. 胃泌素瘤
D. 胰岛细胞瘤
E. 甲状旁腺腺瘤

73. hypothyroidism
A. 甲状腺功能亢进
B. 甲状腺功能低下
C. 甲状腺肿
D. 甲状腺毒症
E. 糖尿病

74. diadetes mellitus
A. 甲状腺功能亢进
B. 甲状腺功能低下
C. 甲状腺肿
D. 桥本病
E. 糖尿病

75. exophthalmic goiter
A. 甲状腺功能亢进
B. 甲状腺功能低下
C. 突眼性甲状腺肿
D. 单纯性甲状腺肿
E. 糖尿病

76. follicular carcinoma
A. 梭形细胞癌
B. 髓样癌
C. 未分化癌
D. 滤泡癌
E. 乳头状癌

77. Cushing syndrome
    A. 梭形细胞癌
    B. 髓样癌
    C. 未分化癌
    D. 滤泡癌
    E. 乳头状癌
78. adrenal insufficiency
    A. 梭形细胞癌
    B. 髓样癌
    C. 未分化癌
    D. 滤泡癌
    E. 乳头状癌
79. pheochromocytoma
    A. 梭形细胞癌
    B. 髓样癌
    C. 未分化癌
    D. 滤泡癌
    E. 乳头状癌
80. multiple endocrine neoplasia syndrome
    A. 巨人症
    B. 呆小症
    C. 多发性内分泌肿瘤综合征
    D. 急性呼吸窘迫综合征
    E. 继发性免役缺陷综合征
81. pituitary adenoma
    A. 垂体腺癌
    B. 垂体腺瘤
    C. 甲状腺癌
    D. 甲状腺腺瘤
    E. 胰岛细胞瘤
82. hypopituitarism
    A. 甲状腺功能亢进
    B. 甲状腺功能低下
    C. 垂体功能亢进
    D. 垂体功能低下
    E. 尿崩症

英文题

1. A 57-year-old woman presents with difficulty swallowing, drooping eyelids, and double vision. A mass in the anterior mediastinum is found. Biopsies from this mass reveal thymus tissue with scattered reactive lymphoid follicles which contain germinal centers. Which of the following is the most likely diagnosis?
    A. Malignant thymoma
    B. Thymic carcinoma
    C. Thymic hyperplasia
    D. Thymic hypoplasia
    E. Thymic lymphoma
2. A 65-year-old man presents with bone pain and is found to have hypocalcemia and increased parathyroid hormone. Surgical exploration of his neck finds all four of his parathyroid glands to be enlarged. Which of the following disorders is the most likely cause of this patient's enlarged parathyroid glands?
    A. Primary hyperplasia
    B. Parathyroid adenoma
    C. Secondary hyperplasia

D. Parathyroid carcinoma
E. Lung carcinoma

3. A 37-year-old man presents with a single, firm mass within the thyroid gland. Histologic examination of the mass reveals nests of tumor cells separated by broad bands of stroma. The stroma stains positively with Congo red stain. Which of the following is the most likely diagnosis?
A. Follicular carcinoma
B. Papillary carcinoma
C. Squamous cell carcinoma
D. Medullary carcinoma
E. Anaplastic carcinoma

4. A 58-year-old woman presents with a mass in her neck. Physical examination shows a single, nonfunctioning mass within the thyroid. Clinically her serum TSH level is within normal limits. Histologic sections from this mass reveal a single nodule composed of follicles similar to normal thyroid tissue. The nodule is surrounded by a complete fibrous capsule that compresses adjacent normal thyroid tissue. Focal invasion into the capsule is found. Further evaluation of the tumor cells demostrates the presence of the PAX8-PPAR-gamma fusion gene. Which of the following is the most likely diagnosis?
A. Colloid carcinoma
B. Colloid goiter
C. Diffuse nontoxic goiter
D. Follicular adenoma
E. Follicular carcinoma

5. A 35-year-old woman presents with progressive muscle weakness and cold intolerance. Physical examination finds enlargement of her thyroid gland, which is rubbery in consistency. Laboratory evaluation finds decreased serum levels of both triiodothyronine (T_3) and thyroxine (T_4), but serum levels of thyroid-stimulating hormone (TSH) are increased. No thyroid-stimulating immunoglobulins are identified in the serum, but thyroidal peroxidase autoantibodies are present. Which of the following histologic findings is most consistent with a diagnosis of Hashimoto's thyroiditis?
A. Diffuse fibrous deposition between atrophic follicles
B. Follicular cell hyperplasia with scalloping of the colloid
C. Granulomatous inflammation with multinucleated giant cells
D. Lymphoid infiltrate with scattered Hurthle cells
E. Parafollicular hyperplasia with deposition of amyloid

6. Which of the following tumors is not a member of the APUD tumors?
A. Carcinoid tumor
B. Small cell carcinoma of the lung

C. Medullary thyroid carcinoma
D. Papillary thyroid carcinoma
E. Pheochromocytoma

7. Which of the following secretes calcitonin?
A. Follicular carcinoma
B. Papillary carcinoma
C. Medullary carcinoma
D. Squamous cell carcinoma
E. Undifferentiated carcinoma

8. Which of the following thyroid carcinomas is a low grade malignancy with a good prognosis?
A. Follicular carcinoma
B. Papillary carcinoma
C. Medullary carcinoma
D. Squamous cell carcinoma
E. Undifferentiated carcinoma

9. Which of the following is incorrect regarding thyroid papillary carcinoma?
A. Nuclei have an optically clear appearance
B. There is intranuclear inclusion formation
C. Papillary architecture may be absent in some cases
D. Psammoma bodies are often present
E. Tend to metastasize through the bloodstream

10. Which is the following is the most common type of cancer of the thyroid:
A. Follicular carcinoma
B. Papillary carcinoma
C. Medullary carcinoma
D. Squamous cell carcinoma
E. Undifferentiated carcinoma

11. An 8-month-old infant is being evaluated for growth and mental retardation. Physical examination reveals a small infant with dry, rough skin, a protuberant abdomen, periorbital edema, a flattened, broad nose, and a large, protuberant tongue. Which of the following disorders is the most likely cause of this infant's signs and symptoms?
A. Graves' disease
B. Cretinism
C. Toxic multinodular goiter
D. Toxic adenoma
E. Struma ovarii

12. A 25-year-old woman presents with the acute onset of cessation of lactation. She delivered her first child several months ago and has been breast-feeding since then. She reports that she has not menstruated since the delivery. She also says that lately she has been tired and has been "feeling cold" all of the time. Laboratory workup reveals a deficiency of ACTH and other anterior pituitary hormones. Which of the following is the most likely cause of this patient's signs and symptoms?
A. Craniopharyngioma
B. Cushings disease

C. Empty sella syndrome
D. Nonsecretory chromophobe adenoma
E. Sheehan's syndrome

13. A 42-year-old man presents with increasing fatigue and occasional headaches. He states that recently he has had to change his shoe size from 9 to 10, and he also thinks that his hands and jaw are now slightly larger. Physical examination reveals a prominent forehead and lower jaw, enlarged tongue, and large hands and feet. Initial laboratory examination reveals increased serum glucose. Which of the following is the most likely explanation for this constellation of clinical findings?
A. Acromegaly
B. Apoplexy
C. Cretinism
D. Diabetes
E. Gigantism

14. Which kind of thyroiditis is associated with increased risk for the development of B cell lymphoma?
A. Subacute thyroiditis
B. Hashimoto thyroiditis
C. Acute thyroiditis
D. Subacute thyroiditis
E. Riedel thyroiditis

15. What is the major pathological change of subacute thyroiditis?
A. Fibrosis of thyroid tissue
B. Destruction of follicle and granulomatous inflammation
C. Abscess formation
D. Abundant lymphocyte infiltration and follicle atrophy
E. Multinodular enlargement of the thyroid

16. Which of the following is incorrect regarding chronic lymphocytic thyroiditis?
A. It is also called Hashimoto thyroiditis
B. It is caused by autoimmune inflammatory disorder
C. It is associated with enlargement of thyroid
D. There are large numbers mononuclear inflammatory infiltrate
E. There is hyperplasia of follicle epithelium

17. Which of the following is the typical morphological feature of Grave's disease?
A. Multinodular enlargement of the thyroid
B. Colloid-rich follicles lined by flattened, inactive epithelium
C. Regenerative changes are common
D. Diffuse hypertrophy and hyperplasia of follicular epithelial cells with formation of small papillae
E. Destruction of follicles and formation of granulomas

18. Which of the following autoantibodies results in the increased thyroxin synthesis in Grave's disease?

A. Thyroid stimulating immunoglobulin (TSI)
B. Anti-thyroglobulin antibody
C. Thyroid growth stimulating immunoglobulin (TGI)
D. TSH binding inhibitor immunoglobin (TBIIs)
E. Anti-thyroid peroxisomes antibody

19. Which of the following is the cause of puberty sporadic goiter?
A. Lack of iodine associated with geographic region
B. Increased physiologic demand for thyroxin
C. Impairment in the synthesis of thyroxin
D. Hereditary enzyme defect
E. Taking medication that interferes with thyroid hormone synthesis

20. An eosinophilic adenoma of the pituitary gland is not associated with:
A. Enlargement of the sella turcica
B. Hypertrophy and hyperplasia of the soft tissues throughout the body
C. Excessive growth of acral parts
D. Premature closure of the epiphyses
E. Impaired glucose tolerance

21. Which of the following is not likely to occur in patients with pheochromocytoma?
A. Paroxysmal hypertention
B. Sweating
C. Neurofibromatosis
D. A fall in blood pressure on palpating the abdomen
E. Paroxysmal hypotension

22. A 25-year-old woman who has never been pregnant presents with 3 months of amenorrhea and a milky discharge from her nipple. She states that her menstrual cycles have been irregular for the past year. Laboratory tests show that her serum LH and estradiol levels are below normal, and a pregnancy test is negative. Which of the following is the most likely cause of these signs and symptoms?
A. Craniopharyngioma of the hypothalamus
B. Germinoma of the pineal gland
C. Islet cell adenoma of the pancreas
D. Medullary carcinoma of the thyroid gland
E. Prolactinoma of the pituitary gland

23. A 40-year-old male patient presents with an abrupt elevation of blood pressure. A CT scan reveals a mass in his adrenal region. Laboratory examination reveals increased urinary levels of metanephrine and vanillylmandelic acid (VMA). A histologic section from the mass reveals a tumor composed of polygonal and spindle cells with abundant granular cytoplasm, arranged in a nested pattern. Electron microscopy reveals variable numbers of electron-dense granules. Which of the fol-

lowing is the most likely diagnosis?

A. Adrenal cortical carcinoma

B. Ganglioneuroma

C. Nephroblastoma

D. Neuroblastoma

E. Pheochromocytoma

答案

中文题

1.C	2.A	3.E	4.C	5.A	6.B	7.A	8.B	9.A	10.E
11.C	12.A	13.A	14.A	15.C	16.D	17.C	18.B	19.D	20.B
21.C	22.B	23.B	24.B	25.C	26.E	27.E	28.B	29.C	30.A
31.D	32.B	33.B	34.A	35.B	36.C	37.D	38.E	39.B	40.B
41.C	42.D	43.B	44.C	45.D	46.E	47.A	48.A	49.B	50.B
51.B	52.C	53.D	54.B	55.C	56.B	57.B	58.D	59.E	60.E
61.B	62.A	63.C	64.E	65.D	66.E	67.E	68.C	69.D	70.C
71.C	72.A	73.B	74.E	75.C	76.D	77. 删除	78. 删除	79.B	80.C
81.B	82.D								

英文题

1.C	2.C	3.D	4.E	5.E	6.E	7.C	8.B	9.E	10.B
11.B	12.E	13.A	14.B	15.B	16.E	17.D	18.C	19.B	20.D
21.D	22.E	23.E							

12　神经系统疾病

中文题

1. 红色神经元是指
   A. 神经元萎缩
   B. 神经元尼氏小体溶解
   C. 神经元胞体固缩
   D. 神经元水肿
   E. 神经元坏死
2. 神经元胞浆或核内出现包含体的原因是
   A. 病毒感染
   B. 细菌感染
   C. 立克次体感染
   D. 螺旋体感染
   E. 真菌感染
3. Wallerian 变性是指
   A. 神经元变性
   B. 神经元尼氏小体溶解
   C. 髓鞘崩解消失
   D. 轴突断裂后的一系列变化
   E. 神经原纤维缠结
4. 小胶质细胞吞噬变性、坏死神经细胞的现象，被称为
   A. 卫星现象
   B. 嗜神经细胞现象
   C. 胶质瘢痕形成
   D. 包含体形成
   E. 胶质结节形成
5. 神经系统中可转化为格子细胞的是
   A. 神经细胞
   B. 星形胶质细胞
   C. 小胶质细胞
   D. 室管膜细胞
   E. 施万细胞
6. 构成胶质结节的细胞成分是
   A. 神经细胞
   B. 星形胶质细胞
   C. 小胶质细胞
   D. 室管膜细胞
   E. 少突胶质细胞
7. 围绕在变性、坏死神经元形成卫星现象的细胞是
   A. 神经细胞
   B. 星形胶质细胞
   C. 小胶质细胞
   D. 室管膜细胞
   E. 少突胶质细胞
8. 卫星现象是指围绕变性、坏死神经元的少突胶质细胞数大于或等于
   A. 1 个
   B. 2 个
   C. 3 个
   D. 4 个
   E. 5 个
9. 细胞毒性脑水肿的主要镜下改变是
   A. 细胞体积增大、水样肿胀
   B. 细胞和血管周间隙增大
   C. 胶质瘢痕形成
   D. 胶质结节形成
   E. 髓鞘崩解消失

10. 血管源性脑水肿的主要镜下改变是
    A. 细胞体积增大、水样肿胀
    B. 细胞和血管周间隙增大
    C. 胶质瘢痕形成
    D. 胶质结节形成
    E. 髓鞘崩解消失
11. 颅内压增高是指侧卧位时，颅内压持续超过
    A. 0.5 kPa
    B. 1.0 kPa
    C. 1.5 kPa
    D. 2.0 kPa
    E. 2.5 kPa
12. 枕骨大孔疝可以引起
    A. 同侧眼上视和内视障碍
    B. 扣带回受压
    C. 延髓受压
    D. 中脑和脑干受压
    E. 同侧肢体瘫痪
13. 大脑镰下疝可以引起
    A. 同侧眼上视和内视障碍
    B. 扣带回受压
    C. 延髓受压
    D. 中脑和脑干受压
    E. 同侧肢体瘫痪
14. 小脑天幕疝不会引起
    A. 同侧眼上视和内视障碍
    B. 扣带回受压
    C. 枕叶距状裂出血性梗死
    D. 中脑和脑干受压
    E. 同侧肢体瘫痪
15. 交通性脑积水可见于
    A. 脉络丛乳头状瘤
    B. 室间孔阻塞
    C. 室管膜瘤
    D. 流行性脑脊髓膜炎
    E. 蛛网膜下腔出血
16. 对缺血缺氧最敏感的脑内细胞是
    A. 神经元
    B. 星形胶质细胞
    C. 少突胶质细胞
    D. 室管膜细胞
    E. 小胶质细胞
17. 脑血流自动调节机制发生障碍的收缩压是
    A. ＜ 50mmHg
    B. ＜ 60mmHg
    C. ＜ 70mmHg
    D. ＜ 80mmHg
    E. ＜ 90mmHg
18. 缺血缺氧引起脑组织发生病理组织学改变的时间是
    A. ＞ 2h
    B. ＞ 4h
    C. ＞ 6h
    D. ＞ 12h
    E. ＞ 24h
19. 缺血性脑病最常累及的部位是
    A. 血供边缘带
    B. 顶叶
    C. 颞叶
    D. 枕叶
    E. 脑干
20. 引起脑梗死最常见的原因是

A. 脑细小动脉硬化
B. 脑动脉粥样硬化
C. 脑动脉栓塞
D. 先天性脑血管畸形
E. 脑血管创伤

21. 脑栓塞的最常见的原因是
A. 脂肪栓塞
B. 心源性栓子
C. 血液高凝状态
D. 空气栓塞
E. 羊水栓塞

22. 脑出血的最常见原因是
A. 高血压病
B. 血液病
C. 动脉粥样硬化症
D. 动脉瘤
E. 脑动脉瘤

23. 脑出血最常累及的部位是
A. 脑室出血
B. 小脑出血
C. 基底节出血
D. 中脑出血
E. 脑桥出血

24. 原发性蛛网膜下腔出血的最常见原因是
A. 动静脉畸形
B. 先天性囊状动脉瘤
C. 血液病
D. 脑动脉炎
E. 脑底异常血管网

25. 高度恶性的星形胶质细胞瘤是
A. 纤维型星形胶质细胞瘤
B. 肥胖型星形胶质细胞瘤
C. 原浆型星形胶质细胞瘤
D. 间变型星形胶质细胞瘤
E. 多形性胶质母细胞瘤

26. 据世界卫生组织的分类，属于 III 级的星形胶质细胞瘤是
A. 纤维型星形胶质细胞瘤
B. 肥胖型星形胶质细胞瘤
C. 原浆型星形胶质细胞瘤
D. 间变型星形胶质细胞瘤
E. 多形性胶质母细胞瘤

27. 介于良恶性之间的星形胶质细胞瘤是
A. 纤维型星形胶质细胞瘤
B. 肥胖型星形胶质细胞瘤
C. 原浆型星形胶质细胞瘤
D. 间变型星形胶质细胞瘤
E. 多形性胶质母细胞瘤

28. 室管膜细胞瘤最常见于
A. 侧脑室
B. 第三脑室
C. 第四脑室
D. 中脑水管
E. 脊髓腰骶部

29. 髓母细胞瘤最常见于
A. 小脑引部
B. 小脑半球
C. 额叶
D. 顶叶
E. 枕叶

30. 髓母细胞瘤的组织学特点是
A. 瘤细胞呈栅栏状排列
B. 肿瘤出血、坏死明显

C. 瘤细胞形成纤维心菊形团
D. 瘤细胞围绕血管形成菊形团
E. 瘤细胞呈上皮样

31. 脑膜瘤最常见的部位是
A. 上矢状窦两侧
B. 蝶骨底
C. 嗅沟
D. 脑室
E. 椎管

32. 大多数脑膜瘤的起源于
A. 脉络丛上皮细胞
B. 室管膜上皮细胞
C. 蛛网膜颗粒细胞
D. 血管内皮细胞
E. 脑膜纤维母细胞

33. 多发性硬化属于
A. 肉芽肿性炎
B. 继发性免疫缺陷病
C. 原发性免疫缺陷病
D. 自身免疫性疾病
E. 获得性免疫缺陷病

34. 多发性硬化的中枢神经系统主要病变是
A. 白质脱髓鞘
B. 血管周淋巴细胞、浆细胞浸润
C. 病变区淋巴细胞浸润
D. 病变区可见格子细胞
E. 轴突肿胀、断裂

35. Devic 病的脱髓鞘病变特点是
A. 主要累及视神经和脊髓
B. 主要累及脑室周围白质
C. 主要累及脑干白质
D. 病变呈同心圆分布
E. 主要累及小脑白质

36. Balo 病的病变特点是
A. 小静脉周围淋巴细胞为主的炎细胞浸润
B. 脱髓鞘带和正常髓鞘带呈同心圆分层排列
C. 病变区可见格子细胞
D. 轴突肿胀、断裂
E. 病变区淋巴细胞浸润

37. 病情凶险的中枢神经脱髓鞘病是
A. 多发性硬化
B. 同心圆性硬化
C. 视神经脊髓炎
D. 急性播散性脑脊髓炎
E. 急性出血性白质脑炎

38. 急性播散性脑脊髓炎的病变特点是
A. 主要累及视神经和脊髓
B. 主要累及脑室周围白质
C. 主要累及脑干白质
D. CNS 弥漫受累
E. 主要累及小脑白质

39. 急性出血性坏死性脑脊髓炎的病变特点是
A. 脑白质弥漫分布的点状出血
B. 蛛网膜下腔出血
C. 硬脑膜下出血
D. 硬脑膜外出血
E. 基底节出血

40. 有关流行性乙型脑炎的主要病理变化，哪项是错误的？
A. 神经细胞的变性，坏死

B. 筛状软化灶形成
C. 淋巴细胞套袖状浸润
D. 病变以小脑，延髓和桥脑最严重
E. 胶质结节

41. 流行性乙型脑炎的病变，下列哪个部位最为轻微？
A. 大脑皮质
B. 基底节
C. 丘脑
D. 中脑
E. 脊髓

42. 流行性乙型脑炎属于
A. 化脓性炎症
B. 变质性炎症
C. 纤维素性炎症
D. 浆液性炎症
E. 肉芽肿性炎症

43. 流行性乙型脑炎的主要病理变化是
A. 神经细胞变性坏死
B. 蛛网膜下腔积脓
C. 肉芽肿形成
D. 微小动脉瘤形成
E. 脑疝形成

44. 流行性乙型脑炎的主要侵犯部位是
A. 脊髓灰质
B. 延脑
C. 小脑
D. 大脑灰质及神经核团
E. 神经节

45. 符合流行性乙型脑炎的描述是
A. 经呼吸道传播
B. 冬春发病
C. 可发生华-弗综合征
D. 可引起智力低下
E. 克氏症阳性

46. 不符合流行性乙型脑炎的描述是
A. 大脑内有软化灶形成
B. 小血管周淋巴细胞和单核细胞套袖状浸润
C. 大脑神经元变性坏死
D. 蛛网膜下腔大量白细胞
E. 小胶质细胞结节形成

47. 流行性乙型脑炎的病原体属于
A. RNA 病毒
B. 细菌
C. 真菌
D. 支原体
E. DNA 病毒

48. 流行性乙型脑炎的传播途径是
A. 消化道传播
B. 呼吸道传播
C. 经血液传播
D. 性传播
E. 蚊虫传播

49. 流行性乙型脑炎的患者多为
A. 男性
B. 女性
C. 儿童
D. 成人
E. 老人

50. 不符合流行性乙型脑炎的临床表现是
A. 嗜睡
B. 昏迷
C. 角弓反张

D. 头痛
E. 呕吐

51. 不符合流行性乙型脑炎的预后的描述是
A. 可引起痴呆
B. 可引起语言障碍
C. 可引起智力低下
D. 可引起脑积水
E. 可引起肢体瘫痪

52. 有关急性暴发性流行性脑脊髓膜炎的描述，哪项不正确？
A. 起病急，进展快，死亡率高
B. 脑膜病变重
C. 常伴中毒性休克和 DIC
D. 由脑膜炎双球菌释放的内毒素所致
E. 常伴肾上腺皮质出血和功能衰竭

53. 有关暴发性脊髓脑膜炎时的临床病理表现的描述，哪一项是错误的？
A. 皮肤粘膜广泛淤点及淤斑
B. 双侧肾上腺广泛出血
C. 蛛网膜下腔充满脓性渗出物
D. 起病急骤，高热，头痛，呕吐
E. 伴有周围循环衰竭

54. 不符合流行性脑脊髓膜炎的临床表现是
A. 颈强直
B. 脑脊液淋巴细胞增多
C. 角弓反张
D. 头痛
E. 呕吐

55. 不符合流行性脑脊髓膜炎的预后的是
A. 大多可痊愈
B. 可引起脑积水
C. 可引起面神经麻痹
D. 常引起智力低下
E. 可引起耳聋

56. 流行性脑脊髓膜炎的病原体属于
A. RNA 病毒
B. 细菌
C. 真菌
D. 支原体
E. DNA 病毒

57. 流行性脊髓脑膜炎的主要病理变化是
A. 软化灶形成
B. 神经细胞变性坏死
C. 淋巴细胞血管周围套状浸润
D. 蛛网膜下腔积脓
E. 小胶质细胞结节状增生

58. 流行性脑脊髓膜炎属于
A. 变质性炎症
B. 增生性炎症
C. 化脓性炎症
D. 浆液性炎症
E. 肉芽肿性炎症

59. 流行性脑脊髓膜炎主要侵犯
A. 硬脑膜外
B. 硬脑膜
C. 软脑膜
D. 大脑皮质
E. 蛛网膜和软脑膜

60. 导致脑疝的大脑疾病是
A. 流行性乙型脑炎
B. 脑水肿
C. 脑出血

D. 脑内肿瘤

E. 以上都可能

61. 流行性脑脊髓膜炎一般不引起

A. 皮肤、粘膜有出血点和瘀斑

B. 脑软化灶形成

C. 脑疝形成

D. 高热、头痛、呕吐

E. 脑膜刺激症明显

62. 不符合流行性脑脊髓膜炎的描述是

A. 引起脑膜刺激症

B. 引起颅内压升高症状

C. 引起脑脊液浑浊或脓样

D. 引起神经细胞变性坏死

E. 引起皮肤瘀点和瘀斑

63. 流行性脑脊髓膜炎的病变最明显的部位是

A. 脑底部脑膜

B. 脑顶部脑膜

C. 大脑灰质

D. 脊髓灰质前角

E. 大脑白质

64. 小儿麻痹症病变主要累及

A. 脊髓前角运动神经元

B. 脊髓后角感觉神经元

C. 延髓神经核团

D. 脑干神经核团

E. 大脑灰质神经元

65. 脊髓灰质炎的病变最重的部位是

A. 脊髓腰膨大

B. 脊髓颈膨大

C. 延髓

D. 脑干

E. 大脑

66. 脊髓灰质炎最主要的传播途径是

A. 消化道传播

B. 呼吸道传播

C. 经血液传播

D. 性传播

E. 蚊虫传播

67. 脊髓灰质炎的病原体属于

A. RNA 病毒

B. 细菌

C. 真菌

D. 支原体

E. DNA 病毒

68. 脊髓灰质炎病变最轻的部位是

A. 脊髓腰膨大

B. 脊髓颈膨大

C. 延髓

D. 脑桥

E. 大脑皮质

69. 脊髓灰质炎引起下肢麻痹主要是由于

A. 充血水肿

B. 胶质瘢痕形成

C. 小胶质细胞增生

D. 运动神经元变性坏死

E. 中性粒细胞浸润

70. 脊髓灰质炎病毒感染的大多数表现为

A. 隐性感染

B. 轻型

C. 非麻痹型

D. 麻痹型

E. 显性感染

71. 引起死亡的脊髓灰质炎类型是

A. 隐性感染
B. 顿挫型
C. 非麻痹型
D. 脊髓型
E. 延髓型

72. 狂犬病的特征性病变是
A. 小胶质细胞增生
B. 脑水肿
C. 神经细胞胞浆内出现 Negri 小体
D. 神经细胞变性、坏死
E. 血管周淋巴细胞为主的炎细胞浸润

73. Negri 小体内含有的病毒颗粒是
A. 单纯疱疹病毒
B. 乳头瘤病毒
C. 狂犬病病毒
D. 巨细胞病毒
E. 麻疹病毒

74. 引起海绵状脑病的病原体是
A. DNA 病毒
B. RNA 病毒
C. 慢病毒
D. 朊蛋白
E. 逆转录病毒

75. 不符合朊蛋白的描述是
A. PrP^{c} 是神经元的跨膜蛋白
B. PrP^{sc} 是 β 折叠构型的 PrP
C. PrP^{sc} 具有感染性
D. PrP^{sc} 可被蛋白酶消化
E. PrP^{sc} 可被漂白粉消毒

76. 人类 Prion 病的病变特点是
A. 神经细胞固缩
B. 噬神经细胞现象
C. 神经细胞卫星现象
D. 神经细胞红色变性
E. 神经细胞胞浆及神经毡空泡形成

77. Alzheimer 病常发生于
A. 30 岁以后
B. 40 岁以后
C. 50 岁以后
D. 60 岁以后
E. 70 岁以后

78. 老年人最常见的神经变性疾病是
A. Alzheimer 病
B. Pick 病
C. Lewy body 病
D. Parkinson 病
E. 肌萎缩性侧索硬化

79. 不属于 Alzheimer 病的病理变化是
A. 老年斑
B. 神经原纤维缠结
C. Hirano 小体
D. 颗粒空泡变性
E. Lewy 小体

80. 属于 Parkinson 病的病理变化是
A. 老年斑
B. 神经原纤维缠结
C. Hirano 小体
D. 颗粒空泡变性
E. Lewy 小体

81. 痴呆是指
A. 智能障碍
B. 意识障碍
C. 失语症
D. 失用症

E. 失认症

82. 引起神经原纤维缠结的主要原因是

A. 淀粉样物质沉积

B. τ 蛋白过度磷酸化

C. 早老蛋白基因突变

D. 巨球蛋白基因突变

E. Prion 感染

83. 不属于 Alzheimer 病的病理变化是

A. Hirano 小体

B. 神经原纤维缠结

C. 颗粒空泡变性

D. 神经细胞胞浆及神经毡空泡形成

E. 老年斑

84. 老年斑的主要成分是

A. 微丝蛋白

B. 微管蛋白

C. β 淀粉样物质

D. 胶质纤维酸性蛋白

E. S-100 蛋白

85. Parkinson 病脑损害的部位是

A. 大脑皮质

B. 基底节

C. 纹状体黑质

D. 小脑

E. 脊髓

86. Parkinson 病的特征性病理变化是

A. 黑质多巴胺细胞脱失

B. 额颞叶萎缩

C. 壳核神经元脱失

D. 大脑皮质神经元丧失

E. 脊髓前角运动神经元减少

87. 最常见的神经系统真菌感染性疾病是

A. 曲菌

B. 毛霉菌

C. 放线菌

D. 新型隐球菌

E. 白色念珠菌

英文题

1. A 38-year-old man with AIDS presents with decreasing mental status. Cerebrospinal fluid (CSF) is stained with a mucicarmine stain and India ink. The mucicarmine stain reveals numerous yeast that stain bright red. The India ink staining shows that these yeast have a capsule. Which of the following is the most likely diagnosis?

A. Chromomycosis

B. Coccidioidomycosis

C. Cryptococcosis

D. Cryptosporidiosis

E. Paracoccidioidomycosis

2. A 45-year-old woman presents with unilateral tinnitus and unilateral hearing loss. Physical examination reveals facial weakness and loss of corneal reflex on the same side as the tinnitus and hearing loss. Where would a tumor need to be located to produce these clinical signs and symptoms?

A. Anterior horn of the spinal cord
B. Anterior pituitary
C. Cerebellopontine angle
D. Frontal cortex
E. Lateral ventricle

3. A 38-year-old woman presents with severe headaches which show increasing frequency. A CT scan of her head shows a 2-cm mass attached to the dura in her right frontal area. Which of the following histologic changes is most likely to be seen in a biopsy specimen taken from this tumor?
A. Antoni A areas with Verocay bodies
B. A whorled pattern with psammoma bodies
C. Endothelial proliferation with serpentine areas of necrosis
D. Fried-egg appearance of tumor cells without necrosis
E. True rosettes and pseudorosettes

4. Which of the following is the characteristic pathological change of prion disease?
A. Presence of vacuoles within neuropil and cell bodies in gray matter
B. Coagulation necrosis
C. Red neurons
D. Condensation of nuclear material (pyknosis)
E. Loss of nuclear staining (karyolosis)

5. Which of the following tumors displays the most malignant behavior?
A. Meningioma
B. Diffuse astrocytoma
C. Anaplastic astrocytoma
D. Oligodendroglia
E. Glioblastoma multiforme

6. Within the central nervous system, which of the following regions is most susceptible to hypoxic-ischemic injury?
A. Arterial border zones
B. Cerebellum
C. Frontal lobe
D. Temporal lobe
E. Brain stem

7. Brain parenchymal hemorrhages caused by hypertension most commonly occur in which of these sites:maldinisong
A. Basal ganglia
B. Thalamus
C. Cerebellum
D. Pons
E. Cerebral white matter

8. Which of the following disorders is not the cause of communicating hydrocephalus:
A. Papilloma of choroid plexus
B. Obstruction of foramen of Luschka
C. Ependymoma
D. Meningitis
E. Subarachnoid hemorrhage

9. Which of the following descriptions is incorrect regarding tonsillar herniation:
A. A life threatening herniation
B. Brain stem compression
C. Hemorrhagic lesions in the midbrain

and pons

D. Respiratory inhibition

E. Impairment of ocular movement on the same side as the lesion

10. A 55-year-old woman presents with a new onset of seizure activity. Computerized tomography (CT) scans and skull x rays demonstrate a mass in the right cerebral hemisphere that is markedly calcific. The mass is removed and histologic sections reveal sheets of cells with clear halos (“fried-egg” appearance) and scattered calcifications. What type of tumor is does this woman have?

A. Oligodendroglima

B. Meningioma

C. Astrocytoma

D. Glioblastoma multiforme

E. Medullobalstoma

11. A 65-year-old man presents with tremors at rest and muscular rigidity. Physical examination reveals the patient has a “masklike” face. In this patient, from which one of the following sites would biopsies most likely reveal intracytoplasmic eosinophilic inclusions within neurons?

A. Basal ganglia

B. Caudate nucleus

C. Hippocampus

D. Midbrain

E. Substantia nigra

12. An elderly woman presented with progressive dementia. Within 2 years she had to be admitted to a nursing home, where she died 3 years later. Autopsy revealed bilateral, symmetrical atrophy of the frontal lobes with wide sulci and narrow gyri. Microscopic examination of tissue taken from this area revealed bundles of filaments in the cytoplasm of neurons and focal collections of neuritic processes surrounding central amyloid cores. Which one of the following diseases is responsible for her death?

A. Parkinson disease

B. Alzheimer disease

C. Hypertention

D. Diabetes mellitus

E. Spongiform encephalopathy

13. Physical examination of a 34-year-old woman with a new onset of intentionmaldinisong tremor finds medial rectus palsy on attempted lateral gaze in the adducting eye and monocular nystagmus in the abducting eye with convergence. Which of the following is the most likely cause of these clinical findings?

A. An apical lung cancer

B. A pituitary adenoma

C. Diabetes mellitus

D. Multiple sclerosis

E. Tertiary syphilis

14. A 62-year-old man presents with rapidly

progressive dementia and dies. At autopsy, marked spongiform degeneration of the brain is found. The familial form of this disorder is associated with mutations of. what gene?

A. FGFR3
B. PRNP
C. PTEN
D. SOD1
E. UBE3A

15. A lumbar puncture is performed on a patient with headaches, photophobia, clouding of consciousness, and stiffness of the neck. If these symptoms are the result of bacterial infection of the meninges, examination of the cerebrospinal fluid (CSF) will most likely reveal which of the following patterns?

A. Increased, Cloudy, Increased, Decreased, Neutrophils
B. Increased, Clear, Increased, Normal, Lymphocytes
C. Increased, Clear, Increased, Normal, Mononuclear cells
D. Decreased, Clear, Decreased, Normal, Lymphocytes
E. Increased, Clear, Increased, Normal, mixed

16. Which of the following clinical findings is not the characteristic of purulent meningitis:

A. Large numbers of lymphocytes in the cerebrospinal fluid (CSF)
B. Stiff neck
C. Headache
D. Turbid appearance of CSF
E. Large numbers of neutrophils in the CSF

17. Which of the following pathological changes is not associated with Herpes Simplex encephalitis:

A. Perivascular mononuclear inflammatory infilation
B. Localized aggregates of microglial cells (microglial nodule)
C. Smudgy inclusion bodies within the nuclei of neurons
D. Necrotizing foci involving the temporal lobe
E. Negri body

18. Which of the following disorders is not included in the spongiform encephalopathies:

A. Creutzfeldt-Jakob disease (CJD)
B. Kuru disease
C. Gerstmann-Straussler syndrome
D. Mad cow disease
E. Guillain-Barre syndrome

19. Which of the following is the pathogen of spongiform encephalopathies:

A. DNA virus
B. RNA virus
C. Reverse transcription virus
D. Prionsmaldinisong

E. Chronic virus

20. Which of the following pathological changes is not associated with Alzheimer disease:
    A. Wild-spread senile plaque
    B. Senile plaque
    C. Neurofibrillary tangles
    D. Amyloid angiopathy
    E. Lewy body
21. Which of the following pathological changes is associated with Parkinson disease:
    A. Hirano body
    B. Russell body
    C. Aschoff body
    D. Mallory body
    E. Lewy body
22. Which of the following is the major component of neurofibrillary tangles:
    A. β-amyloid (Aβ)
    B. Hyperphosphorylated tau protein
    C. Apoprotein E
    D. Glial filament acidic protein
    E. S-100 protein
23. Which of the following morphological changes is not associated with Alzheimer disease:
    A. Cerebral atrophy
    B. Depigmentation of substantia nigra
    C. Dilation of the cerebral ventricles
    D. Neurofibrillary tangles
    E. Senile plaque
24. The substance deposited in senile plaque is the breakdown product of which of the following proteins:
    A. Neurofilaments
    B. Microtubule protein
    C. Amyloid precursor protein (APP)
    D. Glial filament acidic protein
    E. S-100 protein
25. A 79-year-old woman who had a medical history of poorly controlled chronic hypertension died while at home. During the autopsy the pathologist finds a recent intracerebral hemorrhage involving the putamen. Histologic sections from this area reveal hyaline material deposited in the walls of cerebral arterioles, an abnormality that is associated with the formation of which of the following types of aneurysm?
    A. Atherosclerotic aneurysm
    B. Berry aneurysm
    C. Charcot-Bouchard aneurysm formed in chronic hypertension
    D. Septic aneurysm
    E. Saccular aneurysm
26. Which of the following is the most common cause of dementia in the elderly:
    A. Alzheimer disease
    B. Cerebrovascular disease
    C. Parkinson disease
    D. Pick diseasemaldinisong
    E. Prion-associated disorders

27. Which of the following pathological changes is typically found in Alzheimer disease:
    A. Hirano body
    B. Russell body
    C. Aschoff body
    D. Mallory body
    E. Lewy body
28. Which of the following is the major component of senile plaque:
    A. Neurofilaments
    B. Microtubule protein
    C. β-amyloid substance
    D. Glial filament acidic protein
    E. S-100 protein
29. Parkinson disease is an abnormality of which one of the following anatomic sites?
    A. Cerebral cortex
    B. Basal ganglia
    C. Substantia nigra
    D. Cerebellum
    E. Spinal cord
30. Which of the following is the basic pathological change of Parkinson disease:
    A. Loss of dopamine-secreting neurons of the substantia nigra
    B. Atrophy of the cerebral temporal lobe
    C. Loss of neurons in the frontal lobe
    D. Loss of cerebral cortical neurons
    E. Loss of anterior horn motor neurons of the spinal cord

答案

中文题

1.E	2.A	3.D	4.B	5.B	6.B	7.E	8.E	9.A	10.B
11.D	12.C	13.B	14.B	15.A	16.A	17.A	18.D	19.A	20.B
21.B	22.A	23.C	24.B	25.E	26.D	27.B	28.C	29.A	30.C
31.A	32.C	33.D	34.A	35.A	36.B	37.E	38.D	39.A	40.D
41.E	42.B	43.A	44.D	45.D	46.D	47.A	48.E	49.C	50.C
51.D	52.B	53.C	54.B	55.D	56.B	57.D	58.C	59.E	60.E
61.B	62.D	63.B	64.A	65.A	66.A	67.A	68.E	69.D	70.A
71.E	72.C	73.C	74.D	75.D	76.E	77.C	78.A	79.E	80.E
81.A	82.B	83.A	84.C	85.C	86.A	87.D			

英文题

1.C	2.C	3.B	4.A	5.E	6.A	7.A	8.B	9.E	10.A
11.E	12.B	13.B	14.B	15.A	16.A	17.E	18.E	19.D	20.E
21.E	22.B	23.B	24.C	25.C	26.A	27.A	28.B	29.C	30.A

13 传染病

中文题

1. 结核病的免疫反应属
   A. I 型超敏反应
   B. II 型超敏反应
   C. III 型超敏反应
   D. IV 型超敏反应
   E. 以上都不是
2. 与结核杆菌的毒力有关的是
   A. 脂质
   B. 多糖
   C. 蛋白
   D. 外毒素
   E. 内毒素
3. 结核结节主要细胞成分是
   A. 纤维母细胞
   B. 类上皮细胞
   C. Langhans 巨细胞
   D. 淋巴细胞
   E. 中性粒细胞
4. 不符合结核病的描述是
   A. 主要由鸟型结核杆菌引起
   B. 主要经呼吸道传播
   C. 形成结核性肉芽肿
   D. 结核性肉芽肿具有诊断意义
   E. 干酪样坏死具有特异性
5. 当机体抵抗力低下、结核杆菌毒力强时，结核病变呈
   A. 渗出性病变
   B. 增生性病变
   C. 纤维化
   D. 吸收、消散
   E. 钙化
6. 人体内吞噬消灭结核杆菌的细胞是
   A. T 淋巴细胞
   B. 浆细胞
   C. 巨噬细胞
   D. 中性粒细胞
   E. 嗜酸粒细胞
7. 结核病的特征性病变是
   A. 浆液性炎
   B. 纤维素性炎
   C. 纤维素样坏死
   D. 结核结节和干酪样坏死
   E. 空洞形成
8. 当机体抵抗力强、结核杆菌数量少、毒力弱时，结核病变呈
   A. 渗出性病变
   B. 增生性病变
   C. 坏死性病变
   D. 病变扩大
   E. 蔓延扩散
9. 符合原发性肺结核病的描述是
   A. 不易发生血道播散
   B. 病变多从肺尖开始
   C. 不易发生支气管播散
   D. 不易经淋巴道播散
   E. 大多数自然痊愈
10. 不符合原发性肺结核病的描述是
    A. 多见于儿童

B. 肺内原发灶多位于上叶下部或下叶上部近肺膜处
C. 不易经淋巴道播散
D. 原发综合征是其病变特点
E. 多数患者自然痊愈

11. 不符合慢性粟粒性肺结核病的描述的是
A. 多见于成年人
B. 肺原发综合征多已钙化痊愈
C. 患者多因结核性脑膜炎死亡
D. 干酪样坏死破入颈内静脉，血行播散引起
E. 肺内形成新旧不等病变

12. 不符合急性粟粒性肺结核病的是
A. 可以是全身粟粒性结核病的一部分
B. 两肺充血
C. 两肺密布粟粒大小的结节
D. 结核杆菌来自肺外结核病灶
E. 临床上病情危重

13. 不符合全身性粟粒性结核病的是
A. 可发生类白血病反应
B. 累及全身各器官
C. 结核杆菌来自肺门或纵隔淋巴结
D. 受累器官密布粟粒大小结节
E. 常伴有结核性脑膜炎

14. 原发性肺结核病的肺内原发灶常位于
A. 肺尖
B. 肺上叶下部或肺下叶上部靠近胸膜处
C. 肺门
D. 肺背部
E. 肺下部

15. 符合原发性肺结核病的描述是
A. 多发生于成人
B. 结核性脑膜炎是主要死因
C. 病变不播散
D. 难于痊愈
E. 慢性粟粒性肺结核病常见

16. 不符合原发性肺结核病的描述是
A. 多发生于儿童
B. 结核性脑膜炎是主要死因
C. 多数患者自然痊愈
D. 少数患者病情恶化
E. 慢性粟粒性肺结核病常见

17. 不符合继发性肺结核病变特点的是
A. 病变多从肺尖部开始
B. 常为新旧病变交杂
C. 主要通过淋巴道蔓延
D. 全身粟粒性结核病极少见
E. 以增生性病变为主

18. 不符合慢性纤维空洞型肺结核的描述是
A. 可见一个或多个后壁空洞
B. 空洞壁参差不齐
C. 病变上重下轻
D. 可演变为硬化型肺结核
E. 可引起大咯血及脓气胸

19. 男性，31 岁，1 年来，低热、疲倦、乏力、夜间盗汗，并有咳嗽。X 线检查，可见两肺上叶多个大小不等的后壁空洞。结核菌素试验阳性。不符合该患者疾病的描述是
A. 可由浸润性肺结核恶化而来
B. 容易发生血道播散

C. 容易发生气道播散
D. 可引起干酪性肺炎
E. 可引起肠结核

20. 男性，31 岁，1 年来，低热、疲倦、乏力、夜间盗汗，并有咳嗽。X 线检查，可见两肺上叶多个大小不等的后壁空洞。结核菌素试验阳性。不符合该患者疾病的描述是
A. 病变上重下轻
B. 可引起硬变性肺结核
C. 可引起大咯血
D. 属开放性肺结核
E. 易引起全身性粟粒性结核病

21. 不符合局灶性肺结核病的描述是
A. 属于活动性肺结核
B. 病变多位于肺尖部
C. 多为增生性病变
D. 可有干酪样坏死
E. 常发生纤维化、钙化

22. 不符合浸润性结核病的描述是
A. 属于无活动性肺结核
B. 常发生干酪样坏死
C. 常伴有全身症状
D. 可引起干酪样肺炎
E. 可经纤维化、钙化痊愈

23. 不符合急性空洞的描述是
A. 可由干酪样肺炎引起
B. 可由浸润型肺结核引起
C. 多为厚壁空洞
D. 可引起气道播散
E. 可引起结核性脓气胸

24. 不符合干酪样肺炎的描述是
A. 病情急骤
B. 多由气道播散引起
C. 常有急性空洞形成
D. 以增生性病变为主
E. 以渗出和坏死病变为主

25. 不是经血源播散引起的肺外器官结核病是
A. 肠结核
B. 肾结核
C. 结核性脑膜炎
D. 骨结核
E. 输卵管结核

26. 男性，8 岁，1 年来，低热、疲倦、乏力、夜间盗汗，并有咳嗽。X 线检查，可见右肺下叶上部近肺膜处见一边界模糊的团状阴影，肺门多数淋巴结肿大。结核菌素试验阳性。不符合该患者疾病的描述是
A. 可发生淋巴道播散
B. 可发生血道播散
C. 可发生气道播散
D. 可发生干酪样坏死
E. 多数患者发生急性空洞

27. 女性，40 岁，左侧腰部疼痛 1 年，尿中带血，时有脓尿，尿中可查见结核菌。肾盂造影检查，左肾可见空洞。符合该患者疾病的描述是
A. 左肾结核病
B. 可引起膀胱结核
C. 可引起附睾结核
D. 可引起右肾结核
E. 易引起腹膜结核

28. 关于肺外器官结核病的描述，正确的是
    A. 肠结核以结肠肝曲为其好发部位
    B. 附睾结核由血源播散而来
    C. 结核性脑膜炎的脑顶病变最明显
    D. 肾结核病变开始于肾盂
    E. 脊椎结核多累及第 10 胸椎至第 2 腰椎
29. 肠结核形成
    A. 环形溃疡
    B. 烧瓶状溃疡
    C. 与肠的长轴平行的椭圆形溃疡
    D. 火山口状溃疡
    E. 大小不等、形态不一的浅溃疡
30. 结核性脑膜炎的病变
    A. 以脑底部最明显
    B. 以脑顶部最明显
    C. 以大脑灰质最明显
    D. 以脊髓灰质前角最明显
    E. 主要累及 Willis 环和大脑中动脉
31. 女性生殖系统结核最多发生于
    A. 阴道
    B. 子宫颈
    C. 子宫内膜
    D. 输卵管
    E. 卵巢
32. 男性生殖系统结核病多见于
    A. 前列腺
    B. 精囊
    C. 输精管
    D. 附睾
    E. 睾丸
33. 肠结核的好发部位为
    A. 空肠
    B. 回肠
    C. 升结肠
    D. 回盲肠
    E. 降结肠
34. 不符合结核球的描述是
    A. 境界清楚的球形病变
    B. 中央为干酪样坏死
    C. 可由慢性空洞演化而来
    D. 可由浸润型肺结核演化而来
    E. 不恶化、扩散
35. 不符合结核性胸膜炎的描述是
    A. 可由继发性肺结核引起
    B. 可由原发性肺结核引起
    C. 多为渗出性胸膜炎
    D. 多为增生性胸膜炎
    E. 可引起胸膜粘连
36. 不符合结核性腹膜炎的描述是
    A. 可由肠结核引起
    B. 可由肠系膜淋巴结核引起
    C. 多伴有剧烈的压痛和反跳痛
    D. 可为浆液性或纤维素性炎
    E. 可由输卵管结核引起
37. 不符合关节结核的描述是
    A. 多经血道播散而来
    B. 可为浆液性炎
    C. 可为纤维素性炎
    D. 可形成干酪样坏死
    E. 可引起关节强直
38. 不符合骨结核的描述是
    A. 多经血道播散而来
    B. 可形成死骨

C. 可形成冷脓肿
D. 可形成干酪样坏死
E. 脊椎结核以上部胸椎多见

39. 符合结核性脑膜炎的描述是
A. 成人多见
B. 多由脑内结核球恶化而来
C. 以脑顶部病变最显著
D. 可引起脑积水
E. 以中性粒细胞渗出为主

40. 符合肠结核的描述是
A. 可为原发性结核病变
B. 主要累及空肠
C. 形成纵行溃疡
D. 可引起肠癌
E. 不累及肠系膜淋巴结

41. 伤寒病的肠道病变特点是
A. 好发于结肠
B. 不累及肠系膜淋巴结
C. 形成与肠长轴平行的椭圆形溃疡
D. 溃疡愈合后形成瘢痕引起肠狭窄
E. 病变愈复后不会再次复发

42. 伤寒小结的主要细胞是
A. 类上皮细胞
B. 多核巨细胞
C. 淋巴细胞
D. 巨噬细胞
E. 浆细胞

43. 伤寒病人肥达反应（Widal reaction）阳性出现于病程的
A. 第一周以后
B. 第二周以后
C. 第三周以后
D. 第四周以后
E. 第五周以后

44. 伤寒病的潜伏期，伤寒杆菌寄生、繁殖的细胞是
A. 淋巴细胞
B. 巨噬细胞
C. 中性粒细胞
D. 嗜酸粒细胞
E. 浆细胞

45. 伤寒带菌者，伤寒杆菌寄生、繁殖的器官是
A. 回肠下段
B. 淋巴结
C. 胆囊
D. 肝脏
E. 脾脏

46. 伤寒小结一般不发生在
A. 回肠
B. 脾脏
C. 骨髓
D. 胆囊
E. 肝脏

47. 不符合伤寒的描述是
A. 可引起肠出血和肠穿孔
B. 可引起支气管肺炎
C. 可引起心肌炎
D. 可引起胆囊炎
E. 可引起肾盂肾炎

48. 不符合伤寒杆菌的描述是
A. 革蓝染色阳性菌
B. 有强烈的内毒素
C. 经口感染

D. O 和 H 抗原的抗原性较强
E. 肥达反应可辅助临床诊断

49. 肠伤寒形成
A. 环形溃疡
B. 烧瓶状溃疡
C. 与肠的长轴平行的椭圆形溃疡
D. 火山口状溃疡
E. 大小不等、形态不一的浅溃疡

50. 伤寒的传染途径是
A. 消化道传播
B. 呼吸道传播
C. 经血液传播
D. 性传播
E. 内源性感染

51. 伤寒肠道病变的好发部位是
A. 空肠
B. 回肠下段
C. 回盲部
D. 横结肠
E. 乙状结肠

52. 伤寒细胞是指
A. 类上皮细胞
B. 多核巨细胞
C. 泡沫细胞
D. 浆细胞
E. 吞噬伤寒杆菌、淋巴细胞、红细胞的巨噬细胞

53. 伤寒的合并症不包括
A. 肠出血
B. 肠穿孔
C. 支气管肺炎
D. 大叶性肺炎
E. 中毒性心肌炎

54. 伤寒可引起心肌的病变是
A. 细菌性心肌炎
B. 中毒性心肌炎
C. 心肌梗死
D. 心肌肥大
E. 肉芽肿性心肌炎

55. 伤寒并发肠穿孔和出血容易发生在
A. 潜伏期
B. 髓样肿胀期
C. 坏死期
D. 溃疡期
E. 愈合期

56. 不符合伤寒的临床表现的是
A. 持续高热
B. 肝脾肿大
C. 外周血白细胞计数增高
D. 皮肤玫瑰疹
E. 相对缓脉

57. 细菌性痢疾传播途径是
A. 消化道传播
B. 呼吸道传播
C. 经血液传播
D. 性传播
E. 内源性感染

58. 细菌性痢疾的肠道病变特点是
A. 出血性炎
B. 蜂窝组织炎
C. 假膜性炎
D. 肉芽肿性炎
E. 浆液性炎

59. 细菌性痢疾的临床表现不包括

A. 发热
B. 压痛、反跳痛
C. 腹泻
D. 脓血便
E. 里急后重

60. 细菌性痢疾可引起
A. 环形溃疡
B. 烧瓶状溃疡
C. 与肠的长轴平行的椭圆形溃疡
D. 火山口状溃疡
E. 大小不等、形态不一的浅溃疡

61. 急性细菌性痢疾可引起
A. 滤泡性肠炎
B. 肠壁的蜂窝织炎
C. 肠壁的出血性梗死
D. 肠壁肉芽肿形成
E. 肠梗阻

62. 中毒型细菌性痢疾的特点是
A. 常好发生于老年人
B. 常由毒力强的志贺氏菌引起
C. 有明显的腹泻及脓血便
D. 可以出现滤泡性肠炎的变化
E. 容易转为慢性细菌性痢疾

63. 男性，7 岁，因饮食不洁，1 天来发热、腹痛、腹泻、脓血便。大便检查 RBC、WBC 满视野，并有脓球。符合该患者疾病的描述是
A. 直肠粘膜高度充血、水肿、增厚
B. 直肠粘膜大量纤维素及中性粒细胞渗出，构成伪膜
C. 直肠粘膜粘膜多数浅表溃疡形成
D. 直肠粘膜炎症细胞浸润
E. 以上病变都存在

64. 急性细菌性痢疾的肠粘膜表面的假膜不具有
A. 纤维素
B. 坏死组织
C. 中性粒细胞
D. 细菌
E. 真菌

65. 流行性乙型脑炎是
A. 化脓性炎症
B. 变质性炎症
C. 纤维素性炎症
D. 浆液性炎症
E. 肉芽肿性炎症

66. 流行性乙型脑炎的主要病理变化是
A. 神经细胞变性坏死
B. 蛛网膜下腔积脓
C. 肉芽肿形成
D. 微小动脉瘤形成
E. 脑疝形成

67. 流行性乙型脑炎的主要侵犯部位是
A. 脊髓灰质
B. 延脑
C. 小脑
D. 大脑灰质及神经核团
E. 神经节

68. 符合流行性乙型脑炎的描述是
A. 经呼吸道传播
B. 冬春发病
C. 可发生华-弗综合征
D. 可引起智力低下
E. 克氏症阳性

69. 不符合流行性乙型脑炎的描述是
A. 大脑内有软化灶形成
B. 淋巴细胞和单核细胞浸润并有血管套形成
C. 大脑神经元变性坏死
D. 蛛网膜下腔大量白细胞
E. 小胶质细胞结节形成

70. 流行性乙型脑炎的病原体是
A. RNA 病毒
B. 细菌
C. 真菌
D. 支原体
E. DNA 病毒

71. 流行性乙型脑炎的传播途径是
A. 消化道传播
B. 呼吸道传播
C. 经血液传播
D. 性传播
E. 蚊虫传播

72. 流行性乙型脑炎的患者多为
A. 男性
B. 女性
C. 儿童
D. 成人
E. 老人

73. 流行性乙型脑炎的病变最轻的部位是
A. 大脑皮质
B. 基底节
C. 丘脑
D. 中脑
E. 脊髓

74. 不属于流行性乙型脑炎的病变是
A. 血管周围淋巴细胞浸润及血管套形成
B. 筛网状软化灶
C. 蛛网膜下腔中性粒细胞浸润
D. 胶质结节形成
E. 小脑蒲肯野细胞大量丢失

75. 不符合流行性乙型脑炎的临床表现是
A. 嗜睡
B. 昏迷
C. 角弓反张
D. 头痛
E. 呕吐

76. 不符合流行性乙型脑炎的预后的描述是
A. 可引起痴呆
B. 可引起语言障碍
C. 可引起智力低下
D. 可引起脑积水
E. 可引起肢体瘫痪

77. 不符合流行性脑脊髓膜炎的临床表现是
A. 颈强直
B. 脑脊液淋巴细胞增多
C. 角弓反张
D. 头痛
E. 呕吐

78. 不符合流行性脑脊髓膜炎的预后的是
A. 大多可痊愈
B. 可引起脑积水
C. 可引起面神经麻痹
D. 常引起智力低下
E. 可引起耳聋

79. 流行性脑脊髓膜炎的病原体是
A. RNA 病毒
B. 细菌
C. 真菌
D. 支原体
E. DNA 病毒
80. 属于脑膜刺激症状的是
A. Kernig 症
B. 耳聋
C. 面神经麻痹
D. 头痛
E. 呕吐
81. 流行性脑膜炎的主要病理变化是
A. 软化灶形成
B. 神经细胞变性坏死
C. 淋巴细胞血管周围套状浸润
D. 蛛网膜下腔积脓
E. 小胶质细胞结节状增生
82. 流行性脑膜炎属
A. 变质性炎症
B. 增生性炎症
C. 化脓性炎症
D. 浆液性炎症
E. 肉芽肿性炎症
83. 流行性脑膜炎主要侵犯
A. 硬脑膜外
B. 硬脑膜
C. 软脑膜
D. 大脑皮质
E. 蛛网膜和软脑膜
84. 导致脑疝的大脑疾病是
A. 流行性乙型脑炎
B. 脑水肿
C. 脑出血
D. 脑内肿瘤
E. 以上都可能
85. 不符合流行性脑膜炎的描述是
A. 皮肤、粘膜有出血点和瘀斑
B. 脑软化灶形成
C. 脑疝形成
D. 高热、头痛、呕吐
E. 脑膜刺激症明显
86. 不符合流行性脑膜炎的描述是
A. 可引起脑膜刺激症
B. 可引起颅内压升高症状
C. 可引起脑脊液浑浊或脓样
D. 常引起神经细胞变性坏死
E. 可引起皮肤瘀点和瘀斑
87. 流行性脑膜炎的病变
A. 以脑底部最明显
B. 以脑顶部最明显
C. 以大脑灰质最明显
D. 以脊髓灰质前角最明显
E. 主要累及 Willis 环和大脑中动脉
88. 不符合暴发性脑膜炎双球菌性败血症的描述是
A. 皮肤和粘膜出现淤点和淤斑
B. 双侧肾上腺广泛出血
C. 蛛网膜下腔大量脓性渗出物
D. 高热、头痛、呕吐
E. 可发生周围循环衰竭
89. 脊髓灰质炎的病变
A. 以脑底部最明显
B. 以脑顶部最明显

C. 以大脑灰质最明显
D. 以脊髓灰质前角最明显
E. 以小脑最明显

90. 脊髓灰质炎最主要的传播途径是
A. 消化道传播
B. 呼吸道传播
C. 经血液传播
D. 性传播
E. 蚊虫传播

91. 不符合脊髓灰质炎的病变特点的是
A. 以脊髓腰膨大最为严重
B. 以脊髓颈膨大最为严重
C. 主要累及运动神经元
D. 神经元变性坏死
E. 中性粒细胞和淋巴细胞浸润

92. 脊髓灰质炎的病原体是
A. RNA 病毒
B. 细菌
C. 真菌
D. 支原体
E. DNA 病毒

93. 脊髓灰质炎病变最轻的部位是
A. 脊髓腰膨大
B. 脊髓颈膨大
C. 延髓
D. 脑桥
E. 大脑皮质

94. 脊髓灰质炎引起下肢麻痹主要是由于
A. 充血水肿
B. 胶质瘢痕形成
C. 小胶质细胞增生
D. 运动神经元变性坏死
E. 中性粒细胞浸润

95. 脊髓灰质炎病毒感染的大多数表现为
A. 隐性感染
B. 轻型
C. 非麻痹型
D. 麻痹型
E. 显性感染

96. 脊髓灰质炎病毒感染最少引起
A. 隐性感染
B. 轻型
C. 非麻痹型
D. 麻痹型
E. 显性感染

97. 麻风病最常见的类型是
A. 结核样型麻风
B. 瘤型麻风
C. 界线类偏结核型麻风
D. 界线类偏瘤型麻风
E. 未定类麻风

98. 麻风的早期诊断依赖于
A. 血液检查
B. 分泌物检查
C. 粪便检查
D. 尿液检查
E. 活体组织检查

99. 麻风细胞来源于
A. 中性粒细胞
B. 淋巴细胞
C. 巨噬细胞
D. 浆细胞
E. 肥大细胞

100. 麻风的传染途径主要是

A. 消化道传播
B. 呼吸道传播
C. 密切接触传播
D. 性传播
E. 蚊虫传播

101. 最易查见麻风杆菌和最具传染性的麻风类型是
A. 结核样型麻风
B. 瘤型麻风
C. 未定类麻风
D. 界限类麻风偏瘤型
E. 界限类麻风偏结核样型

102. 麻风病属于
A. 纤维素性炎
B. 变质性炎
C. 肉芽肿性炎
D. 浆液性炎
E. 化脓性炎

103. 麻风病的传播途径主要是
A. 接触传播
B. 蚊虫传播
C. 消化道传播
D. 呼吸道传播
E. 经血传播

104. 麻风病患者四肢畸形是由于
A. 多发性肌炎
B. 纤维素性关节炎
C. 瘢痕形成
D. 神经损害
E. 血栓闭塞性脉管炎

105. 肠阿米巴病最常发生在
A. 空肠
B. 回肠
C. 盲肠和升结肠
D. 横结肠
E. 乙状结肠和直肠

106. 溶组织内阿米巴滋养体对组织的破坏主要通过
A. 水解酶作用
B. 毒素作用
C. 继发细菌感染
D. 机械损伤作用
E. 吞噬作用

107. 不属于肠阿米巴病的并发症为
A. 肺脓肿
B. 脑脓肿
C. 肝脓肿
D. 结肠癌
E. 肠狭窄

108. 肠阿米巴病溃疡的特点是
A. 环形溃疡
B. 烧瓶状溃疡
C. 与肠的长轴平行的椭圆形溃疡
D. 火山口状溃疡
E. 大小不等、形态不一的浅溃疡

109. 阿米巴滋养体所引起的组织坏死为
A. 凝固性坏死
B. 干酪样坏死
C. 纤维素样坏死
D. 液化性坏死
E. 湿性坏疽

110. 诊断肠阿米巴病的依据是
A. 肠穿孔
B. 暗红色、糊状大便

C. 大便检查可见阿米巴滋养体
D. 腹泻
E. 里急后重

111. 诊断阿米巴肝脓肿的依据是
A. 多发性脓肿
B. 多位于肝右叶
C. 脓液呈果酱样
D. 脓肿壁呈破絮样
E. 脓肿周围组织可见阿米巴滋养体

112. 阿米巴病的传播途径是
A. 消化道传播
B. 呼吸道传播
C. 密切接触传播
D. 性传播
E. 蚊虫传播

113. 血吸虫病对机体影响最严重的病变是
A. 尾蚴性皮炎
B. 童虫引起的肺病变
C. 成虫引起的静脉内膜炎
D. 虫卵引起的虫卵结节
E. 成虫引起的贫血

114. 血吸虫病的虫卵结节中，不具备的成分是
A. 干酪性坏死
B. 类上皮细胞
C. 多核巨细胞
D. 淋巴、单核细胞
E. 嗜酸性粒细胞

115. 侵入人体皮肤引起血吸虫病的是
A. 虫卵
B. 毛蚴
C. 胞蚴
D. 尾蚴
E. 成虫

116. 易并发结肠癌的疾病是
A. 阿米巴病
B. 血吸虫病
C. 细菌性痢疾
D. 伤寒
E. 肠结核

117. 肠道血吸虫病，一般不发生
A. 肠粘膜溃疡
B. 肠粘膜息肉
C. 肠粘膜虫卵结节形成
D. 肠穿孔
E. 肠壁增厚、肠腔狭窄

118. 我国流行的血吸虫病是
A. 曼氏血吸虫
B. 埃及血吸虫
C. 日本血吸虫
D. 湄公血吸虫
E. 间插血吸虫

119. 日本血吸虫引起肉芽肿病变的是
A. 虫卵
B. 毛蚴
C. 尾蚴
D. 童虫
E. 成虫

120. 日本血吸虫病引起的肝硬化为
A. 胆汁性肝硬化
B. 门脉性肝硬化
C. 坏死后性肝硬化
D. 干线型肝硬化
E. 淤血性肝硬化

121. 日本血吸虫虫卵可引起
A. 皮炎
B. 嗜酸性脓肿和假结核结节
C. 肠系膜静脉及门静脉炎
D. 肺炎
E. 胆管炎
122. 日本血吸虫尾蚴可引起
A. 皮炎
B. 嗜酸性脓肿和假结核结节
C. 肠系膜静脉及门静脉炎
D. 肺炎
E. 胆管炎
123. 日本血吸虫成虫可引起
A. 皮炎
B. 嗜酸性脓肿和假结核结节
C. 肠系膜静脉及门静脉炎
D. 肺炎
E. 胆管炎
124. 日本血吸虫童虫可引起
A. 皮炎
B. 嗜酸性脓肿和假结核结节
C. 肠系膜静脉及门静脉炎
D. 多器官点状出血
E. 胆管炎
125. 日本血吸虫虫卵结节的形成属于
A. I 型超敏反应
B. II 型超敏反应
C. III 型超敏反应
D. IV 型超敏反应
E. 以上都不是
126. 引起急性血吸虫病的主要是
A. 虫卵
B. 毛蚴
C. 尾蚴
D. 童虫
E. 成虫
127. 不符合急性血吸虫病的描述是
A. 起病急骤
B. 发热
C. 肝脾肿大
D. 腹泻
E. 血中中性粒细胞升高
128. 血吸虫性肝硬化的病变特点是
A. 结节大小一致，纤维间隔薄而均匀
B. 结节大小不一，纤维间隔厚而不匀
C. 肝脏呈深绿色
D. 大量假小叶形成
E. 汇管区虫卵沉积，纤维组织增生
129. 艾滋病的病原体是
A. HIV
B. HCV
C. HBV
D. HPV
E. CMV
130. 人类免疫缺陷病毒（HIV）属于
A. DNA 病毒
B. 缺陷病毒
C. 逆转录病毒
D. 肠道病毒
E. 肿瘤病毒
131. HIV 病毒主要感染
A. B 细胞
B. T 细胞
C. NK 细胞

D. T_8 细胞

E. T_4 细胞和单核巨噬细胞

132. 与 HIV 病毒包鞘的 GP120 蛋白具有高度亲和力的淋巴细胞表面分子是

A. TCR

B. CD3

C. CD4

D. CD8

E. CD28

133. 不符合艾滋病传播途径的是

A. 接吻传播

B. 母婴传播

C. 输血传播

D. 异性性交传播

E. 同性恋性交传播

134. 艾滋病传播途径的是

A. 消化道传播

B. 呼吸道传播

C. 接触传播

D. 性传播

E. 蚊虫传播

135. 艾滋病免疫缺陷的主要原因是

A. 吞噬细胞功能障碍

B. B 细胞功能障碍

C. 补体缺陷

D. T_4 细胞大量减少

E. T_8 细胞大量减少

136. HIV 感染最常用的实验室检查方法是

A. 酶联免疫吸附法

B. 免疫荧光检查

C. 放射免疫测定法

D. 免疫印迹检测法

E. 放射免疫沉淀法

137. 艾滋病患者外周血的 T_4 细胞

A. ＜ 200/μl

B. ＜ 400/μl

C. ＜ 600/μl

D. ＜ 800/μl

E. ＜ 1000/μl

138. 艾滋病最常见的肺部感染是

A. 曲菌感染

B. 新形隐球菌感染

C. 巨细胞病毒感染

D. 卡氏肺孢子虫感染

E. 白色念珠菌感染

139. 艾滋病最常见的恶性肿瘤是

A. 鳞状细胞癌

B. 腺癌

C. 未分化癌

D. 恶性淋巴瘤

E. Kaposi 肉瘤

140. 晚期艾滋病患者淋巴结的病变是

A. 淋巴滤泡增生

B. 副皮质区淋巴细胞增生

C. 窦组织细胞增生

D. 淋巴细胞消失殆尽

E. 淋巴结结构正常

141. Kaposi 肉瘤是

A. 血管内皮起源的恶性肿瘤

B. 纤维组织起源的恶性肿瘤

C. 横纹肌起源的恶性肿瘤

D. 淋巴细胞起源的恶性肿瘤

E. 平滑肌起源的恶性肿瘤

142. 艾滋病机会性感染的特点是

A. 单一病原体感染
B. 单一器官受累
C. 感染易于控制
D. 多病原、多器官、多系统受累
E. 炎症反应典型

143. 不符合艾滋病机会性感染的特点是
A. 多种病原体感染
B. 多器官受累
C. 感染易于控制
D. 多系统受累
E. 出现复杂的临床表现

144. 不符合艾滋病机会性感染的特点是
A. 单一病原体感染
B. 多器官受累
C. 感染不易控制
D. 多系统受累
E. 出现复杂的临床表现

145. 在梅毒病变中，浸润的炎细胞主要是
A. 中性粒细胞
B. 嗜酸粒细胞
C. 淋巴细胞及浆细胞
D. 肥大细胞
E. 巨噬细胞

146. 第一期梅毒的特征病变是
A. 树胶肿形成
B. 主动脉瘤形成
C. 梅毒疹
D. 外生殖器硬性下疳
E. 脊髓痨

147. 引起口腔鹅口疮的真菌为
A. 放线菌
B. 曲菌
C. 毛霉菌
D. 念珠菌
E. 隐球菌

148. 不符合淋病的描述是
A. 性交传播
B. 常为急性化脓性炎
C. 多累及泌尿生殖道
D. 可引起尿道狭窄
E. 与盆腔炎症性疾病无关

149. 深部真菌病不易发生于
A. 恶性肿瘤患者
B. 长期使用广谱抗生素患者
C. 糖尿病患者
D. 使用免疫抑制药物患者
E. 免疫功能正常者

150. 不符合白色念珠菌感染的描述是
A. 多为内源性感染
B. 常形成白色假膜
C. 病变中可见芽生孢子和假菌丝
D. 多发生在消化道、呼吸道、泌尿生殖道
E. 消化道以直肠病变多见

151. 不符合曲菌病的描述是
A. 多经呼吸道感染
B. 可引起支气管炎
C. 可引起支气管肺炎
D. 大量菌丝聚集形成曲菌团
E. 曲菌丝有隔，呈直角分支

152. 不符合隐球菌病的描述是
A. 主要经呼吸道传播
B. 在肺内引起原发病灶
C. 经血道播散到脑和脑膜

D. 引起隐球菌性脑膜炎
E. 病变中异物巨细胞内可见菌丝

153. tuberculoma
A. 结核病
B. 结核瘤
C. 结核结节
D. 结核空洞
E. 结核肉芽肿

154. leprosy
A. 伤寒
B. 痢疾
C. 梅毒
D. 麻风
E. 淋病

155. gonorrhea
A. 伤寒
B. 痢疾
C. 梅毒
D. 麻风
E. 淋病

156. syphilis
A. 伤寒
B. 痢疾
C. 梅毒
D. 麻风
E. 淋病

157. typhoid fever
A. 伤寒
B. 痢疾
C. 梅毒
D. 麻风
E. 淋病

158. schistosomiasis
A. 血吸虫病
B. 阿米巴病
C. 曲菌病
D. 念珠菌病
E. 隐球菌病

159. meningitis
A. 脑炎
B. 脑膜炎
C. 脊髓灰质炎
D. 脑疝
E. 脑病

160. encephalitis
A. 脑炎
B. 脑膜炎
C. 脊髓灰质炎
D. 脑疝
E. 脑病

英文题

如果你已经刷到了这里

却没有发现其实传染病相关的英文题散落在各章之中

这说明你肯定没有好好刷英文题

（笑）

答案

中文题

1.D	2.A	3.B	4.A	5.A	6.C	7.D	8.B	9.E	10.C
11.D	12.D	13.C	14.B	15.B	16.E	17.C	18.B	19.C	20.E
21.A	22.A	23.C	24.D	25.A	26.E	27.E	28.E	29.A	30.A
31.D	32.D	33.D	34.E	35.D	36.C	37.A	38.E	39.D	40.A
41.E	42.D	43.B	44.B	45.C	46.D	47.E	48.A	49.C	50.A
51.B	52.E	53.D	54.B	55.D	56.C	57.A	58.C	59.B	60.E
61.A	62.D	63.E	64.E	65.B	66.A	67.D	68.D	69.D	70.A
71.E	72.C	73.E	74.C	75.C	76.D	77.B	78.D	79.B	80.A
81.D	82.C	83.E	84.E	85.B	86.D	87.B	88.C	89.D	90.A
91.B	92.A	93.E	94.D	95.A	96.D	97.A	98.E	99.C	100.C
101.B	102.C	103.A	104.D	105.C	106.A	107.D	108.B	109.D	110.C
111.E	112.A	113.D	114.A	115.D	116.B	117.D	118.C	119.A	120.D
121.B	122.A	123.C	124.D	125.D	126.A	127.E	128.E	129.A	130.C
131.E	132.C	133.A	134.D	135.D	136.A	137.A	138.D	139.E	140.D
141.A	142.D	143.C	144.A	145.C	146.D	147.D	148.E	149.E	150.E
151.E	152.E	153.B	154.D	155.E	156.C	157.A	158.A	159.B	160.A