

Physics-inspired priors for neutron star mergers improves equation of state constraints

SPENCER J. MAGNALL ^{1,2} CHRISTIAN ECKER ³ LUCIANO REZZOLLA ^{3,4,5} PAUL D. LASKY ^{1,2} AND
SIMON R. GOODE ^{1,2}

¹*School of Physics and Astronomy, Monash University, Clayton VIC 3800, Australia*

²*OzGrav: The ARC Centre of Excellence for Gravitational Wave Discovery, Clayton VIC 3800, Australia*

³*Institut für Theoretische Physik, Max-von-Laue-Strasse 1, 60438 Frankfurt, Germany*

⁴*Frankfurt Institute for Advanced Studies, Ruth-Moufang-Strasse 1, 60438 Frankfurt, Germany*

⁵*School of Mathematics, Trinity College, Dublin 2, Ireland*

ABSTRACT

Gravitational-wave astronomy shows great promise in determining nuclear physics in a regime not accessible to terrestrial experiments. We introduce physics-inspired priors informed by nuclear theory and perturbative quantum chromodynamics calculations, as well as astrophysical measurements of neutron-star masses and radii. When these priors are used in gravitational-wave astrophysical inference, we show a significant improvement on nuclear equation of state constraints. Applying these to the first observed gravitational-wave binary neutron-star merger GW170817, the constraints on the radius of a $1.4 M_{\odot}$ neutron star improve from $R_{1.4} = 12.54^{+1.05}_{-1.54}$ km to $R_{1.4} = 12.11^{+0.91}_{-1.11}$ km and those on the tidal deformability from $\tilde{\Lambda}_{1.186} < 720$ to $226 \leq \tilde{\Lambda}_{1.186} \leq 690$. (90% confidence intervals). We also show these priors can be used to perform model selection between binary neutron star and neutron star-black hole mergers; in the case of GW190425, the results are inconclusive with a Bayes factor of 1.33 in favour of the binary neutron star merger hypothesis, representing only marginal evidence. We advocate for these physics-inspired priors to be used as standard in the literature, and provide open-source code for reproducibility and further adaptation of the method.

Keywords: neutron stars — gravitational waves — equation of state

1. INTRODUCTION

One of the main goals of gravitational wave (GW) astronomy is to infer constraints on the neutron-star equation of state (EOS) from observations of binary neutron star mergers (BNS). The baryon density in neutron stars and their mergers can reach several times the nuclear saturation density $n_s = 0.16 \text{ fm}^{-3}$ and temperatures of the order of tens of MeV (see, e.g., Baiotti & Rezzolla 2017; Paschalidis 2017; Radice et al. 2020, for reviews). Such densities and temperatures are inaccessible to terrestrial experiments and also to ab initio calculations due to the infamous fermionic sign problem, which prevents direct quantum Chromodynamics (QCD) solutions on the lattice.

A common approach to astrophysical GW inference is to use agnostic priors for some GW observables. For example, analyses of the first two observed BNS GW170817 (e.g., Abbott et al. 2017, 2018, 2019) and GW190425 (e.g., Abbott et al. 2020) employed uniform priors on the chirp mass $\mathcal{M}$ and the dimensionless tidal deformability $\tilde{\Lambda}$, implicitly assuming these parameters are uncorellated. These are the two most informative parameters for the neutron star EOS.

The above prior assumptions are overly conservative. Basic physical principles such as causality, consistency with nuclear-physics experiments, and perturbative QCD calculations, imply a strong correlation between $\mathcal{M}$ and $\tilde{\Lambda}$ in neutron-star binaries. Constraints from astrophysical measurements of neutron-star masses and radii further tightens this correlation (Altiparmak et al. 2022a; Ecker & Rezzolla 2022; Ecker & Rezzolla 2022).

Motivated by this, we introduce physics-inspired joint priors on $\mathcal{M}$ and $\tilde{\Lambda}$ that incorporate microphysical constraints from nuclear theory and perturbative QCD, as well as astrophysical constraints on neutron-star masses and radii into astrophysical GW parameter inference. We show a significant improvement on the measurement of the marginalised posterior on $\tilde{\Lambda}$ from the GW observation of GW170817 from $\tilde{\Lambda} = 466^{+308}_{-434}$ with the uniform priors to $\tilde{\Lambda} = 384^{+306}_{-158}$ with the new physics-inspired prior (confidence intervals are 90% throughout). We infer a posterior on the EOS, and find that the neutron-star radius measurement of an equivalent $1.4 M_{\odot}$ star improves from $R_{1.4} = 12.54^{+1.05}_{-1.54}$ km when using uniform priors (see also Most et al. 2018; Capano et al. 2020, for some of the first alternative estimates) to

$R_{1.4} = 12.11^{+0.91}_{-1.11}$ km with our physics-inspired priors. For GW190425, the effective tidal deformability improves from $\tilde{\Lambda} = 381^{+1065}_{-271}$ to $\tilde{\Lambda} = 183^{+180}_{-100}$, a fractional improvement of XXX on the 90% uncertainties.

In addition to astrophysical inference, we show that these priors can be used to perform model selection, and hence distinguish between GW-source types. We perform model selection between a BNS and a neutron star – black hole (NSBH) origin for GW190425 and find a Bayes factor of 1.33 in favour of a BNS origin. When folding in the rates as the prior odds, we find an odds ratio of XXX, indicating

In advocating for this prior becoming standard in the literature, we provide an open-source configuration for the `Bilby` Bayesian inference library (Ashton et al. 2019; Romero-Shaw et al. 2020) currently used for the majority of GW parameter estimation by the LIGO-Virgo-KAGRA (LVK) collaboration (e.g., Aasi et al. 2015; Acernese et al. 2015; Akutsu et al. 2020). This code provides a numerical implementation of the joint probability distribution as a `Bilby` constrained-prior object for $\mathcal{M}$ and $\tilde{\Lambda}$. We describe details of the method and implementation in the next Section.

2. METHODS

The construction of our physics-inspired prior closely follows that of Altiparmak et al. (2022b), which we briefly review here (see Ecker & Rezzolla 2022; Ecker & Rezzolla 2022, for additional details). We begin by building a large set of EOSs by stitching together various components. At the lowest densities ($n < 0.5 n_s$) we adopt the Baym-Pethick-Sutherland (BPS) model (Baym et al. 1971) for the neutron-star crust. In the range $0.5 n_s \leq n < 1.1 n_s$, we randomly sample polytropes to span the range between the softest and stiffest EOSs from Hebeler et al. (2013). At high densities ($n \gtrsim 40 n_s$), corresponding to a baryon chemical potential of $\mu = 2.6$ GeV, we impose the perturbative QCD constraint from Fraga et al. (2014) on the pressure $p(X, \mu)$ of cold quark matter, with the renormalization scale parameter X sampled uniformly in the range $[1, 4]$.

For the intermediate density range ($1.1 n_s < n \lesssim 40 n_s$), we use the parametrization method of Annala et al. (2020), which models the sound speed as a function of the chemical potential, $c_s^2(\mu)$, using piecewise-linear segments:

$$c_s^2(\mu) = \frac{(\mu_{i+1} - \mu) c_{s,i}^2 + (\mu - \mu_i) c_{s,i+1}^2}{\mu_{i+1} - \mu_i}, \quad (1)$$

where μ_i and $c_{s,i}^2$ are parameters defining the i -th segment in the range $\mu_i \leq \mu \leq \mu_{i+1}$. Throughout this work, we adopt natural units where $c = G = 1$, unless otherwise stated.

The number density follows as

$$n(\mu) = n_1 \exp \left(\int_{\mu_1}^{\mu} \frac{d\mu'}{\mu' c_s^2(\mu')} \right), \quad (2)$$

where $n_1 = 1.1 n_s$, and $\mu_1 = \mu(n_1)$ is set by the corresponding polytropic EOS. The pressure is then obtained via

$$p(\mu) = p_1 + \int_{\mu_1}^{\mu} d\mu' n(\mu'), \quad (3)$$

where the integration constant p_1 matches the pressure of the polytrope at $n = n_1$. We numerically integrate Eq. (3) using seven segments for $c_s^2(\mu)$.

Using this framework, we generate approximately 10^6 EOSs by randomly selecting the maximum sound speed $c_{s,\text{max}}^2 \in [0, 1]$ and uniformly sampling the free parameters $\mu_i \in [\mu_1, \mu_{N+1}]$ (where $\mu_{N+1} = 2.6$ GeV) and $c_{s,i}^2 \in [0, c_{s,\text{max}}^2]$. These EOSs are, by construction, consistent with nuclear theory and perturbative QCD uncertainties.

To incorporate astrophysical constraints, we solve the Tolman-Oppenheimer-Volkoff (TOV) equations for each EOS and retain only those satisfying the mass measurements of J0348+0432 (Antoniadis et al. 2013, $M = 2.01 \pm 0.04 M_\odot$) and J0740+6620 (Cromartie et al. 2020; Fonseca et al. 2021, $M = 2.08 \pm 0.07 M_\odot$), discarding EOSs with a maximum mass $M_{\text{TOV}} < 2.0 M_\odot$. In addition, we impose NICER radius constraints from J0740+6620 (Miller et al. 2021; Riley et al. 2021) and J0030+0451 (Riley et al. 2019; Miller et al. 2019), rejecting EOSs with $R < 10.75$ km at $M = 2.0 M_\odot$ or $R < 10.8$ km at $M = 1.1 M_\odot$. We note that, unlike what was done in Altiparmak et al. (2022a) and Ecker & Rezzolla (2022), we do not impose any GW-informed constraints on the binary tidal deformability in the construction of our prior as we are reanalysing these data using the constructed prior.

In the BNS case, the binary tidal deformability of the prior is computed as

$$\tilde{\Lambda}_{\text{NSNS}} := \frac{16}{13} \frac{(12M_2 + M_1)M_1^4 \Lambda_1 + (12M_1 + M_2)M_2^4 \Lambda_2}{(M_1 + M_2)^5}, \quad (4)$$

where M_i , R_i , and $\Lambda_i = \frac{2}{3} k_2 (R_i/M_i)^5$ are the component source-frame masses, radii, and tidal deformabilities, respectively, and k_2 is the second tidal Love number. The source-frame chirp mass is $\mathcal{M}_{\text{chirp}} := (M_1 M_2)^{3/5} (M_1 + M_2)^{-1/5}$ and we define the mass ratio $q := M_2/M_1$ with $M_2 \leq M_1$.

By definition, the tidal deformability of a black hole is zero $\Lambda_{\text{BH}} = 0$, implying the total tidal deformability for a NSBH binary is

$$\tilde{\Lambda}_{\text{NSBH}} := \frac{16}{13} \frac{(12M_{\text{BH}} + M_{\text{NS}})M_{\text{NS}}^4 \Lambda_{\text{NS}}}{(M_{\text{NS}} + M_{\text{BH}})^5}, \quad (5)$$

where M_{NS} and Λ_{NS} are the neutron-star mass and tidal deformability, respectively, and M_{BH} is the black-hole mass. The expressions for the chirp mass and mass ratio are the same, with the understanding that $M_2 = M_{\text{NS}}$ and $M_1 = M_{\text{BH}}$.




Figure 1. Coupled priors for the source-frame chirp mass $\mathcal{M}$ and dimensionless tidal deformability $\tilde{\Lambda}$. Top: prior for BNS mergers. Bottom: prior for NSBH mergers. The horizontal grey bands are the 90% credible intervals for the measured chirp masses of GW170817 (Abbott et al. 2017), GW190425 (Abbott et al. 2020), GW200115 (Abbott et al. 2021) and GW230529 (Abac et al. 2024).

2.1. BNS and NSBH mergers

We use the probability distributions described in the previous section as priors on the source-frame chirp mass and effective tidal deformability as shown in Fig. 1. We implement this in *Bilby* using the `conditional_prior` function, interpolating a 200×200 grid to draw from a joint prior on $\mathcal{M}$ and $\tilde{\Lambda}$. The 90% probability contours (dashed black lines) can be fitted by a simple power law

$$\tilde{\Lambda}_{\min(\max)} = a + b(\mathcal{M}_{\text{chirp}})^c, \quad (6)$$

where the fitting coefficients are $a = -57(-23)$, $b = 583(2690)$ and $c = -3.9(-4.5)$ for the BNS case and $a = 6(-5)$, $b = 296(1700)$ and $c = -6.6(-4.9)$ for the BHNS case. For comparison we also show contours for $a = -41(-0.55)$, $b = 539(2091)$ and $c = -4.5(-5.5)$ (dashed gray lines) corresponding to a prior in which the constraint on the tidal deformability from the GW170817 event, i.e., $\tilde{\Lambda} < 720$, is imposed; this corresponds to what done in Altiparmak et al. (2022a).

We sample directly in source-frame chirp mass, mass ratio and the two tidal deformability parameters $\tilde{\Lambda}$ and $\delta\tilde{\Lambda}$.¹ We

note that q is not a variable that depends on the EOS, and we therefore take the standard approach of choosing a prior on q uniform in the range $q \in [0.125, 1]$ (e.g., Romero-Shaw et al. 2020)². The choice of a lower limit of $q = 0.125$ is due to waveform limitations and is not astrophysically motivated.

Furthermore, while both $\tilde{\Lambda}$ and $\delta\tilde{\Lambda}$ are related to the EOS, we only impose a prior here on $\tilde{\Lambda}$, as $\tilde{\Lambda}$ is measured much better than $\delta\tilde{\Lambda}$ (e.g., Wade et al. 2014). In principle, one could set a combined prior on the three parameters ($\mathcal{M}$, $\tilde{\Lambda}$, $\delta\tilde{\Lambda}$), however, adding this latter parameter would not add significantly to parameter estimation in the relatively low signal-to-noise regime. We therefore leave exploration of physically-motivated priors on $\delta\tilde{\Lambda}$ for future work.

What discussed so far refers to BNS mergers and the corresponding parameterisation does not go to large enough chirp-mass values. Hence, in the case of NSBH mergers we need to derive a new distribution by going back to the raw distributions of progenitor stellar masses and radii, and combine that progenitor distribution with that of a BH of mass M_1 and tidal deformability $\Lambda_1 = 0$. In principle, in this case we could use more sophisticated and realistic mass-ratio distributions, such as those informed by state-of-the-art binary population synthesis (e.g., Broekgaarden et al. 2021). In practice, we have decided to use here a uniformly-distributed mass-ratio prior as we have found that our results do not depend significantly on the choice made for prior distribution on mass ratios. Finally, also for NSBHs we choose to work with priors in $\mathcal{M}-\tilde{\Lambda}$, enforcing $\tilde{\Lambda}(\Lambda_2, \mathcal{M}, q)$ where Λ_2 is the tidal deformability of the neutron star.

2.2. Parameter estimation

We perform a Bayesian parameter-estimation analysis on the GW strain data of the two (likely) BNS merger events GW170817 and GW190425 using the publicly-available data from the GWOSC repository³. We utilise the *Bilby* Bayesian inference framework (Ashton et al. 2019; Romero-Shaw et al. 2020) and priors on source-frame chirp mass and tidal-deformability as discussed in Section 2. Priors for all renaming parameters assume the so-called low-spin ($\chi \leq 0.05$) default priors as defined in *Bilby*. We use Planck 2018 cosmological parameters (Planck Collaboration et al. 2020) such that Hubble’s constant and the matter density are respectively $H_0 = 67.66 \text{ km s}^{-1}$, $\Omega_M = 0.30966$.

Parameter estimation for each run is performed using the *dynesty* nested sampler (Speagle 2020) and the *IMRPhenomPv2_NRTidal* waveform (Dietrich et al.

¹ The quantity $\delta\tilde{\Lambda}$ is another tidal deformability parameter that enters at sixth-order post-Newtonian, cf. $\tilde{\Lambda}$ that enters at fifth order (see Favata 2014; Wade et al. 2014, for definitions of these quantities).

² One could choose a prior on q motivated by astrophysical formation scenarios (e.g.,), however we choose here to do use the common approach instead PL: Are there references for this? If not, I suggest we just delete this footnote.

³ <https://gwosc.org>

2019). Analysis of BNS merger signals is computationally expensive and as such we utilise parallel implementation of Bilby, i.e., pBilby (Smith et al. 2020). More specifically, we reconstruct pressure–density and mass–radius curves from posterior samples of source–frame chirp mass, mass ratio, and effective tidal deformability by performing an EOS inference on the posterior samples of mass and tidal deformability. We utilise a four-piece spectral parameterisation (Lindblom 2010) stitched to an SLy (Douchin & Haensel 2001) crust at low densities (further details can be found in Appendix B). **LR: note important caveat in the paragraph below!** We should remark that the choice of adopting an approach for the EOS inference (i.e., four-piece spectral parameterisation) that is different from that used in the EOS construction (i.e., seven-segment sound-speed parameterisation) follows a precise logic. In particular, this has the advantage that it is possible to compare directly with the LVK inference analysis and hence clearly distinguish the assumptions behind the definition of the priors from those employed in the inference. At the same time, this also implies that the posteriors obtained for the EOSs will be different from the corresponding priors. While these differences are very small, we plan to explore the use of the same set of assumptions for the EOS priors and inference in future work.

3. RESULTS

Figure 2 shows the posterior distributions of the tidal deformability of GW170817 analyzed using the physics-inspired BNS (black), the publicly-released uniform priors from the LIGO-Virgo-KAGRA collaboration (blue; Abbott et al. 2017), and the marginalised prior for $\tilde{\Lambda}$ (red). While we show only the tidal posteriors, we sample over all binary parameters. Posterior distributions for all other binary parameters, including the chirp mass and mass ratio, are identical. Note that, as expected, the marginalised posterior is not significantly different at large tidal deformabilities (compare blue and red lines), but shows significant changes at small values of $\tilde{\Lambda}$, essentially excluding values $\tilde{\Lambda} \lesssim 200$. As a result, the 90% lower and upper limits on the tidal deformability are now set to $226 \leq \tilde{\Lambda}_{1.186} \leq 690$. These arguably represent the most interesting consequence of the physics-informed prior on the $\mathcal{M}-\tilde{\Lambda}$ relation.

Figure 3 shows the impact of the $\mathcal{M}-\tilde{\Lambda}$ relation on the reconstructed posterior distributions of the pressure and energy density (top panel) and of the mass and radius (bottom panel) relative to the GW170817 event. More specifically, the shaded regions show the 90% credible intervals for the posteriors samples obtained using the physics inspired priors (red) and uniform priors (blue). **LR: up to here**

The dotted lines represent the 90% credible intervals of the effective priors on the physics prior and uniform prior when sampled with the spectral parameterisation. The grey




Figure 2. Marginalised posterior distributions of the tidal deformability $\tilde{\Lambda}$ obtained from Bayesian parameter estimation for GW170817. The posterior using a nuclear, microphysics, and astrophysics inspired prior is shown in teal, while the posterior using uniform priors from the LIGO-Virgo-KAGRA collaboration is shown in midnight blue. We show the (marginalised) prior with the light blue curve. Note, the posterior samples from the LVK show support up to $\tilde{\Lambda} = 2000$.

curves show the 90% credible interval of the true EOS distribution of our priors; i.e., the distribution used to generate our chirp-mass–tidal-deformability relationship. We find that the radius of a $1.4 M_{\odot}$ neutron star can be constrained to $12.11^{+0.91}_{-1.11}$ km.

Figure 4 shows the posterior distribution of the intrinsic tidal parameter of GW190425 when analyzed using both the BNS prior (teal) and NSBH prior (pink). We show the official posterior distribution of the LVK obtained using uniform priors in blue. Bayesian-model selection between the BNS and NSBH merger gives a Bayes factor of 1.33, marginally in favour of a BNS merger origin.

Figure 5 shows the reconstructed pressure–density (top) and mass–radius (bottom) posteriors of GW190425. The EOS 90% credible intervals of posteriors obtained using the BNS physics inspired prior are shown in teal, while the NSBH posteriors are shown in pink. Once again, the dashed curve represents our effective prior on the EOS, while the grey curve shows the true prior. We find that the radius of $1.4 M_{\odot}$ neutron star is constrained to $12.69^{+1.24}_{-1.22}$ km using the BNS prior, and $12.06^{+1.44}_{-1.15}$ km using the NSBH prior.

4. CONCLUSIONS

We have constructed new physics inspired priors for the analysis of GW strain data of BNS mergers. Analysis of the neutron star mergers GW170817 and GW190425 with our priors, constrains the effective tidal-deformability of GW170817 to $\tilde{\Lambda} = 384^{+306}_{-158}$ and GW190425 to $\tilde{\Lambda} = 183^{+180}_{-100}$. Corresponding to a 37%, and 80% decrease in the width of 90% credible interval respectively when compared to uniform priors. Performing EOS inference, we find




Figure 3. Posterior distributions of the cold nuclear EOS from analysis of GW170817 using the spectral parameterisation (Lindblom 2010). The top panel shows the posterior distributions in pressure–density space, while the bottom panel shows the reconstructed mass–radius curves. The shaded regions show the 90% credible intervals of the posterior distributions obtained from the analysis using the physics inspired priors (teal) and uniform priors (orange). The dashed curves show the 90% credible intervals of the effective priors, whilst the grey curve shows the 90% credible interval of the true distribution used to generate the priors.




Figure 4. The posterior distributions of the tidal deformability $\tilde{\lambda}$ obtained from Bayesian parameter estimation for G190425. The posterior using a nuclear, microphysics, and astrophysics inspired BNS prior is shown in teal, and the corresponding NSBH prior is shown in pink. The posterior using uniform priors from the LVK is shown in midnight blue.




Figure 5. Posterior distributions of the cold nuclear EOS from analysis of GW190425 using the spectral parameterisation (Lindblom 2010). The top panel shows the posterior distributions in pressure–density space, while the bottom panel shows the reconstructed mass–radius curves. The shaded regions show the 90% credible intervals of the posterior distributions obtained from the analysis using the physics inspired BNS priors (teal) and the NSBH prior (orange). The black dashed curve shows the 90% credible intervals of the effective priors, whilst the grey curve shows the 90% credible interval of the true distribution used to generate the priors.

CE: Same comment as for Fig.3.

that the radius of a $1.4M_{\odot}$ neutron star can be constrained to $12.11^{+0.91}_{-1.11}$ km at a 90% credible interval. Which suggests a preference for softer/stiffer equations of state ...

Model selection between a BNS and NSBH origin on GW190425, yields marginal support for a BNS origin. However, such a result is not unexpected given the low signal to noise of the event and lack of significant tidal measurements.

In this work, we have constructed priors using binaries containing neutron stars with low spins ($\chi \leq 0.05$) only.

APPENDIX

A. MODEL SELECTION

We perform parameter estimation of gravitational-wave strain data using the `dynesty` nested sampler. Unlike standard MCMC samplers, `dynesty` calculates the evidence for each parameter estimation run. To perform Bayesian model selection between BNS and NSBH mergers, we calculate the Bayes factor

$$\text{BF}_{\text{NSBH}}^{\text{BNS}} = \frac{\mathcal{Z}_{\text{BNS}}}{\mathcal{Z}_{\text{NSBH}}}, \quad (\text{A1})$$

where $\mathcal{Z}_{\text{BNS}}$ and $\mathcal{Z}_{\text{NSBH}}$ are the Bayesian evidences obtained for the parameter estimation runs using the BNS and NSBH priors respectively.

Formally, for model selection we should be computing the odds ratio, which is defined as

$$\mathcal{O}_{\text{NSBH}}^{\text{BNS}} := \frac{\mathcal{Z}_{\text{BNS}}}{\mathcal{Z}_{\text{NSBH}}} \frac{\pi_{\text{BNS}}}{\pi_{\text{NSBH}}}, \quad (\text{A2})$$

where π_{BNS} and π_{NSBH} are the prior odds for each hypothesis; i.e our relative belief about which hypothesis is more likely. In this work we set the prior odds to unity, so the odds ratio is the Bayes factor. We could in principle construct an odds ratio where the prior odds was informed by the relative rates of BNS and NSBH mergers. However since these rates are still highly uncertain, we choose the more agnostic approach of setting the prior odds to unity.

B. EQUATION OF STATE INFERENCE

The posterior on the EOS from the GW data is

$$p(\epsilon|d) = \frac{p(d|\epsilon)\pi(\epsilon)}{\mathcal{Z}}, \quad (\text{B3})$$

where $\pi(\epsilon)$ is the prior on the EOS parameters ϵ , and $\mathcal{Z}$ is the evidence. The likelihood is given by

$$p(d|\epsilon) = \int d\theta p(d|\theta) \pi(\theta|\epsilon), \quad (\text{B4})$$

where θ are our nuisance parameters which we marginalise over. Our likelihood reduces to

$$p(d|\epsilon) = \int d\theta p(d|\mathcal{M}, q, \tilde{\Lambda}) \pi(\mathcal{M}, q, \tilde{\Lambda}|\epsilon, \theta) d\mathcal{M} dq d\tilde{\Lambda}. \quad (\text{B5})$$

Since we are dealing with a small number of events (i.e 1-2), we may ignore the modelling of overall neutron-star population parameters without introducing significant biases into the EOS inference (e.g [Agathos et al. 2015](#); [Wysocki et al. 2020](#); [Landry et al. 2020](#)). The marginalised likelihood transforms to

$$p(d|\epsilon) = \int p(d|\mathcal{M}, q, \tilde{\Lambda}) \pi(\mathcal{M}, q, \tilde{\Lambda}|\epsilon) d\mathcal{M} dq d\tilde{\Lambda}, \quad (\text{B6})$$

We have two terms that need to be calculated; $\pi(\mathcal{M}, q, \tilde{\Lambda}|\epsilon)$ a prior on chirp mass, mass-ratio and $\tilde{\Lambda}$ for a given EOS ϵ , and $p(d|\mathcal{M}, q, \tilde{\Lambda})$ the likelihood. Typically, this likelihood would be obtained by reweighing posterior samples of parameter estimation runs

$$p(d|\mathcal{M}, q, \tilde{\Lambda}) = \frac{p(\mathcal{M}, q, \tilde{\Lambda}|d)}{\pi_{\text{PE}}(\mathcal{M}, q, \tilde{\Lambda})}, \quad (\text{B7})$$

however if we did this we would be removing any of our prior information from the posterior samples. Since our prior on $\tilde{\Lambda}$ and $\mathcal{M}$ is actually physical motivated, we set

$$\pi_{\text{PE}}(\mathcal{M}, q, \tilde{\Lambda}) = \pi(\mathcal{M}, q, \tilde{\Lambda}|\epsilon), \quad (\text{B8})$$

which then implies that

$$p(d|\epsilon) = \int p(\mathcal{M}, q, \tilde{\Lambda}|d) d\mathcal{M} dq d\tilde{\Lambda}. \quad (\text{B9})$$

Since the chirp mass is often very-well constrained by parameter estimation, we take the median value of chirp mass $\bar{\mathcal{M}}$, and rewrite our posterior distribution as

$$p(\mathcal{M}, q, \tilde{\Lambda}|d) = p(q, \tilde{\Lambda}) \delta(\mathcal{M} - \bar{\mathcal{M}}), \quad (\text{B10})$$

which simplifies the integral to

$$p(d|\epsilon) = \int p(\bar{\mathcal{M}}, q, \tilde{\Lambda}|d) dq d\tilde{\Lambda}. \quad (\text{B11})$$

We note that we can also express the tidal deformability as a function of $\bar{\mathcal{M}}$, q and ϵ which removes another dimension from our integration

$$p(d|\epsilon) = \int p(q, \tilde{\Lambda}(\bar{\mathcal{M}}, q, \epsilon)|d) dq, \quad (\text{B12})$$

which is now just an integral over mass ratio q .

We evaluate the probability density of our marginalised posterior samples using a bounded kernel-density estimate, and parameterise our EOS using the spectral parameterisation of [Lindblom \(2010\)](#). We use uniform priors for each EOS parameter, and enforce a minimum TOV mass $M_{\text{TOV}} \geq 2.0M_{\odot}$ and causality.

ACKNOWLEDGEMENTS

The authors acknowledges insightful discussions with xxx. Partial funding comes from the ERC Advanced Grant “JETSET: Launching, propagation and emission of relativistic jets from binary mergers and across mass scales” (Grant No. 884631), and the European Horizon Europe staff exchange (SE) programme HORIZON-MSCA2021-SE-01 Grant No. NewFunFiCO-101086251. CE acknowledges support by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) through the CRC-TR 211 “Strong-interaction matter under extreme conditions” – project number 315477589 – TRR 211. LR acknowledges the Walter Greiner Gesellschaft zur Förderung der physikalischen Grundlagenforschung e.V. through the Carl W. Fueck Laureatus Chair.

DATA AVAILABILITY

LR: Here we can add information about the code repository on github

REFERENCES

- Aasi, J., Abbott, B. P., Abbott, R., et al. 2015, *Classical and Quantum Gravity*, 32, 074001, doi: [10.1088/0264-9381/32/7/074001](#)
- Abac, A. G., Abbott, R., Abouelfettouh, I., et al. 2024, *ApJL*, 970, L34, doi: [10.3847/2041-8213/ad5beb](#)
- Abbott, B. P., Abbott, R., Abbott, T. D., et al. 2017, *PhRvL*, 119, 161101, doi: [10.1103/PhysRevLett.119.161101](#)
- , 2018, *PhRvL*, 121, 161101, doi: [10.1103/PhysRevLett.121.161101](#)
- , 2019, *Physical Review X*, 9, 011001, doi: [10.1103/PhysRevX.9.011001](#)
- , 2020, *ApJL*, 892, L3, doi: [10.3847/2041-8213/ab75f5](#)
- Abbott, R., Abbott, T. D., Abraham, S., et al. 2021, *ApJL*, 915, L5, doi: [10.3847/2041-8213/ac082e](#)
- Acernese, F., Agathos, M., Agatsuma, K., et al. 2015, *Classical and Quantum Gravity*, 32, 024001, doi: [10.1088/0264-9381/32/2/024001](#)
- Agathos, M., Meidam, J., Del Pozzo, W., et al. 2015, *PhRvD*, 92, 023012, doi: [10.1103/PhysRevD.92.023012](#)
- Akutsu, T., Ando, M., Arai, K., et al. 2020, *Progress of Theoretical and Experimental Physics*, 2021, 05A101, doi: [10.1093/ptep/ptaa125](#)
- Altıparmak, S., Ecker, C., & Rezzolla, L. 2022a, *ApJL*, 939, L34, doi: [10.3847/2041-8213/ac9b2a](#)
- , 2022b, *Astrophys. J. Lett.*, 939, L34, doi: [10.3847/2041-8213/ac9b2a](#)
- Annala, E., Gorda, T., Kurkela, A., Nättilä, J., & Vuorinen, A. 2020, *Nature Physics*, 16, 907, doi: [10.1038/s41567-020-0914-9](#)
- Antoniadis, J., Freire, P. C. C., Wex, N., et al. 2013, *Science*, 340, 448, doi: [10.1126/science.1233232](#)
- Ashton, G., Hübner, M., Lasky, P. D., et al. 2019, *ApJS*, 241, 27, doi: [10.3847/1538-4365/ab06fc](#)
- Baiotti, L., & Rezzolla, L. 2017, *Rept. Prog. Phys.*, 80, 096901, doi: [10.1088/1361-6633/aa67bb](#)
- Baym, G., Pethick, C., & Sutherland, P. 1971, *ApJ*, 170, 299, doi: [10.1086/151216](#)
- Broekgaarden, F. S., Berger, E., Neijssel, C. J., et al. 2021, *MNRAS*, 508, 5028, doi: [10.1093/mnras/stab2716](#)
- Capano, C. D., Tews, I., Brown, S. M., et al. 2020, *Nature Astronomy*, doi: [10.1038/s41550-020-1014-6](#)
- Cromartie, H. T., Fonseca, E., Ransom, S. M., et al. 2020, *Nature Astronomy*, 4, 72, doi: [10.1038/s41550-019-0880-2](#)
- Dietrich, T., Samajdar, A., Khan, S., et al. 2019, *PhRvD*, 100, 044003, doi: [10.1103/PhysRevD.100.044003](#)

- Douchin, F., & Haensel, P. 2001, *A&A*, 380, 151,
doi: [10.1051/0004-6361:20011402](https://doi.org/10.1051/0004-6361:20011402)
- Ecker, C., & Rezzolla, L. 2022, *Astrophys. J. Lett.*, 939, L35,
doi: [10.3847/2041-8213/ac8674](https://doi.org/10.3847/2041-8213/ac8674)
- Ecker, C., & Rezzolla, L. 2022, *Mon. Not. Roy. Astron. Soc.*, 519,
2615, doi: [10.1093/mnras/stac3755](https://doi.org/10.1093/mnras/stac3755)
- Favata, M. 2014, *PhRvL*, 112, 101101,
doi: [10.1103/PhysRevLett.112.101101](https://doi.org/10.1103/PhysRevLett.112.101101)
- Fonseca, E., Cromartie, H. T., Pennucci, T. T., et al. 2021,
Astrophys. J. Lett., 915, L12, doi: [10.3847/2041-8213/ac03b8](https://doi.org/10.3847/2041-8213/ac03b8)
- Fraga, E. S., Kurkela, A., & Vuorinen, A. 2014, *ApJL*, 781, L25,
doi: [10.1088/2041-8205/781/2/L25](https://doi.org/10.1088/2041-8205/781/2/L25)
- Hebel, K., Lattimer, J. M., Pethick, C. J., & Schwenk, A. 2013,
Astrophys. J., 773, 11, doi: [10.1088/0004-637X/773/1/11](https://doi.org/10.1088/0004-637X/773/1/11)
- Landry, P., Essick, R., & Chatziioannou, K. 2020, *PhRvD*, 101,
123007, doi: [10.1103/PhysRevD.101.123007](https://doi.org/10.1103/PhysRevD.101.123007)
- Lindblom, L. 2010, *PhRvD*, 82, 103011,
doi: [10.1103/PhysRevD.82.103011](https://doi.org/10.1103/PhysRevD.82.103011)
- Miller, M. C., Lamb, F. K., Dittmann, A. J., et al. 2019, *Astrophys. J. Lett.*, 887, L24, doi: [10.3847/2041-8213/ab50c5](https://doi.org/10.3847/2041-8213/ab50c5)
- , 2021, *Astrophys. J. Lett.*, 918, L28,
doi: [10.3847/2041-8213/ac089b](https://doi.org/10.3847/2041-8213/ac089b)
- Most, E. R., Weih, L. R., Rezzolla, L., & Schaffner-Bielich, J. 2018, *Phys. Rev. Lett.*, 120, 261103,
doi: [10.1103/PhysRevLett.120.261103](https://doi.org/10.1103/PhysRevLett.120.261103)
- Paschalidis, V. 2017, *Classical and Quantum Gravity*, 34, 084002,
doi: [10.1088/1361-6382/aa61ce](https://doi.org/10.1088/1361-6382/aa61ce)
- Planck Collaboration, et al. 2020, *A&A*, 641, A6,
doi: [10.1051/0004-6361/201833910](https://doi.org/10.1051/0004-6361/201833910)
- Radice, D., Bernuzzi, S., & Perego, A. 2020, *Annual Review of Nuclear and Particle Science*, 70, 95,
doi: [10.1146/annurev-nucl-013120-114541](https://doi.org/10.1146/annurev-nucl-013120-114541)
- Riley, T. E., Watts, A. L., Bogdanov, S., et al. 2019, *Astrophys. J. Lett.*, 887, L21, doi: [10.3847/2041-8213/ab481c](https://doi.org/10.3847/2041-8213/ab481c)
- Riley, T. E., Watts, A. L., Ray, P. S., et al. 2021, *Astrophys. J. Lett.*, 918, L27, doi: [10.3847/2041-8213/ac0a81](https://doi.org/10.3847/2041-8213/ac0a81)
- Romero-Shaw, I. M., Talbot, C., Biscoveanu, S., et al. 2020,
MNRAS, 499, 3295, doi: [10.1093/mnras/staa2850](https://doi.org/10.1093/mnras/staa2850)
- Smith, R. J. E., Ashton, G., Vajpeyi, A., & Talbot, C. 2020,
MNRAS, 498, 4492, doi: [10.1093/mnras/staa2483](https://doi.org/10.1093/mnras/staa2483)
- Speagle, J. S. 2020, *MNRAS*, 493, 3132,
doi: [10.1093/mnras/staa278](https://doi.org/10.1093/mnras/staa278)
- Wade, L., Creighton, J. D. E., Ochsner, E., et al. 2014, *PhRvD*, 89,
103012, doi: [10.1103/PhysRevD.89.103012](https://doi.org/10.1103/PhysRevD.89.103012)
- Wysocki, D., O’Shaughnessy, R., Wade, L., & Lange, J. 2020,
arXiv e-prints, arXiv:2001.01747,
doi: [10.48550/arXiv.2001.01747](https://doi.org/10.48550/arXiv.2001.01747)