

Critical Review

How Neuroimaging Studies Have Challenged Us to Rethink: Is Chronic Pain a Disease?

Irene Tracey* and M. Catherine Bushnell†

*Oxford Centre for Functional Magnetic Resonance Imaging of the Brain, Nuffield Department of Anaesthetics and Department of Clinical Neurology, University of Oxford, John Radcliffe Hospital, Headington, England, UK.

†Alan Edwards Centre for Research on Pain, Department of Anesthesia and Faculty of Dentistry, McGill University, Montreal, Quebec, Canada.

Abstract: In this review, we present data from functional, structural, and molecular imaging studies in patients and animals supporting the notion that it might be time to reconsider chronic pain as a disease. Across a range of chronic pain conditions, similar observations have been made regarding changes in structure and function within the brains of patients. We discuss these observations within the framework of the current definition of a disease.

Perspective: Neuroimaging studies have made a significant scientific impact in the study of pain. Knowledge of nociceptive processing in the noninjured and injured central nervous system has grown considerably over the past 2 decades. This review examines the information from these functional, structural, and molecular studies within the framework of a disease state.

© 2009 by the American Pain Society

Key words: Chronic pain, disease, neuroimaging, central nervous system.

Editor's Note: This article is 1 in a series of invited Critical Review articles designed to celebrate The Journal of Pain's 10th year anniversary of publication.

Neuroimaging studies have made a huge impact scientifically. The techniques and paradigms are now penetrating the fields of clinical medicine,⁶⁵ diagnosis,¹² and even drug discovery.^{13,88,103}

The pain field is no exception to these exciting developments, and our knowledge of nociceptive processing in the noninjured and injured central nervous system has grown considerably over the past 2 decades. To date, the focus has been to measure functional correlates of the human pain experience using either blood flow based methods, such as Positron Emission Tomography (PET) and functional Magnetic Resonance Imaging (fMRI), or via electrophysiological methods, such as magnetoence-

phalography (MEG) and electroencephalography (EEG). Many excellent reviews and several meta-analyses have been written summarizing the findings to date,^{1,2,16,99} with more recent reviews focusing on the neural basis of pain modulation and its relief.^{10,60,93,100,101} Techniques that focus on the structural architecture of the brain, in terms of gray matter density,^{3,4} white matter connections,^{8,47} receptor density,^{50,56} brain biochemistry,^{41,43} and neurotransmitter availability^{56,91,105,108} have been applied also to the field of pain with often surprising results.

In this review, rather than regurgitate much of the information already reviewed and current regarding human central pain processing, we want to examine the information from these functional, structural, and molecular studies within the framework of a disease state. This is partly motivated by the observation that treatment options are pharmacologically and behaviorally similar for many patients, despite aetiologies for the pain, particularly when neuropathic, being different.^{49,70} This has been taken as evidence that the symptoms likely share some overlapping mechanisms, common to the chronic-pain condition, which the various drugs target irrespective of the cause. Certainly, the focus on mechanism-based analgesic drug development and treatment,^{73,106} reinforces this concept that some shared changes occur during the

Address reprint requests to Irene Tracey, Oxford Centre for Functional Magnetic Resonance Imaging of the Brain, Nuffield Department of Anaesthetics and Department of Clinical Neurology, University of Oxford, John Radcliffe Hospital, Headington, OX3 9DU, England, UK. E-mail: irene@fmrib.ox.ac.uk or M. Catherine Bushnell, Alan Edwards Centre for Research on Pain, Department of Anesthesia and Faculty of Dentistry, McGill University, 3640 University Street, Room M19, Montreal, Quebec, Canada H3A 2B2. E-mail: catherine.bushnell@mcgill.ca

1526-5900/\$36.00

© 2009 by the American Pain Society

doi:10.1016/j.jpain.2009.09.001

transition to a chronic-pain status. The question is whether these changes are detectable using current or advanced techniques, and if so, do the changes that reflect common underlying mechanisms, constitute a disease-like process.

Why is any of this important? Well, chronic pain is an enormous medical-health problem. Current statistics estimate that approximately 20% of the adult population have chronic pain,¹⁴ and separate to the physical and emotional burden it brings, the financial cost to society is huge, currently estimated at over €200 billion per annum in Europe, and \$150 billion per annum in the USA. Treatment options are limited with many patients either not responding or having incomplete pain reduction.^{31,71,96} A paradigm shift in our thinking is needed if we are to better diagnose, manage, and treat chronic pain. Certainly beyond the immediate pain community, we need to encourage people to consider chronic pain in a new light and very possibly as a disease in its own right. This review presents the data, but we leave the reader to decide whether sufficient evidence exists to reclassify chronic pain as a disease.

Distinguishing Disease From Syndrome

According to the Compact Oxford English Dictionary (<http://www.askoxford.com/>), the most common definition of the noun 'disease' is "a disorder of structure or function in a human, animal, or plant, especially one that produces specific symptoms." A more expansive definition includes it being a "cause of discomfort or distress" (Oxford English Dictionary). In contrast, the definition of syndrome is "a group of symptoms which consistently occur together." The main distinction here is that in order for something to be a disease there must be an identifiable disorder of structure or function and not just a grouping of symptoms. The factors leading to the disorder of structure or function might vary, as is the case with cancer, but the end result must be a disordered system. In the case of chronic pain, the disorder would be within the nervous system. Historically, chronic pain has been labeled as a syndrome (or group of syndromes), but recent evidence, mainly from neuroimaging studies, strongly suggests that chronic pain could be labeled as a disease.

Disordered Function Producing Discomfort and Distress: Evidence from Functional Imaging

Chronic pain is discomforting and distressing for most patients. Providing objective proof that this is the case, in addition to listening to the patient or examining their behavior, can be obtained using functional imaging. Areas of the brain involved in processing and controlling affect, negative emotions like depression, anxiety, and aversion, are now better understood and include structures like the amygdala, anterior insular cortex, prefrontal cortices, parahippocampal region, amongst others. Recruitment of these regions could be taken as evidence

for the patient's pain causing discomfort or distress. For instance, the processing of experimental heat pain in patients with somatoform pain disorder compared to matched controls revealed, despite similar behavioral ratings, a hypoactive state of the ventromedial prefrontal/orbitofrontal cortex (BA 10/11) and a hyperactive state of the parahippocampal gyrus, amygdala, and anterior insular.⁴⁵ An earlier study by Gracely et al⁴⁰ on fibromyalgia patients showed that pain catastrophizing, independent of depression, was significantly associated with increased activity in similar brain regions, particularly those associated with attention and anticipation to pain, as well as emotional aspects of pain. Interestingly, such findings are found across different patient types supporting common disturbances in function. For instance, in patients with Irritable Bowel Syndrome (IBS), Mayer et al⁶⁸ found that compared to patients with ulcerative colitis and control subjects, the IBS group had increased activity in response to rectal distention within the amygdala and prefrontal cortices, amongst other limbic and paralimbic regions. Kulkarni et al found that osteoarthritic knee pain was associated with increased activity in the cingulate cortex, thalamus, and the amygdala when compared to experimental knee pain.⁵⁸ Finally, focusing on the neural correlates underpinning the patients' ongoing, tonic pain, Baliki et al⁵ again emphasized the relevance of the medial prefrontal cortex, including rostral ACC, during episodes of sustained, high, ongoing pain. Furthermore, its activity was strongly related to the intensity of chronic back pain.

These findings of an altered cerebral processing of either experimentally induced or disease-related pain in patients support previous findings identifying the relevance of these structures in pain anticipation,^{15,77,79,80} and anxiety-induced pain amplification.^{29,76}

Evidence of disturbed prefrontal activity and a dysfunction of emotion regulation during experimental pain stimulation in depressed patients have been shown in recent studies.^{7,95}

Such data forces us to think about how factors associated with chronic pain conditions, like depression, can become part of the overall condition itself and contribute to the discomfort and distress via increased activity within relevant brain regions. For instance, in patients with fibromyalgia, it has been shown that their degree of depression was related to amygdala and anterior insular activity during experimental pain,³⁸ as well as medial prefrontal cortex activity during disease-relevant induced pain in patients with rheumatoid arthritis and suffering depression.⁹⁰

The anterior insular cortex is particularly intriguing because of its role beyond pain perception. Current thought links activity within the anterior insula to, among other factors, interoception, body awareness, anxiety, depression, fear, and possibly even consciousness.¹⁸⁻²¹ There is therefore a potential link to a sense of body disturbance, discomfort, and distress. In a meta-analysis performed by Schweinhardt et al,⁸⁹ it was shown that the peak coordinate of activity in clinical pain was shifted to the anterior insular cortex compared to nociceptive pain in healthy volunteers, whose activity was more

midposterior insular. This apparent maladaptive plasticity and shift in brain activity towards the more affective division of the insular cortex is perhaps indicative of a functional disturbance.

One consequence of this discomfort and distress is the impact it has on a patient's cognition,²⁸ and is perhaps a direct way of confirming the presence of centrally induced alterations in normal cognitive functioning due to pain. While many neuroimaging studies examine the neural basis for how attention and distraction modulate the pain experience, they have largely been done in healthy control subjects and not patients.¹⁰¹ The reverse has rarely been examined,⁹³ namely identifying a disruption in normal cognitive brain processing due to the presence of pain, except for 1 study in healthy controls.¹¹ Studies looking at how pain alters your capacity to make rational decisions due to biases in cost and reward calculations identify, again in healthy subjects, that pain is clearly disruptive to normal cognition,⁹⁴ and one can readily extrapolate these findings to testable experiments in patients that might produce further evidence to support functions being disturbed.

In summary, these findings strongly support the case for dysfunctional pain processing, especially in affect-regulating regions, and that these patterns of brain activity strongly reflect patients being in true discomfort and distress.

Disordered Functions and Departure from State of Health: Evidence from Functional Imaging

We shall examine 3 areas where normal physiological functions have been shown to be disturbed in chronic pain states, indicating a departure from a state of health: 1) resting state networks; 2) descending inhibition and facilitation; and 3) thalamic asymmetry.

Resting State Networks

Several years ago, it was observed that subjects undergoing neuroimaging data collection while at rest displayed functional connectivity of specific cortical regions,^{32,44} and that this observation was robust across subjects and modalities. These connectivities are now considered as components of the default mode network (DMN), a set of brain regions including medial prefrontal cortex, medial temporal lobes, and posterior cingulate cortex /retrospenial cortex that display balanced positive and negative correlations and are disrupted in several neurological and psychiatric disorders.^{22,32,82,84} In response to task performance, certain areas within the DMN reliably deactivate, and in an early study, we found complete abolishment of normal nociceptive-induced deactivation in the capsaicin model of central sensitization in the presence of gabapentin,⁵⁴ suggesting a possible interaction with the default mode network. Baliki et al⁶ more specifically investigated whether long-term pain alters the functional connectivity of these cortical regions known to be active at rest. During execution of a simple visual task, which patients with chronic back

pain performed as well as controls, they found that patients displayed reduced deactivation in several key default-mode network regions. Their findings demonstrate that chronic pain, like other major neurological and psychiatric diseases, has a widespread impact on overall normal brain function.

Descending Inhibition and Facilitation

The descending pain modulatory system is a well-characterized anatomical network that enables us to regulate, largely within the dorsal horn, nociceptive processing in varying circumstances to produce either facilitation or inhibition.^{30,48,100} The relevance of descending facilitation in chronic-pain states has gathered considerable momentum over the past few years,^{37,78,97} and our human work in models of central sensitization confirm that this facilitatory system becomes active and underpins the maintenance of the centrally sensitized state.^{54,59,107} In parallel, many studies in chronic-pain patients have highlighted also a dysfunction in the normal descending inhibition displayed by healthy volunteers, indicating a dysfunction in this powerful and dedicated endogenous pain modulatory system in chronic pain.^{9,39,46,63,83,92}

Thalamic Asymmetry

Experimentally induced tonic pain has previously been reported to result in less thalamic activation when compared to acute phasic pain in PET studies.²⁵ However, controversy exists, as increased blood flow to the thalamus has also been reported and thought to reflect an arousal reaction to pain,⁷⁵ and to be involved in the processes of attention and vigilance.^{33,81} Evidence from patient studies, however, supports the fact that blood flow to the thalamus is reduced: A common finding has been a relative decrease in thalamic CBF during ongoing pain, which then receded after analgesia and symptom improvement.^{26,34,35,52,53,74} Indeed, historically, thalamic infarcts have long been recognized as a cause of spontaneous pain²⁴ and more recently, atrophy of the thalamus has also been reported in patients with chronic back pain using voxel-based morphometry (see section below).

Therefore, perhaps the most convincing data directly to support the idea that pain is a disease, rather than a syndrome, involves evidence that it is a disorder of structure, as well as function. This is consistent with the definition of disease ("a disorder of structure or function in a human, animal, or plant, especially one that produces specific symptoms.")

Disturbed Structure of the Brain: Evidence from Anatomical MRI

In 2004, Apkarian et al¹³ reported that chronic-pain patients had less brain gray matter than age-matched control subjects. That study was conducted in chronic low-back pain patients and showed that such patients had reduced gray matter in the thalamus and in the lateral prefrontal cortex, a region involved in descending pain modulation. The gray-matter loss was greater in patients who had neuropathic type symptoms than ones who did not, and the

decrease in gray matter correlated with the duration of the symptoms. Similar studies have now been conducted in patients with chronic headache, fibromyalgia and irritable bowel syndrome (IBS), and chronic regional pain syndrome (CRPS).⁶⁶ Although the details of which brain regions show the largest effects differ among studies, gray-matter decreases have been observed in all of these populations.^{23,36,57,61,85,86} The predominant gray-matter decreases observed in patients with chronic pain contrast with usage-related increases in brain gray matter that has been observed during learning,²⁷ and during repeated painful stimulation in healthy subjects.⁹⁸ Similarly, patients with cluster headache who show increased hypothalamic activation have increased gray matter in the region of increased activation.⁶⁷

Another anatomical neuroimaging method, diffusion tensor imaging (DTI), allows in vivo mapping of the anatomical connections in the human brain. Hadjipavlou et al⁴⁷ used this method to identify anatomical circuitry involved in the top-down influence on pain processing, involving the periaqueductal grey (PAG) and its connections with the prefrontal cortex, amygdala, thalamus, and rostroventral medulla. This method has now been applied to chronic-pain patients, where we see disruptions of structure within brain regions involved in normal modulatory influences on pain.^{36,61}

Together, these anatomical studies show that chronic pain is associated with structural changes in the brain. Nevertheless, the current cross-sectional studies tell us little about cause and effect. This is particularly important, since chronic-pain patients frequently have comorbid conditions, including anxiety and mood disorders, altered life-styles so are generally more sedentary, and are also taking various drugs that themselves might be contributing to these measured changes. Thus, it is possible that the gray matter changes are related to the comorbid factors and not to the pain itself. Although some studies have excluded patients with major comorbid conditions such as depression,⁵⁷ other studies suggest that such factors are important to the gray-matter changes. For example, Schmidt-Wilcke et al⁸⁷ observed that when depression and age were included as nuisance factors, most of the observed gray-matter changes were no longer significant. Similarly, we cannot identify from these neuroimaging studies the cellular basis for changes in grey-matter size—is this a neurodegenerative phenomenon, or are the structural changes related to non-neural cells? In order to answer these questions, animal studies are needed. Animal models are available to address a number of chronic pain conditions, including neuropathic pain, headache, CRPS, arthritis, inflammatory visceral pain conditions, and back pain. On the other hand, such models do not completely mimic functional pain conditions, where the etiology is unknown. Nevertheless, the use of rodents with short life spans will allow us to conduct longitudinal studies lasting only months, but covering a large part of the animal's life span. A recent 5-month longitudinal study of rats undergoing a nerve injury (spared nerve injury—SNI) revealed reductions in the size of frontal cortex, but not until 20 weeks after the injury. Although the rats showed mechanical

and thermal hyperalgesia from the time of the injury, they began demonstrating anxiety-like behavior at approximately the same time as the changes in frontal cortex became manifest (Seminowicz et al, in press). Many chronic-pain patients show anxiety-like behavior after their pain has persisted for months or years, so it is interesting to speculate that some of these secondary effects of chronic pain may well be associated with structural changes in the brain.

Rodent studies also allow for histological analysis of the tissue, thus helping us interpret the anatomical changes seen with neuroimaging methods. Metz et al⁶⁹ investigated layer 2/3 pyramidal neurons in acute slices of the medial prefrontal cortex (mPFC) in the rat SNI model of neuropathic pain. These investigators found changes in dendritic branching and spine density of the neurons, providing the first direct evidence of anatomical changes at the cellular basis associated with chronic pain.

Neurochemical Disruptions in the Brain: Evidence from PET Studies

Studies are now beginning to show that chronic-pain patients may have altered brain neurochemistry. Using in vivo proton magnetic-resonance spectrometry (¹H-MRS), Grachev et al⁴² observed altered brain chemistry in the frontal cortices of chronic back-pain patients. Decreased levels of the neuronal marker N-Acetyl aspartate were observed in the dorso-lateral prefrontal cortex, a region in which gray-matter decreases were also observed in back-pain patients. Using similar techniques, Mullins et al⁷² showed that glutamate is elevated in the cingulate cortex in response to painful stimuli in healthy humans, and Harris et al⁵¹ showed that reductions in glutamate in the posterior insula in fibromyalgia patients is associated with reduced experimental and clinical pain. These neurochemical findings add further evidence to the idea that reduced gray-matter density in chronic-pain patients may be related to possible excitotoxicity and neuronal loss.

Other studies show possible changes in neurochemicals involved in pain modulation in chronic-pain patients. Two seminal molecular-imaging studies using positron emission tomography (PET) in chronic-pain patients showed cerebral decreases in opioid receptor binding in patients with central neuropathic pain and with rheumatoid arthritis.^{55,56} More recent PET studies in fibromyalgia patients show alterations in both dopamine and opioid availability in the forebrain.^{50,104} In the absence of external painful stimuli, fibromyalgia patients appear to have a reduction in the receptor availability of both dopamine D2 receptors and opioid mu-receptors in parts of the forebrain. For dopamine, it has also been shown that patients do not release dopamine in the basal ganglia in response to an external pain stimulus, whereas healthy subjects do release dopamine in that situation.^{91,105} Such results may mean that patients have decreased receptor availability or have a heightened background tone or endogenous release of these neurotransmitters that is known to produce a reduced phasic release, but in either case, it appears that neurochemicals important for pain modulation are not responding as

they do in healthy individuals. Again, these findings are not specific to fibromyalgia but are shown for other chronic-pain conditions. Willoch et al¹⁰² found in patients with central poststroke pain reduced opioid binding in pain-processing regions and Maarrawi et al⁶² showed differential opioid-receptor availability in central and peripheral neuropathic-pain patients. Combined, these studies strongly support the case for disorder or dysfunction in the neurochemistry of chronic-pain patients' brains.

Does the Evidence Prove that Chronic Pain is a Disease?

This review has presented substantial functional, anatomical, and neurochemical evidence that chronic-pain patients have altered brains. But is what we see as altered and dysfunctional central nervous system processing more an adaptive response to the constant nociceptive barrage rather than a diseaselike process? The chicken and egg problem here is that for most "diseases" normally something within the body alters and changes function and possibly structure, and this process itself largely produces the symptoms and condition. For pain, we only have evidence of such changes after the transition to chronicity; therefore, it's difficult to know whether these mechanism-based changes are simply a normal adaptive response or are critical for the chronic-pain state itself. Reversibility of such changes with symptom improvement might help clarify this issue. However, if these changes could be induced without any initiating nociceptive input, would a chronic-pain-like state occur? In many conditions, chronic pain results

after a clear tissue-damaging event, leading first to acute pain and then to chronic pain. Most neuropathic pain conditions have a clear nerve injury that precipitated the pain. Nevertheless, other painful conditions come about without a clear precipitating injury. These conditions, such as fibromyalgia, vulvodynia, interstitial cystitis, and irritable bowel syndrome, are sometimes referred to as functional pain syndromes, because the patient presents with pain without an obvious physiological cause. Could these conditions be related to pathophysiology of the central nervous system that is similar to that caused by a constant nociceptive barrage in other pain conditions? Recent data from animal studies investigating stress-induced hyperalgesia⁶⁴ and peripheral hypersensitivity without peripheral inflammation after amygdala activation¹⁷ provide support for this concept. Could an excitotoxicity of pain-modulatory circuitry be evoked not only by hyperexcitability of the afferent nociceptive system, but also, given the right genetic susceptibility, by activation of the stress, arousal, or attentional circuitry in humans? At this time, we can only speculate about these mechanisms.

Conclusion

By taking a multifactorial and longitudinal approach to the study of chronic pain, including in our analyses genetic and environmental factors, and merging data from the molecular to the clinical level, we may someday unravel the complexities of chronic pain. But for now, imaging studies have shown that chronic pain is associated with functional, structural, and chemical changes in the brain, thus putting it into the realm of a disease state.

References

1. Apkarian AV, Baliki MN, Geha PY: Towards a theory of chronic pain. *Prog Neurobiol* 87:81-97, 2009
2. Apkarian AV, Bushnell MC, Treede RD, Zubieta JK: Human brain mechanisms of pain perception and regulation in health and disease. *Eur J Pain* 9:463-484, 2005
3. Apkarian AV, Sosa Y, Sonty S, Levy RM, Harden RN, Parrish TB, Gitelman DR: Chronic back pain is associated with decreased prefrontal and thalamic gray matter density. *J Neurosci* 24:10410-10415, 2004
4. Ashburner J, Friston KJ: Voxel-based morphometry—the methods. *Neuroimage* 11:805-821, 2000
5. Baliki MN, Chialvo DR, Geha PY, Levy RM, Harden RN, Parrish TB, Apkarian AV: Chronic pain and the emotional brain: Specific brain activity associated with spontaneous fluctuations of intensity of chronic back pain. *J Neurosci* 26:12165-12173, 2006
6. Baliki MN, Geha PY, Apkarian AV, Chialvo DR: Beyond feeling: Chronic pain hurts the brain, disrupting the default-mode network dynamics. *J Neurosci* 28:1398-1403, 2008
7. Bär KJ, Wagner G, Koschke M, Boettger S, Boettger MK, Schlösser R, Sauer H: Increased prefrontal activation during pain perception in major depression. *Biological psychiatry* 62:1281-1287, 2007
8. Behrens TE, Johansen-Berg H, Woolrich MW, Smith SM, Wheeler-Kingshott CA, Boulby PA, Barker GJ, Sillery EL, Sheehan K, Ciccarelli O, Thompson AJ, Brady JM, Matthews PM: Non-invasive mapping of connections between human thalamus and cortex using diffusion imaging. *Nat Neurosci* 6:750-757, 2003
9. Berman SM, Naliboff BD, Suyenobu B, Labus JS, Stains J, Ohning G, Kilpatrick L, Bueller JA, Ruby K, Jarcho J, Mayer EA: Reduced brainstem inhibition during anticipated pelvic visceral pain correlates with enhanced brain response to the visceral stimulus in women with irritable bowel syndrome. *J Neurosci* 28:349-359, 2008
10. Bingel U, Tracey I: Imaging CNS modulation of pain in humans. *Physiology (Bethesda)* 23:371-380, 2008
11. Bingel U, Rose M, Glascher J, Buchel C: fMRI reveals how pain modulates visual object processing in the ventral visual stream. *Neuron* 55:157-167, 2007
12. Bookheimer S: Pre-surgical language mapping with functional magnetic resonance imaging. *Neuropsychol Rev* 17:145-155, 2007
13. Borsook D, Becerra L, Hargreaves R: A role for fMRI in optimizing CNS drug development. *Nat Rev Drug Discov* 5:411-424, 2006

14. Breivik H, Collett B, Ventafridda V, Cohen R, Gallacher D: Survey of chronic pain in Europe: Prevalence, impact on daily life, and treatment. *Eur J Pain* 10:287-333, 2006
15. Brown CA, Seymour B, El-Deredy W, Jones AK: Confidence in beliefs about pain predicts expectancy effects on pain perception and anticipatory processing in right anterior insula. *Pain* 139:324-332, 2008
16. Bushnell M, Apkarian AV: Representation of pain in the brain. *Wall and Melzack's Textbook of Pain* 107-124 2006, 2006
17. Carrasquillo Y, Gereau RW 4th: Activation of the extracellular signal-regulated kinase in the amygdala modulates pain perception. *J Neurosci* 27:1543-1551, 2007
18. Craig AD: Interoception: The sense of the physiological condition of the body. *Curr Opin Neurobiol* 13:500-505, 2003
19. Craig AD: How do you feel? Interoception: the sense of the physiological condition of the body. *Nature Rev Neurosci* 3:655-666, 2002
20. Craig AD: How do you feel—now? The anterior insula and human awareness. *Nature Rev Neurosci* 10:59-70, 2009
21. Critchley HD, Wiens S, Rotshtein P, Ohman A, Dolan RJ: Neural systems supporting interoceptive awareness. *Nat Neurosci* 7:189-195, 2004
22. Damoiseaux JS, Beckmann CF, Arigita EJ, Barkhof F, Scheltens P, Stam CJ, Smith SM, Rombouts SA: Reduced resting-state brain activity in the "default network" in normal aging. *Cereb Cortex* 18:1856-1864, 2008
23. Davis KD, Pope G, Chen J, Kwan CL, Crawley AP, Diamant NE: Cortical thinning in IBS: Implications for homeostatic, attention, and pain processing. *Neurology* 70:153-154, 2008
24. Dejerine, J: Le Syndrome Thalamique. *Rev Neurol* 14, 1906
25. Derbyshire SW, Jones AK: Cerebral responses to a continual tonic pain stimulus measured using positron emission tomography. *Pain* 76:127-135, 1998
26. Di Piero V, Jones AK, Iannotti F, Powell M, Perani D, Lenzi GL, Frackowiak RS: Chronic pain: A PET study of the central effects of percutaneous high cervical cordotomy. *Pain* 46:9-12, 1991
27. Draganski B, Gaser C, Kempermann G, Kuhn HG, Winkler J, Büchel C, May A: Temporal and spatial dynamics of brain structure changes during extensive learning. *J Neurosci* 26:6314-6317, 2006
28. Eccleston C, Crombez G: Pain demands attention: a cognitive-affective model of the interruptive function of pain. *Psychol Bull* 125:356-366, 1999
29. Fairhurst M, Wiech K, Dunckley P, Tracey I: Anticipatory brainstem activity predicts neural processing of pain in humans. *Pain* 128:101-110, 2007
30. Fields H: In P. Melzack P, Wall, R (eds): *Textbook of Pain*, 4th ed. Churchill Livingstone, 2005, pp 125-142
31. Finnerup NB, Otto M, McQuay HJ, Jensen TS, Sindrup SH: Algorithm for neuropathic pain treatment: An evidence based proposal. *Pain* 118:289-305, 2005
32. Fox MD, Raichle ME: Spontaneous fluctuations in brain activity observed with functional magnetic resonance imaging. *Nat Rev Neurosci* 8:700-711, 2007
33. Fredrikson M, Wik G, Fischer H, Andersson J: Affective and attentive neural networks in humans: A PET study of Pavlovian conditioning. *Neuroreport* 7:97-101, 1995
34. Garcia-Larrea L, Maarrawi J, Peyron R, Costes N, Mertens P, Magnin M, Laurent B: On the relation between sensory deafferentation, pain and thalamic activity in Wallenberg's syndrome: A PET-scan study before and after motor cortex stimulation. *Eur J Pain* 10:677-688, 2006
35. García-Larrea L, Peyron R, Mertens P, Gregoire MC, Lavenne F, Le Bars D, Convers P, Mauguière F, Sindou M, Laurent B: Electrical stimulation of motor cortex for pain control: A combined PET-scan and electrophysiological study. *Pain* 83:259-273, 1999
36. Geha PY, Baliki MN, Harden RN, Bauer WR, Parrish TB, Apkarian AV: The brain in chronic CRPS pain: Abnormal gray-white matter interactions in emotional and autonomic regions. *Neuron* 60:570-581, 2008
37. Gebhart GF: Descending modulation of pain. *Neurosci Biobehav Rev* 27:729-737, 2004
38. Giesecke T, Gracely RH, Williams DA, Geisser ME, Petzke FW, Clauw DJ: The relationship between depression, clinical pain, and experimental pain in a chronic pain cohort. *Arthritis Rheum* 52:1577-1584, 2005
39. Goadsby PJ: Recent advances in understanding migraine mechanisms, molecules and therapeutics. *Trends Mol Med* 13:39-44, 2007
40. Gracely RH, Geisser ME, Giesecke T, Grant MA, Petzke F, Williams DA, Clauw DJ: Pain catastrophizing and neural responses to pain among persons with fibromyalgia. *Brain* 127:835-843, 2004
41. Grachev ID, Fredrickson BE, Apkarian AV: Abnormal brain chemistry in chronic back pain: An in vivo proton magnetic resonance spectroscopy study. *Pain* 89:7-18, 2000
42. Grachev ID, Fredrickson BE, Apkarian AV: Brain chemistry reflects dual states of pain and anxiety in chronic low back pain. *J Neural Transm* 109:1309-1334, 2002
43. Grachev ID, Fredrickson BE, Apkarian AV: Dissociating anxiety from pain: Mapping the neuronal marker N-acetyl aspartate to perception distinguishes closely interrelated characteristics of chronic pain. *Mol Psychiatry* 6:256-258, 2001
44. Greicius MD, Krasnow B, Reiss AL, Menon V: Functional connectivity in the resting brain: A network analysis of the default mode hypothesis. *Proc Natl Acad Sci USA* 100:253-258, 2003
45. Gündel H, Valet M, Sorg C, Huber D, Zimmer C, Sprenger T, Tölle TR: Altered cerebral response to noxious heat stimulation in patients with somatoform pain disorder. *Pain* 137:413-421, 2008
46. Gwilym SE, Keltner JR, Warnaby CE, Carr AJ, Chizh B, Chessell I, Tracey I: Psychophysical and functioning imaging evidence supporting the presence of central sensitization in a cohort of osteoarthritis patients. *Arthritis Rheum* 61:1226-1234, 2009
47. Hadjipavlou G, Dunckley P, Behrens E, Tracey I: Determining anatomical connectivities between cortical and brainstem pain processing regions in humans: A diffusion tensor imaging study in healthy controls. *Pain* 123:169-178, 2006

48. Hagbarth KE, Kerr DI: Central influences on spinal afferent conduction. *J Neurophysiol* 17:295-307, 1954
49. Hansson PT, Dickenson AH: Pharmacological treatment of peripheral neuropathic pain conditions based on shared commonalities despite multiple etiologies. *Pain* 113: 251-254, 2005
50. Harris RE, Clauw DJ, Scott DJ, McLean SA, Gracely RH, Zubieta JK: Decreased central mu-opioid receptor availability in fibromyalgia. *J Neurosci* 27:10000-10006, 2007
51. Harris RE, Sundgren PC, Pang Y, Hsu M, Petrou M, Kim SH, McLean SA, Gracely RH, Clauw DJ: Dynamic levels of glutamate within the insula are associated with improvements in multiple pain domains in fibromyalgia. *Arthritis Rheum* 58:903-907, 2008
52. Hsieh JC, Belfrage M, Stone-Elander S, Hansson P, Ingvar M: Central representation of chronic ongoing neuropathic pain studied by positron emission tomography. *Pain* 63:225-236, 1995
53. Iadarola MJ, Max MB, Berman KF, Byas-Smith MG, Coghill RC, Gracely RH, Bennett GJ: Unilateral decrease in thalamic activity observed with positron emission tomography in patients with chronic neuropathic pain. *Pain* 63: 55-64, 1995
54. Iannetti GD, Zambreanu L, Wise RG, Buchanan TJ, Hugdins JP, Smart TS, Vennart W, Tracey I: Pharmacological modulation of pain-related brain activity during normal and central sensitization states in humans. *Proc Natl Acad Sci USA* 102:18195-18200, 2005
55. Jones AK, Cunningham VJ, Ha-Kawa S, Fujiwara T, Luthra SK, Silva S, Derbyshire S, Jones T: Changes in central opioid receptor binding in relation to inflammation and pain in patients with rheumatoid arthritis. *Br J Rheumatol* 33:909-916, 1994
56. Jones AK, Watabe H, Cunningham VJ, Jones T: Cerebral decreases in opioid receptor binding in patients with central neuropathic pain measured by [¹¹C]diprenorphine binding and PET. *Eur J Pain* 8:479-485, 2004
57. Kuchinad A, Schweinhardt P, Seminowicz DA, Wood PB, Chizh BA, Bushnell MC: Accelerated brain gray matter loss in fibromyalgia patients: Premature aging of the brain? *J Neurosci* 27:4004-4007, 2007
58. Kulkarni B, Bentley DE, Elliott R, Julyan PJ, Boger E, Watson A, Boyle Y, El-Deredy W, Jones AK: Arthritic pain is processed in brain areas concerned with emotions and fear. *Arthritis Rheum* 56:1345-1354, 2007
59. Lee MC, Zambreanu L, Menon DK, Tracey I: Identifying brain activity specifically related to the maintenance and perceptual consequence of central sensitization in humans. *J Neurosci* 28:11642-11649, 2008
60. Leknes S, Tracey I: A common neurobiology for pain and pleasure. *Nat Rev Neurosci* 9:314-320, 2008
61. Lutz J, Jäger L, de Quervain D, Krauseneck T, Padberg F, Wichnalek M, Beyer A, Stahl R, Zirngibl B, Morhard D, Reiser M, Schelling G: White and gray matter abnormalities in the brain of patients with fibromyalgia: A diffusion-tensor and volumetric imaging study. *Arthritis Rheum* 58: 3960-3969, 2008
62. Maarrawi J, Peyron R, Mertens P, Costes N, Magnin M, Sindou M, Laurent B, Garcia-Larrea L: Differential brain opioid receptor availability in central and peripheral neuropathic pain. *Pain* 127:183-194, 2007
63. Mainero C, Zhang WT, Kumar A, Rosen BR, Sorensen AG: Mapping the spinal and supraspinal pathways of dynamic mechanical allodynia in the human trigeminal system using cardiac-gated fMRI. *Neuroimage* 35:1201-1210, 2007
64. Martenson ME, Cetas JS, Heinricher MM: A possible neural basis for stress-induced hyperalgesia. *Pain* 142:236-244, 2009
65. Matthews PM, Honey GD, Bullmore ET: Applications of fMRI in translational medicine and clinical practice. *Nat Rev Neurosci* 7:732-744, 2006
66. May A: Chronic pain may change the structure of the brain. *Pain* 137:7-15, 2008
67. May A, Ashburner J, Büchel C, McGonigle DJ, Friston KJ, Frackowiak RS, Goadsby PJ: Correlation between structural and functional changes in brain in an idiopathic headache syndrome. *Nat Med* 5:836-838, 1999
68. Mayer EA, Berman S, Suyenobu B, Labus J, Mandelkern MA, Naliboff BD, Chang L: Differences in brain responses to visceral pain between patients with irritable bowel syndrome and ulcerative colitis. *Pain* 115:398-409, 2005
69. Metz AE, Yau HJ, Centeno MV, Apkarian AV, Martina M: Morphological and functional reorganization of rat medial prefrontal cortex in neuropathic pain. *Proc Natl Acad Sci USA* 106:2423-2428, 2009
70. Molton IR, Graham C, Stoelb BL, Jensen MP: Current psychological approaches to the management of chronic pain. *Curr Opin Anaesthesiol* 20:485-489, 2007
71. Moore RA, Moore OA, Derry S, McQuay HJ: Numbers needed to treat calculated from responder rates give a better indication of efficacy in osteoarthritis trials than mean pain scores. *Arthritis Res Ther* 10:R39, 2008
72. Mullins PG, Rowland LM, Jung RE, Sibbitt WL Jr: A novel technique to study the brain's response to pain: Proton magnetic resonance spectroscopy. *Neuroimage* 26:642-646, 2005
73. Patapoutian A, Tate S, Woolf CJ: Transient receptor potential channels: Targeting pain at the source. *Nat Rev Drug Discov* 8:55-68, 2009
74. Peyron R, Garcia-Larrea L, Deiber MP, Cinotti L, Convers P, Sindou M, Mauguière F, Laurent B: Electrical stimulation of precentral cortical area in the treatment of central pain: Electrophysiological and PET study. *Pain* 62: 275-286, 1995
75. Peyron R, García-Larrea L, Grégoire MC, Costes N, Convers P, Lavenne F, Mauguière F, Michel D, Laurent B: Haemodynamic brain responses to acute pain in humans: sensory and attentional networks. *Brain* 122:1765-1780, 1999
76. Ploghaus A, Narain C, Beckmann CF, Clare S, Bantick S, Wise R, Matthews PM, Rawlins JN, Tracey I: Exacerbation of pain by anxiety is associated with activity in a hippocampal network. *J Neurosci* 21:9896-9903, 2001
77. Ploghaus A, Tracey I, Gati JS, Clare S, Menon RS, Matthews PM, Rawlins JN: Dissociating pain from its anticipation in the human brain. *Science* 284:1979-1981, 1999
78. Porreca F, Ossipov MH, Gebhart GF: Chronic pain and medullary descending facilitation. *Trends Neurosci* 25: 319-325, 2002
79. Porro CA, Baraldi P, Pagnoni G, Serafini M, Facchin P, Maieron M, Nichelli P: Does anticipation of pain affect cortical nociceptive systems? *J Neurosci* 22:3206-3214, 2002

80. Porro CA, Cettolo V, Francescato MP, Baraldi P: Functional activity mapping of the mesial hemispheric wall during anticipation of pain. *Neuroimage* 19:1738-1747, 2003
81. Portas CM, Rees G, Howseman AM, Josephs O, Turner R, Frith CD: A specific role for the thalamus in mediating the interaction of attention and arousal in humans. *J Neurosci* 18: 8979-8989, 1998
82. Raichle ME, Snyder AZ: A default mode of brain function: A brief history of an evolving idea. *Neuroimage* 37: 1083-1090. discussion 1097-1089.
83. Sandrini G, Rossi P, Milanov I, Serrao M, Cecchini AP, Nappi G: Abnormal modulatory influence of diffuse noxious inhibitory controls in migraine and chronic tension-type headache patients. *Cephalalgia* 26:782-789, 2006
84. Sheline YI, Barch DM, Price JL, Rundle MM, Vaishnavi SN, Snyder AZ, Mintun MA, Wang S, Coalson RS, Raichle ME: The default mode network and self-referential processes in depression. *Proc Natl Acad Sci USA* 106:1942-1947, 2009
85. Schmidt-Wilcke T, Leinisch E, Gänssbauer S, Draganski B, Bogdahn U, Altmeyen J, May A: Affective components and intensity of pain correlate with structural differences in gray matter in chronic back pain patients. *Pain* 125:89-97, 2006
86. Schmidt-Wilcke T, Leinisch E, Straube A, Kämpfe N, Draganski B, Diener HC, Bogdahn U, May A: Gray matter decrease in patients with chronic tension type headache. *Neurology* 65:1483-1486, 2005
87. Schmidt-Wilcke T, Luerding R, Weigand T, Jürgens T, Schuierer G, Leinisch E, Bogdahn U: Striatal grey matter increase in patients suffering from fibromyalgia—a voxel-based morphometry study. *Pain* 132(Suppl 1):S109-S116, 2007
88. Schweinhardt P, Bountra C, Tracey I: Pharmacological fMRI in the development of new analgesic compounds. *NMR Biomed* 19:702-711, 2006
89. Schweinhardt P, Glynn C, Brooks J, McQuay H, Jack T, Chessell I, Bountra C, Tracey I: An fMRI study of cerebral processing of brush-evoked allodynia in neuropathic pain patients. *Neuroimage* 32:256-265, 2006
90. Schweinhardt P, Kalk N, Wartolowska K, Chessell I, Wordsworth P, Tracey I: Investigation into the neural correlates of emotional augmentation of clinical pain. *Neuroimage* 40:759-766, 2008
91. Scott DJ, Heitzeg MM, Koeppe RA, Stohler, Zubieta JK: Variations in the human pain stress experience mediated by ventral and dorsal basal ganglia dopamine activity. *J Neurosci* 26:10789-10795, 2006
92. Seifert F, Maihofner C: Representation of cold allodynia in the human brain—a functional MRI study. *Neuroimage* 35:1168-1180, 2007
93. Seminowicz DA, Davis KD: A re-examination of pain-cognition interactions: Implications for neuroimaging. *Pain* 130:8-13, 2007
94. Seymour B, O'Doherty JP, Koltzenburg M, Wiech K, Frackowiak R, Friston K, Dolan R: Opponent appetitive-aversive neural processes underlie predictive learning of pain relief. *Nat Neurosci* 8:1234-1240, 2005
95. Strigo IA, Simmons AN, Matthews SC, Craig AD, Paulus MP: Association of major depressive disorder with altered functional brain response during anticipation and processing of heat pain. *Arch Gen Psychiatry* 65:1275-1284, 2008
96. Sultan A, Gaskell H, Derry S, Moore RA: Duloxetine for painful diabetic neuropathy and fibromyalgia pain: Systematic review of randomised trials. *BMC Neurol* 8:29, 2008
97. Suzuki R, Rygh LJ, Dickenson AH: Bad news from the brain: Descending 5-HT pathways that control spinal pain processing. *Trends Pharmacol Sci* 25:613-617, 2004
98. Teutsch S, Herken W, Bingel U, Schoell E, May A: Changes in brain gray matter due to repetitive painful stimulation. *Neuroimage* 42:845-849, 2008
99. Tracey I: Nociceptive processing in the human brain. *Current opinion in neurobiology* 15:478-487, 2005
100. Tracey I, Mantyh PW: The cerebral signature for pain perception and its modulation. *Neuron* 55:377-391, 2007
101. Wiech K, Ploner M, Tracey I: Neurocognitive aspects of pain perception. *Trends Cogn Sci* 12:306-313, 2008
102. Willoch F, Schindler F, Wester HJ, Empl M, Straube A, Schwaiger M, Conrad B, Tölle TR: Central poststroke pain and reduced opioid receptor binding within pain processing circuitries: A [11C]diprenorphine PET study. *Pain* 108: 213-220, 2004
103. Wise RG, Tracey I: The role of fMRI in drug discovery. *J Magn Reson Imaging* 23:862-876, 2006
104. Wood PB, Patterson JC 2nd, Sunderland JJ, Tainter KH, Glabus MF, Lilien DL: Reduced presynaptic dopamine activity in fibromyalgia syndrome demonstrated with positron emission tomography: A pilot study. *J Pain* 8:51-58, 2007
105. Wood PB, Schweinhardt P, Jaeger E, Dagher A, Hakyemez H, Rabiner EA, Bushnell MC, Chizh BA: Fibromyalgia patients show an abnormal dopamine response to pain. *Eur J Neurosci* 25:3576-3582, 2007
106. Woolf CJ: Pain: Moving from symptom control toward mechanism-specific pharmacologic management. *Ann Intern Med* 140:441-451, 2004
107. Zambreanu L, Wise RG, Brooks JC, Iannetti GD, Tracey I: A role for the brainstem in central sensitisation in humans. Evidence from functional magnetic resonance imaging. *Pain* 114:397-407, 2005
108. Zubieta JK, Heitzeg MM, Smith YR, Bueller JA, Xu K, Xu Y, Koeppe RA, Stohler CS, Goldman D: COMT val158met genotype affects mu-opioid neurotransmitter responses to a pain stressor. *Science* 299:1240-1243, 2003