

Supplementary Information

Table S1 | Dominant ring- and diffuse-porous species at the Smithsonian Conservation Biology Institute (SCBI) and Harvard Forest, along with sample sizes included in this analysis.

site	xylem porosity	species	species code	dendrometer bands		tree cores	
				n trees	n tree-years	n cores	date range
SCBI	ring	<i>Carya cordiformis</i>	CACO	0	0	18	1917-2009
		<i>Carya glabra</i>	CAGL	0	0	39	1901-2009
		<i>Carya ovalis</i>	CAOVL	0	0	24	1896-2009
		<i>Carya tomentosa</i>	CATO	0	0	17	1926-2009
		<i>Fraxinus americana</i>	FRAM	0	0	69	1910-2009
		<i>Quercus alba</i>	QURU	34	197	66	1904-2009
		<i>Quercus montana</i>	QUPR	0	0	67	1893-2009
		<i>Quercus rubra</i>	QUAL	35	229	71	1870-2009
		<i>Quercus velutina</i>	QUVE	0	0	83	1902-2009
	diffuse	<i>Fagus grandifolia</i>	FAGR	13	89	81	1932-2009
		<i>Liriodendron tulipifera</i>	LITU	41	354	109	1920-2009
Harvard	ring	<i>Fraxinus americana</i>	FRAM	9	27	0	1901-2014
		<i>Quercus alba</i>	QURU	118	575	179	
		<i>Quercus velutina</i>	QUVE	11	50	0	
	diffuse	<i>Fagus grandifolia</i>	FAGR	8	45	0	1952-2013
		<i>Betula lenta</i>	BELE	8	44	0	
		<i>Betula populifolia</i>	BEPO	5	24	0	
		<i>Betula papyrifera</i>	BEPA	3	13	0	
		<i>Betula alleghaniensis</i>	BEAL	21	90	44	
		<i>Prunus serotina</i>	PRSE	9	37	0	
		<i>Acer rubrum</i>	ACRU	144	669	59	
		<i>Acer pensylvanicum</i>	ACPE	4	16	0	

Table S2 | Sample size by year

Year	SCBI	Harvard Forest
1998	NA	755
1999	NA	733
2000	NA	711
2001	NA	704
2002	NA	701
2003	NA	700
2011	105	NA
2012	99	NA
2013	145	NA
2014	146	NA
2015	144	NA
2016	145	NA
2017	145	NA
2018	143	NA
2019	142	NA

Table S3 | Summary of parameters describing the phenology and rate of growth for ring- and diffuse- porous species at SCBI and Harvard Forest.

	SCBI		Harvard Forest	
	ring	diffuse	ring	diffuse
Stem Growth				
critical T_{max} window	3/22-4/9	2/19-5/21	4/2-5/7	3/19-5/7
DOY_{25}	123 (May 4)	154 (June 4)	132 (May 15)	164 (June 14)
DOY_{50}	152 (June 2)	172 (June 22)	159 (June 9)	182 (July 2)
DOY_{75}	180 (June 30)	190 (July 9)	186 (July 6)	199 (July 19)
$DOY_{g_{max}}$	152 (June 2)	173 (June 23)	161 (June 11)	183 (July 3)
g_{max} (mm/day)	0.046	0.061	0.03	0.025
L_{pgs}	56.5	35.8	54.5	35.1
ΔDBH (mm/yr)	4.7	3.6	3.1	1.4
Leaf Phenology (ecosystem level)				
Greenup	101 (April 11)		115 (April 25)	
Mid-greenup	120 (April 30)		137 (May 17)	
Peak	173 (June 22)		182 (July 1)	
Senescence	215 (Aug. 3)		218 (Aug. 6)	

(placeholder for chronologies table.)






Figure S1 | Growth trajectories for ring- and diffuse-porous trees, as both relative and cumulative fractions of total annual growth. Each line represents one year's growth for a given tree, fit with McMahon model.

\begin{figure}



\caption{Figure S2 | Landscapes of relationships between the day of year on which 25% of annual growth is achieved (DOY_{25}) and temperature in prior weeks for ring- and diffuse-porous trees at SCBI and Harvard Forest. Shown are matrices of linear coefficients of first-order linear regressions between temperature and DOY_{25} , where Window Open and Window Close indicate number of weeks prior to DOY_{25} (ring-porous: May 5 at SCBI, May 13 at HF; diffuse-porous: June 4 at SCBI, June 14 at HF). Black circles indicate the critical T_{max} window (ring-porous: March 22- April 9 at SCBI, April 2 - May 07 at HF; diffuse-porous: Feb. 19- May 21 at SCBI, March 19 - May 07 at HF).}

\begin{figure}[H]



{

}

\caption{Figure S3 | Sensitivity of annual growth, as derived from tree-rings, to monthly minimum temperatures, for ## chronologies from ## sites across eastern North America. Chronologies are grouped by xylem porosity and ordered by mean April T_{max} . Chronology details are given in Table S4. (**NOTE: Figure still needs some work. Chronology numbers are off. See GitHub Issue #49.**)}