

roseatespoonbillIslandPresent model summary

Contents

Overview	2
Variables:	2
Components:	3
Model equation:	3
Global sensitivity and uncertainty analysis:	3
Model uncertainty	3
Model sensitivity	5
Original model	6
Arithmetic mean model	7
Geometric mean model	8
Limiting factor model	9
Multiplicative model	10
Summary of influential variables	11
Original model	11
Arithmetic mean model	11
Geometric mean model	11
Limiting factor model	11
Multiplicative model	12
References	12

Overview

This document summarizes the results of a global sensitivity and uncertainty analysis for the **roseatespoonbillIslandPresent** habitat suitability index (HSI) model for *Platalea ajaja*. Metadata for the model is stored in the `ecorest` package in R.

The original documentation for this model can be found [here](#)¹.

Sub-model: **Island, key or islet habitats evaluated under present conditions**

The `roseatespoonbillIslandPresent` model is comprised of **4** variables and **1** components.

Variables:

Table 1. SIV variables included in the `roseatespoonbillIslandPresent` model. Type indicates whether a variable is continuous or categorical and breakpoints indicates the number of distinct breakpoints in suitability graphs.

	Variable name	Type	Breakpoints
SIV1	dist.to.mainland.SIV	continuous	3
SIV2	island.sa.SIV	categorical	3
SIV3	h.woody.veg.island.SIV	continuous	4
SIV4	roost.nest.hab.SIV	categorical	3




Figure 1. Suitability index graphs for variables included in the `roseatespoonbillIslandPresent` model in `ecorest`.

¹<https://ecolibrary.sec.usace.army.mil/resource/e1901cf8-7e92-428c-a955-f6b6594e4d0b>

Components:

Table 2. Components included in the roseatespoonbillIslandPresent model in ecorest.

Component	Equation
Roosting and nesting component	$(SIV1*SIV2*SIV3*SIV4)^{(1/4)}$

Model equation:

The equation to calculate an overall HSI index for the roseatespoonbillIslandPresent model is:

CRN

According to our classification, this model's format is: **geometric mean**

Global sensitivity and uncertainty analysis:

We ran global sensitivity and uncertainty analyses on the roseatespoonbillIslandPresent model using the sensobol package in R (Puy et al. 2022). The following parameters were used for the sensobol analysis:

Table 3. Parameters and settings used for sensobol sensitivity and uncertainty analyses.

Parameter	Equation	Value
Number of input variables (M)	-	4
Base sample size (n)	-	107000
Number of model evaluations (N)	$n*(M+2)$	642000
First order estimator	See Puy et al. (2022)	Saltelli
Total order estimator	See Puy et al. (2022)	Jansen
Number of bootstrap replications	-	1000
Sampling scheme	-	Quasi-random
Matrices	-	A, B, AB

We ran a sensitivity and uncertainty analysis for the roseatespoonbillIslandPresent model using the original equation outlined in the documentation from Lewis (1983) and using arithmetic mean, geometric mean, limiting factor, and multiplicative equations to contrast the results across different equation structures.

Model uncertainty

We ran the roseatespoonbillIslandPresent model using 642000 combinations of its SIV variables, which were sampled from a uniform distribution spanning the range of possible values listed in the roseatespoonbillIslandPresent documentation. We limited the range of possible values for each parameter to the range in which the SIV values were greater than zero to prevent HSI score distributions with primarily zero values.




Figure 2. Empirical distributions of HSI scores for the roseatespoonbillIslandPresent model using the original geometric mean model equation from Lewis (1983), and an arithmetic mean, geometric mean, limiting factor, and multiplicative structure incorporating all SIV variables. Note differences in the y axis.

We assumed a uniform distribution for all parameters because we evaluated all ecorest models in batch. Should you decide to run your own sensitivity analysis, this assumption should be evaluated independently for each parameter in the model.

Table 4. Quantiles from the empirical distribution of HSI scores for the original roseatespoonbillIslandPresent model structure, an arithmetic mean equation, a geometric mean equation, a limiting factor equation, and a multiplicative equation structure.

	1%	2.5%	5%	25%	50%	75%	95%	97.5%	99%	100%
Original	0.20	0.26	0.32	0.53	0.67	0.79	0.94	0.97	1	1
Arithmetic	0.37	0.43	0.48	0.62	0.72	0.82	0.94	0.97	1	1
Geometric	0.20	0.26	0.32	0.53	0.67	0.79	0.94	0.97	1	1
Limiting	0.01	0.02	0.04	0.23	0.40	0.50	0.80	0.88	1	1
Multiplicative	0.00	0.00	0.01	0.08	0.20	0.38	0.78	0.87	1	1

Generally, models with outputs that span the entire range of possible outcomes (*i.e.*, 0-1) may be better at discriminating between different restoration actions or across habitat quality. The empirical distribution of HSI scores can also provide you with information about whether HSI scores for a given model tend to be skewed (*e.g.*, many low scores and few high scores, uniform probability of obtaining scores from 0-1, *etc.*)

Model sensitivity

Below are the results of the global sensitivity analysis for the roseatespoonbillIslandPresent model using the original equation, an arithmetic mean, a geometric mean, a limiting factor, and a multiplicative model structure. The sensobol package uses variance-based sensitivity metrics, so the model’s sensitivity to a given parameter is a measure of how much variance in the HSI score decreases in response to that parameter being fixed (Puy et al. 2022). For each parameter, the observed changes in the variance of the HSI score can be described with a first order sensitivity index (S_i) that accounts for the influence of a single parameter of interest on variance in HSI, or with a total order index (T_i) that accounts for the influence of a single parameter on its own and in combination with all other parameters (*i.e.*, interactions) (Puy et al. 2022). We can compare the 95% confidence intervals for the first and total order indices to a dummy parameter, which represents a parameter that has no influence on the variance in a model’s output. While an uninfluential variable should theoretically have an S_i and T_i of zero, small approximation errors can lead variables to have a non-zero influence on a model’s output (Puy et al. 2022). If the confidence interval of the S_i and T_i index for a given parameter overlaps the confidence interval of the dummy parameter, we can deduce that the parameter has a negligible effect on variance in HSI scores, both on its own and in combination with all other variables.

To ensure that the results of our sensitivity analysis converged, we determined whether the range of the 95% confidence interval for each sensitivity index was less than or equal to 0.05 (Sarrazin et al., 2016). Models that did not converge should be interpreted with caution.

Table 5. Model convergence records for the original roseatespoonbillIslandPresent model structure, an arithmetic mean equation, a geometric mean equation, a limiting factor equation, and a multiplicative equation structure.

	Did the model converge? (T/F)
Original	TRUE
Arithmetic	TRUE
Geometric	TRUE
Limiting	TRUE
Multiplicative	TRUE

Original model




Figure 3. Results of a sensitivity analysis for the roseatespoonbillIslandPresent model based on the original geometric mean model outlined in Lewis (1983). Dashed lines represent baseline numerical approximation error for S_i and T_i (*i.e.*, dummy variables).

Arithmetic mean model




Figure 4. Results of a sensitivity analysis for the roseatespoonbillIslandPresent model based on an arithmetic mean structure. Dashed lines represent baseline numerical approximation error for S_i and T_i (*i.e.*, dummy variables).

Geometric mean model




Figure 5. Results of a sensitivity analysis for the roseatespoonbillIslandPresent model based on a geometric mean structure. Dashed lines represent baseline numerical approximation error for S_i and T_i (*i.e.*, dummy variables).

Limiting factor model




Figure 6. Results of a sensitivity analysis for the roseatespoonbillIslandPresent model based on a limiting factor structure. Dashed lines represent baseline numerical approximation error for S_i and T_i (*i.e.*, dummy variables).

Multiplicative model




Figure 7. Results of a sensitivity analysis for the roseatespoonbillIslandPresent model based on a multiplicative mean structure. Dashed lines represent baseline numerical approximation error for S_i and T_i (*i.e.*, dummy variables).

Summary of influential variables

Original model

In the **original roseatespoonbillIslandPresent** model, **4 of 4** variables are influential.

Variable with the highest first order sensitivity: h.woody.veg.island.SIV

Variable with the highest total order sensitivity: h.woody.veg.island.SIV

Un-influential variables in original model:

None

Arithmetic mean model

In the **arithmetic mean roseatespoonbillIslandPresent** model, **4 of 4** variables are influential.

Variable with the highest first order sensitivity: h.woody.veg.island.SIV

Variable with the highest total order sensitivity: h.woody.veg.island.SIV

Un-influential variables in arithmetic mean model:

None

Geometric mean model

In the **geometric mean roseatespoonbillIslandPresent** model, **4 of 4** variables are influential.

Variable with the highest first order sensitivity: h.woody.veg.island.SIV

Variable with the highest total order sensitivity: h.woody.veg.island.SIV

Un-influential variables in geometric mean model:

None

Limiting factor model

In the **limiting factor roseatespoonbillIslandPresent** model, **4 of 4** variables are influential.

Variable with the highest first order sensitivity: h.woody.veg.island.SIV

Variable with the highest total order sensitivity: h.woody.veg.island.SIV

Un-influential variables in limiting factor mean model:

None

Multiplicative model

In the **multiplicative roseatespoonbillIslandPresent** model, **4 of 4** variables are influential.

Variable with the highest first order sensitivity: h.woody.veg.island.SIV

Variable with the highest total order sensitivity: h.woody.veg.island.SIV

Un-influential variables in multiplicative model:

None

References

1. Lewis, JC. 1983. Habitat suitability index models: roseate spoonbill. U.S. Dept. Int. Fish. Wildl. Serv. FWS/OBS-82/10.50. 16 pp.
2. McKay S, D Hernandez-Abrams, and K Cushway. 2025. ecorest: conducts analyses informing ecosystem restoration decisions. R package version 2.0.1, <https://CRAN.R-project.org/package=ecorest>.
3. Puy, A, S Lo Piano, A Saltelli, and SA Levin. 2022. sensobol: an R package to compute variance based sensitivity indices. Journal of Statistical Software 102(5):1-37. doi: 10.18637/jss.v102.i05
4. Sarrazin, F., Pianosi, F., & Wagener, T. (2016). Global sensitivity analysis of environmental models: Convergence and validation. Environmental Modelling & Software, 79, 135–152. <https://doi.org/10.1016/j.envsoft.2016.02.005>