



A tool to estimate and reduce GHGs from farms

The purpose of this document is to provide an introduction on how to use the Holos model (version 4) and the required vs. optional inputs.

For the purpose of this training, we are going to create a farm that has an annual beef production system, and a feed crop production system. The farm is located in Manitoba near Portage La Prairie.

Launch Holos

Please note that Holos 4 can be installed on a Microsoft Windows PC only. Mac OS will be supported in the next version.

Launch Holos by double-clicking on the Holos desktop icon. If there are no saved farms in the system, Holos will create a new farm and ask the user for a farm name and an optional comment (Figure 1). If there is already a saved farm in the system, Holos will ask the user to open the existing farm or to create a new farm (Figure 2).

Enter “**Holos 2022**” as a farm name and “**training version**” as the “**comment**”. Ensure to click the “**Advanced Mode**” button so that additional features and functionality will be available. Click “**Ok**” to proceed to the next screen

Ensure “**Metric**” is selected as the unit of measurement type and then click the “**Next**” button at the bottom of the screen (Figure 3).

Enter a name for this new farm

Name

Comments

Figure 1: Entering a name for the new farm.

Would you like to open an existing farm or create a new one?


Figure 2: If a farm has been previously saved, Holos will prompt to re-open that farm.

Select units of measurement

Figure 3: Select metric as the unit of measurement.

Creating and locating the new beef farm

The beef farm that we will create for this exercise is located in the province of Manitoba. Select “**Manitoba**” on the “**Select a province**” screen, and then click the “**Next**” button.



The screenshot shows a web interface for selecting a province. The title "Select a province" is displayed in a bold, dark font. Below the title is a dropdown menu with a blue border and a small downward arrow on the right, showing "Manitoba" as the selected option. At the bottom of the interface, there is a light gray bar containing a "Next" button.

Figure 4: Select manitoba as the province.

Holos uses **Soil Landscapes of Canada (SLC)**, which are a series of GIS coverages that show the major characteristics of soils and land for all of Canada (compiled at a scale of 1:1 million). SLC polygons may contain one or more distinct soil landscape components.

The “Farm Location” screen brings up a map of Canada with the province of Manitoba centered on the screen (Figure 5).

The map contains red colored polygons that can be selected by moving the cursor over the region that contains the location of your farm. You can zoom in or out of the map by using the mouse wheel or by hovering the cursor over the zoom icon at the bottom of the screen.

The beef farm for this example is located between Winnipeg and Portage la Prairie (Portage) with SLC polygon number **851003**.

1. Find and right-click on this polygon to select it on the map. Note that at this point daily climate data will be downloaded from [NASA](#). You can also select a specific polygon by clicking the “Enter a Polygon” button (Figure 6).

Note: Climate data is central to most calculations performed by Holos. For the most accurate estimation of farm emissions, measured climate data should be provided by the user which will override the default data obtained from the NASA weather API.

Holos will use daily precipitation, temperature, and potential evapotranspiration values to model soil carbon

change (climate parameter), nitrous oxide emissions, as well as ammonia volatilization.

Farm Location

Select the location of your farm by right-clicking on a region (use mouse wheel to zoom in or out)




Figure 5: Map of the Manitoba province showing the different selectable polygons.

Farm Location

Select the location of your farm by right-clicking on a region (use mouse wheel to zoom in or out)




Figure 6: Specifying a polygon ID using the option button.


Once the farm location is selected, soil information (texture, sand, and clay proportions) for the types of soils found in this region are displayed on the right side of the screen. It's possible that more than one soil type per region will be found and the user is expected to select their soil type from this list or use the default selection. (Figure 7)

For this tutorial, keep the default first selected soil type, and keep the default "Hardiness zone".

Farm Location

Select the location of your farm by right-clicking on a region (use mouse wheel to zoom in or out)

Enter a Polygon



Location Details

SLC polygon : Portage
Ecozone : Subhumid Prairies
Ecodistrict : 851
Province : Manitoba
Polygon : 851003
Latitude : 50.269704602224

Soil Zones Hardiness Zones Climate Data

Soil types in selected region

Multiple soil types found at this location, please select one from the following list

Black

Soil great group: Regosol
Soil texture: Fine
Proportion of clay in soil % : 30
Proportion of sand in soil % : 20
Drainage class: Well drained

Black

Soil great group: Black chernozem
Soil texture: Fine
Proportion of clay in soil % : 47
Proportion of sand in soil % : 28
Drainage class: Well drained

53° 45' 46.88" N 104° 49' 14.00" W

Back Next

Canada

851003 (Portage, Manitoba) | Total emissions: 0 (kg CO₂e) | Profit : \$ 0.00 | Single-Year Mode

Figure 7: Multiple soil types might be available for a given region.

Note: Soil data obtained from the user's selected location will be used in the calculation of location-specific N₂O emission factors. Properties such as soil texture, top layer thickness, and soil pH are required for these calculations, and can be overwritten.

Click the "Next" button to proceed to the next step.

Selecting Farm Components

Now that the farm location has been selected, we can move on to the “Component Selection” screen. This is where the user can select different components for their farm. Holos will display all available components on the left side of the screen under the “All Available Components” column (Figure 8). These components are grouped into various categories including Land Management, Beef Production and Dairy Cattle.

If we click on the drop down button next to a categories' name, we can then see the available components in that category. For this portion of the training section, we will be are working with the “Land management” and “Beef production” categories.




Figure 8: The available components screen. Specific components can be chosen here to include in the farm.

The Holos model is designed to define the land management before livestock. This is because we are allowing livestock to be placed onto a specific Pasture (field) for grazing, and that is easier done when a pasture field has been defined already (otherwise the user would have to interrupt the livestock setup to setup a field).

Crop and Hay Production

Note: In order to calculate soil carbon change for fields and crop rotations, click on the "Settings" menu and select the "Multi-year" option (default is Single-Year).

Now we can add our first component to the farm. Drag a "Field" component from the left side of the screen and drop it on the "My Farm" on the right side (Figure 6). The screen will now update to reflect this new component that you have added to your farm. Holos will label the field as "Field #1". At this point, we can now enter production information related to the crop being grown on this field.




Figure 9: Adding a component to the farm.

Wheat with Cover Crop

Our first field on the farm will grow continuous wheat with a cover crop of hairy vetch. Change the following elements in the "Field #1" component.

1. Rename the field to "**Wheat & Hairy Vetch**" in the "**Step 1**" section of the screen. Change the area of the field to **18 ha**.
2. Select "**Wheat**" as the main crop and "**Hairy Vetch**" as the cover crop in "**Step 2**".
3. Under the "**General**" tab:
 - Enter a yield of **3000 kg/ha** (wet weight). The dry weight value will automatically be calculated based on the moisture content of crop value.
 - Select "**Reduced Tillage**" as the tillage type.

- Enter "200" as the amount of irrigation.
- Select '0' as the pesticide passes.
- Leave Harvest method as default selection.

File
Settings
Tools
View
Help

Component Selection

Select components for your farm by dragging and dropping items from the "Available Components" column to the "My Farm" column. Click on an item under the "My Farm" column to enter details.

Available Components

Land management

Crop Rotation
Shelterbelt
Field

Beef production
Dairy cattle
Swine
Sheep
Other livestock
Poultry
Infrastructure

My Farm

Land management

Wheat & Hairy vetch

Wheat & Hairy vetch

Enable Multi-Year Mode : Yes No

Step 1: Define the general properties for this field

Field name : Wheat & Hairy vetch
Total area of this field : 18 (ha)
Start year of field history : 1985
End year : 2022

Step 2: Click the 'Add Crop' button to describe the crops grown on this field. Enter one crop for continuous fields (i.e. continuous wheat) and the first phase for crop rotations (i.e. wheat, fallow). Holos will use this entry to populate the field history.

Add Crop

Year	Crop	Winter/Cover/Undersown Crop
2022	Wheat	Hairy Vetch (Vicia villosa roth)

Step 3: Adjust details of the crop (wheat) grown on this field

Show Additional Information : Yes No

General
Fertilizer
Manure
Winter & Cover Crops
Residue
Economics

General Properties

Amount of irrigation, pesticide usage, and yield for the selected crop can be entered on this tab. Harvested yield can be entered as either wet weight or dry weight. Additional information can be entered by clicking on the 'Show Additional Information' button.

Yield (wet weight) : 3,000 (kg ha⁻¹)
Yield (dry weight) : 2,640 (kg DM ha⁻¹)
Moisture content of crop : 12 (%)
Tillage : Reduced tillage
Harvest method : Cash crop
Amount of irrigation : 200 (mm)
Number of pesticide passes : 0

Back
Next

Canada

851003 (Portage, Manitoba)
Total emissions: 15.90 (Mg CO2e)
Loss : \$ 17,076.24
Multi-Year Mode

Figure 10: Field Component of the farm.

4. Select the 'Fertilizer' tab and click the "Add Fertilizer Application" button. Holos has now added a new fertilizer application for this field and will suggest Urea as the fertilizer blend. A default application rate is calculated based on the yield value entered for this field. Details of this fertilizer application can be changed by clicking the "Show Additional Information" button (e.g., season of application, different fertilizer blend, etc.).

File
Settings
Tools
View
Help

Component Selection

Select components for your farm by dragging and dropping items from the "Available Components" column to the "My Farm" column. Click on an item under the "My Farm" column to enter details.

Available Components

Land management

Crop Rotation

Shelterbelt

Field

Beef production

Dairy cattle

Swine

Sheep

Other livestock

Poultry

Infrastructure

My Farm

Land management

Wheat & Hairy Vetch

Wheat & Hairy Vetch

Step 1: Define the general properties for this field

Field name :

Wheat & Hairy Vetch

Total area of this field :

1 (ha)

Start year of field history :

1985

End year :

2022

Step 2: Click the 'Add Crop' button to describe the crops grown on this field. Enter one crop for continuous fields (i.e. continuous wheat) and the first phase for crop rotations (i.e. wheat, fallow). Holos will use this entry to populate the field history.

Add Crop

Year	Crop	Winter/Cover/Undersown Crop
2022	Wheat	Hairy Vetch (<i>Vicia villosa</i> roth)

Step 3: Adjust details of the crop (wheat) grown on this field

Show Additional Information : Yes No

General Fertilizer Manure Winter & Cover Crops Residue Economics

2022 Wheat - Fertilizer Management

Multiple fertilizer applications can be made to this field throughout the year. To add a fertilizer application, click 'Add Fertilizer Application' to specify the details.

Add Fertilizer Application

Season of application	Blend	Application rate (kg ha ⁻¹)
Spring	Urea	52.5

A suggested fertilizer application rate of 52.5 (kg ha⁻¹) for wheat with a yield of 3000 (kg ha⁻¹) has been set. This value can be adjusted as needed.

Back

Next

Canada

731014 (Daysland Plain, Alberta)
Total emissions: 505 (kg CO₂e)
Loss : \$ 711.60
Multi-Year Mode

Figure 11: Adding fertilizer to a field.

Note: At a minimum, Holos requires the area of the field, type of crop grown, and a field-specific fertilizer application rate to calculate direct and indirect nitrous oxide emissions.

Residue management of each crop (and cover crop) can be adjusted in Holos (see 'Residue' tab). Holos provides default values depending on the type of crop being grown and will set a value for percentage of product returned to soil, percentage of straw returned to soil, etc. These residue input settings will have an impact on the final soil carbon change estimates.

Furthermore, biomass fractions and N concentrations can be overwritten by the user, and in this way 'custom' crops can be added that are currently not available.

Page 10

Pasture (native) / Grasslands Information

The cow-calf operation (defined later on) relies on native pasture for the summer months (May through October).

1. Drag a new "Field" tab component to your list of components. Enter the name "Native Grassland" in the 'Field name' input box.
2. Enter "100" as the total area of the field.
3. Select "Seeded Grassland" from the crop list under 'Crop' column (step 2). Please note that Holos auto populates the 'Winter/Cover/Undersown Crop' area when a perennial crop is selected.
4. Keep '0' as the amount of irrigation and pesticide passes.
5. No fertilizer is used for this crop.

The screenshot displays the 'Holos 2022' software interface. On the left, the 'Component Selection' panel shows 'Available Components' (Land management, Beef production, Dairy cattle, Swine, Sheep, Other livestock, Poultry, Infrastructure) and 'My Farm' (Native Grassland, Wheat & Hairy Vetch). The main panel is titled 'Native Grassland' and contains three steps: Step 1 (Define general properties), Step 2 (Add Crop), and Step 3 (Adjust details of the crop). Step 1 shows 'Field name: Native Grassland', 'Total area of this field: 100 (ha)', 'Start year of field history: 1985', and 'End year: 2022'. Step 2 shows 'Add Crop' button and a table with columns 'Year', 'Crop', and 'Winter/Cover/Undersown Crop'. Step 3 shows 'General Properties' with 'Amount of irrigation: 0 (mm)' and 'Number of pesticide passes: 0'. A footer bar includes 'Canada' logo, location '851003 (Portage, Manitoba)', 'Total sequestration: 91.56 (Mg CO2e)', 'Loss: \$ 17,076.24', and 'Multi-Year Mode'.

Figure 12: Native Grasslands information.

Barley Grain and Mixed Hay Rotation

To demonstrate the crop rotation component (as opposed to using individual field components), we will assume that barley grain and mixed hay are grown in rotation, with the mixed hay under seeded to the barley so that it can be harvested in both main years (example derived from University of Alberta's Breton plots, Figure).

When using the "Crop Rotation" component, any sequence of crops that are input into this components will be applied to each individual field that is part of the rotation setup. This means one field is added for each rotation phase, and the rotation shifts so that each rotation phase is present on one field. Since each field can have a

different historical management, soil carbon algorithms will run for each field.

For this example, we assume that the farm requires **70 ha** of barley grain and mixed hay, which are grown in rotation. We will need to setup three fields where barley grain is rotated in each field every two years (Figure 7). When using the crop rotation component, the crop management input of a specific crop is repeated on each field in the rotation where the crop is grown.

To setup the rotation:

1. Add one "**Crop Rotation**" component from the available components.
2. To expand the horizontal space available in Holos, click on "**View**" from the top menu bar and select "**Hide List of Available Components**".
3. The rotation of this field **begins in 1985 and ends in 2022**. Under step 1, please ensure that these two values are set as the start and end year respectively.
4. Enter "**70**" ha as the **total area** of this field.
5. Under **Step 2** change the crop to **Barley**. The **year** for this crop should be 2022.
 - Under the **General Tab** enter 3000 kg/ha (wet weight) as the **yield** for this crop.
 - Change the **tillage type** to **Reduced Tillage**.
 - Keep **0** as the amount of irrigation and number of pesticide passes.
6. Now add another crop to this rotation. Click on "**Add Crop**" under "**Step 2**" to add a second crop to the rotation. Note that Holos sets the year for this new crop to 2021 or one before the previous crop's year. This means that Holos is expecting the user to enter crops that have been grown in reverse order back to 1985.

***Note:** It is not necessary to enter a crop for each individual year going back to 1985, only enough crops to describe a single phase of the rotation will need to be entered by the user. Holos will then copy the phase information and back-populate the field history (e.g., Holos will copy the rotation back to 1985 on behalf of the user).*

7. For this newly added crop select "**Tame Mixed(grass/legume)**" as the crop type.
8. Click on the "**Add crop**" button one more time. For this third crop, select "**Tame Mixed(grass/legume)**" once again as the crop type.

Holos 2022

File Settings Tools View Help

Component Selection

Select components for your farm by dragging and dropping items from the "Available Components" column to the "My Farm" column. Click on an item under the "My Farm" column to enter details.

My Farm

Land management

- Crop rotation #1
- Native Grassland
- Wheat & Hairy Vetch

Crop rotation #1

Enable Multi-Year Mode: ☐ Yes ☒ No

Step 1: Starting with one field in the rotation, enter the start and end year of this rotation and the size of the field

Start year: 1985
End year: 2022
Total area of this field: 70 (ha)

Step 2: Click the 'Add Crop' button to add crops to this rotation. Keep adding crops until the first phase of this rotation has been listed. Holos will add a new field to your farm for each crop in the rotation.

Add Crop

Year	Crop	Winter/Cover/Undersown Crop		
2020	Tame mixed (grass/legume)	Tame mixed (grass/legume)	X	↓
2021	Tame mixed (grass/legume)	Tame mixed (grass/legume)	X	↑
2022	Barley	None	X	↑

Step 3: Adjust the details of barley. These adjustments will be copied to the other fields in the rotation when barley was grown

Show Additional Information: ☐ Yes ☒ No

General Fertilizer Manure Winter & Cover Crops Residue Economics

General Properties

Amount of irrigation, pesticide usage, and yield for the selected crop can be entered on this tab. Harvested yield can be entered as either wet weight or dry weight. Additional information can be entered by clicking on the "Show Additional Information" button.

Yield (wet weight): 3,000 (kg ha⁻¹)
Yield (dry weight): 2,640 (kg DM ha⁻¹)
Moisture content of crop: 12 (%)
Tillage: Reduced tillage
Harvest method: Cash crop
Amount of irrigation: 0 (mm)
Number of pesticide passes: 0

Back Next

Canada

851003 (Portage, Manitoba) Total sequestration: 198.21 (Mg CO₂e) Loss: \$ 164,504.54 Multi-Year Mode

Figure 13: An example of a crop rotation of three crops.

- Now add harvest data to each of the tame mixed crops. You will need to select each **tame mixed** crop and add the harvest data to that specific crop. So select the first tame mixed crop (2021) and then:
 - Go under the **Harvest Tab** and click **Add Harvest Date** button to create a new harvest.
 - Select a Harvest date of "August 31, 2021", assuming the harvest is done on the same day every year.
 - Select **Mid** for **Forage growth stage**.
 - Enter **5** as the total number of bales.
 - Enter **500** as the **Wet bale weight**.
- Repeat the above steps (step 9) for the second tame mixed crop.

If the tame mixed field is harvested more than once, the "Add Harvest Date" button can be used to add subsequent harvests.

Cow-Calf Operation

Click the view menu item again and uncheck **Hide List of Available Components** option so that we can see all of the available components again.

Adding animal components follows the exact same approach that was used for land management components. Under the **Beef Production** category in the available components, drag and drop one **Beef Cow-Calf** component to the **My farm** section on the right. **Replacement heifers** will not be used in this example, so we will remove this group by clicking the X icon right next to its entry under **Step 1**.

FileSettingsToolsViewHelp

Component Selection

Select components for your farm by dragging and dropping items from the "Available Components" column to the "My Farm" column. Click on an item under the "My Farm" column to enter details.

Available Components

Land management

Beef production

Beef Cow-Calf

Beef Stockers & Backgrounders

Beef Finisher

Dairy cattle

Swine

Sheep

Other livestock

Poultry

Infrastructure

My Farm

Beef production

Beef Cow-Calf

Land management

Crop rotation #1

Native Grassland

Wheat & Hairy vetch

Beef Cow-Calf

Enable Multi-Year Mode : Yes No

Step 1: Define the animal groups for this cow-calf component

Add Group

Group name		
> Bulls		X
Cows		X
Calves		X

Step 2: Define the management periods for the selected animal group (bulls)

Add Management Period

Management period name	Start date	End date	Number of days
> Winter feeding	01-01-2021	04-30-2021	120
Summer grazing	05-01-2021	10-31-2021	184
Extended fall grazing	11-01-2021	12-31-2021	61

Step 3: Define the management of bulls during this selected period of time (2021-01-01 - 2021-04-30)How Additional Information :

GeneralDietHousingManure

General Management

Basic information about this group of animals can be entered here. Values entered here will apply to the animals during the selected management period.

Number of animals :

4

Daily gain :

0

(kg day⁻¹)

Start weight :

820.00

(kg)

End weight :

820.00

(kg)

BackNext

Canada

851003 (Portage, Manitoba) | Total emissions: 443.22 (Mg CO2e) | Loss : \$ 164,504.54 | Multi-Year Mode

Figure 14: The Beef Cow-Calf Component.

Entering Beef Cows, Calves and Bulls Information.

Beef Cows - Winter Feeding

Following the annual feeding cycle, the beef farm we are working with is **divided into three management (production) periods**. We can now enter production and management data corresponding to these three management periods.

1. Under the animal groups section in '**Step 1**', make sure that the "**Cows**" row is selected in order to enter

Page 14

the associated management information for that group.

2. Click the management period named **"Winter Feeding"** in 'Step 2' to activate that management period.
3. Ensure **"January 1, 2021"** is set as the 'Start date' and that **"April 30, 2021"** is set as the 'End date' (120 days). Note that the 'Number of days' being shown will be inclusive of the start and end dates.

Next, we can enter data related to the number of animals, diet, manure system, and housing type.

- **General Tab:**

- Enter **150** as the number of animals.
- Keep the remaining entries at their default values.

***Note:** The number of animals, average daily gain, and feed quality are the minimum required inputs for calculating methane and nitrous oxide emissions. Length of management periods (i.e., duration of grazing) also will be needed. Housing and manure management information are also important inputs but are relatively more impactful on the emissions of monogastrics.*

- **Diet Tab:**

We are going to create a custom diet for our group of cows during the **"Winter feeding"** management period. (Holos incorporates feed ingredient information from the recently published Nutrient Requirements of Beef Cattle book (2016).

Click on the 'Diet' tab. Since we are going to create our own custom diet, we will click on the 'Custom Diet Creator' button. Note that Holos provides a default set of animal diets that can be used.

- Custom-Diet Creator:

- Click the **"Add Custom Diet"** button in the **"Step 1"** section of the screen to create a new custom diet.
- Rename this diet to **"My Custom Cow Diet"** then press the Enter key to save the name.
- Under step 1, ensure that you have the **Animal Type** set to **Beef Cow**.
- To add ingredients to our new diet, select **"Alfalfa hay"** from the ingredient list, and then click the **"Add Selected Ingredient to Diet"** button.
- We will add one more ingredient to our diet. Select **'Barley Hay'** from the ingredient list, and then click the **"Add Selected Ingredient to Diet"** button.
- Enter **50%** for **'Barley Hay'** and **50%** for **'Alfalfa Hay'** in **"Step 3"**. Note that Holos now reports the diet being complete since all ingredients total up to 100%.

- Click the **"OK"** button to save the new custom diet
- Select the **"My Custom Cow Diet"** from the drop down menu in front of **Diet Type**.

***Note:** Diet quality information such as crude protein, total digestible nutrient, and fat are required inputs so that Holos can estimate enteric methane emissions from an animal group.*

Custom Diet Creator

Step 1: To begin creating a diet press "Add Custom Diet"

Diet name	Animal type	Forage (1x, %)	CP (% DM)	TDN (1x, %)	Fat (% DM)	NDF (% DM)	M
My Custom Cow Diet	Beef cow	100	15.35	57.7	1.95	49.3	

Step 3: Diet is complete

Ingredient	Percentage in diet (% DM)		
Alfalfa hay	50	✕	
Barley hay	50	✕	
Total %: 100			

Step 2: Add ingredients to your diet by selecting the ingredient and then click "Add Selected Ingredient to Diet"

Ingredient	Dry matter (% AF)	Forage (%)	CP (% DM)	TDN (% DM)	Starch (% DM)	Fat (% DM)	ME (Mcal kg ⁻¹)
Barley grain	89.7	0	12.8	84.1	56.7	2.2	3
Barley grain flaked	81.1	0	12.5	84	59.3	2	3
Barley hay	88	100	10.9	60.2	5.7	2.4	2.2
Barley silage	33.6	100	12	60.6	9.2	3.5	2.2
Barley straw	85.1	100	6.1	48.3	14.9	1	1.7
Beet pulp, dry	91.5	0	9.1	66.6	0.9	1.1	2.4
Beet pulp, wet	21.9	0	9.5	66.6	1.7	0.9	2.4
Bermudagrass fresh	34.9	100	15.2	57.3	1.8	2.8	2.1

OK

Figure 15: Custom diet creator for Cows animal group.

- Housing Tab:
 - Select **Confined no barn** as the housing type.
- Manure Tab:
 - Select **Deep Bedding** as the manure handling system.

Beef Cows - Summer Grazing

Click on the management period named **Summer Grazing**. Ensure that the start date is set to May 1st, 2021 and the end date is set to October 31st, 2021.

- General Tab:
 - Enter **150** as the number of animals.
- Diet Tab:
 - Select **High energy protein** as the diet type.
- Housing Tab:
 - Select **Pasture** as the housing type.
 - Select **Native Grassland** as the pasture location.
- Manure Tab:

- Select **Pasture** as the manure handling system.

Beef Cows - Extended Fall Grazing

Click on the management period named **Extended Fall Grazing**. Ensure that the start date is set to November 1st, 2021 and the end date is set to December 31st, 2021.

- **General Tab:**
 - Enter **150** as the number of animals.
- **Diet Tab:**
 - Select **Medium energy protein** as the diet type.
- **Housing Tab:**
 - Select **Pasture** as the housing type.
 - Select **Native Grassland** as the pasture location.
- **Manure Tab:**
 - Select **Pasture** as the manure handling system.

File
Settings
Tools
View
Help

Component Selection

Select components for your farm by dragging and dropping items from the "Available Components" column to the "My Farm" column. Click on an item under the "My Farm" column to enter details.

Available Components

Land management

- Crop Rotation
- Shelterbelt
- Field

Beef production

- Beef Cow-Calf
- Beef Stockers & Backgrounders
- Beef Finisher

Dairy cattle

- Swine
- Sheep
- Other livestock
- Poultry

My Farm

Beef production

- Beef Cow-Calf

Land management

- Crop rotation #1
- Native Grassland
- Wheat & Hairy Vetch

Beef Cow-Calf

Step 1: Define the animal groups for this cow-calf component

Add Group

Group name	
Bulls	X
Cows	X
Calves	X

Step 2: Define the management periods for the selected animal group (cows)

Add Management Period

Management period name	Start date	End date	Number of days	
Winter feeding	01-01-2021	04-30-2021	120	X
Summer grazing	05-01-2021	10-31-2021	184	X
Extended fall grazing	11-01-2021	12-31-2021	61	X

Step 3: Define the management of cows during this selected period of time (2021-01-01 - 2021-04-30) Show Additional Information : Yes No

General
Diet
Housing
Manure

General Management

Basic information about this group of animals can be entered here. Values entered here will apply to the animals during the selected management period.

Number of animals : 150
Daily gain : 0 (kg day⁻¹)
Start weight : 610.00 (kg)
End weight : 610.00 (kg)
Milk production : 8.00 (kg day⁻¹)

Back
Next

Canada

851003 (Portage, Manitoba)
Total emissions: 547.95 (Mg CO₂e)
Loss : \$ 164,504.54
Multi-Year Mode

Figure 16: Beef Cow-Calf, Cow group

Bulls

Click on the **Bulls** row in the animal group section **Step 1**. Information related to diet, housing and manure management is identical to the cows group.

- Right click on the **Bulls** animal group. A menu will appear allowing you to select the option to copy management periods from another animal group. Since the management for the bulls is similar to the management for the cows, click the **Copy Management From → Cows** sub-menu item.
- Adjust the **number of bulls** for each of the three management periods to 4.

Component Selection
Select components for your farm by dragging and dropping items from the "Available Components" column to the "My Farm" column. Click on an item under the "My Farm" column to enter details.

Available Components

- Land management
 - Crop Rotation
 - Shelterbelt
 - Field
- Beef production
 - Beef Cow-Calf
 - Crop rotation #1
 - Native Grassland
 - Wheat and Hairy Vetch
- Dairy cattle
- Swine
- Sheep
- Other livestock
- Poultry
- Infrastructure

My Farm

- Beef production
 - Beef Cow-Calf
- Land management
 - Crop rotation #1
 - Native Grassland
 - Wheat and Hairy Vetch

Beef Cow-Calf

Step 1: Define the animal groups for this cow-calf component

Add Group

Group name	Management period name	Start date	End date	Number of days	
Bulls	Winter feeding	01-01-2021	04-30-2021	120	X
Cows	Cows grazing	05-01-2021	10-31-2021	184	X
Calves	Calves grazing	11-01-2021	12-31-2021	61	X

Step 2: Define the management periods for the selected animal group (bulls)

Add Management Period

Step 3: Define the management of bulls during this selected period of time (2021-01-01 - 2021-04-30)

Show Additional Information: Yes No

General Management

Basic information about this group of animals can be entered here. Values entered here will apply to the animals during the selected management period.

Number of animals: 4

Daily gain: 0 (kg day⁻¹)

Start weight: 820.00 (kg)

End weight: 820.00 (kg)

Back Next

Canada

851003 (Portage, Manitoba) Total emissions: 406.98 (Mg CO2e) Loss: \$ 148,376.98 Multi-Year Mode

Figure 17: Copying data from another animal group

Beef Calves

Calves on our farm are born on March 1 and weaned on September 30 at the age of seven months. Using a final weaning rate of 85%, we will have 110 calves from March to September. Following the cows, calves will be in confinement for the months of March and April and will be grazing on pasture from May to September. This will result in two separate management periods.

Click on the **Calves** row in the animal group section **Step 1** to activate the calf group. The first management period will span from **March 1, 2021 to April 30, 2021** and the second management period will span from **May 1, 2021 to September 30, 2021**.

Management Period # 1:

Rename this period from Management Period # 1 to **Confinement**.

- **General Tab:**
 - Enter **110** as the number of animals.
 - All other options as default.
- **Diet Tab:**
 - Select **Medium energy protein** as the diet type.
- **Housing Tab:**
 - Select **Confined no barn** as the housing type.
- **Manure Tab:**
 - Select **Deep bedding** as the manure handling system.

Management Period # 2:

Rename this period from Management Period # 2 to **Grazing**.

- **General Tab:**
 - Enter **110** as the number of animals.
- **Diet Tab:**
 - Select **High energy protein** as the diet type.
- **Housing Tab:**
 - Select **Pasture** as the housing type.
 - Select **Native Grassland** as the pasture location.
- **Manure Tab:**
 - Select **Pasture** as the manure handling system.

Beef Stocker & Backgrounder operation

To enter information on backgrounder and stocker animals, we will add a new "**Beef Stockers & Backgrounders**" component to our farm.

- Drag and drop a new **Beef Stockers & Backgrounders** component. The beef farm manages 200 backgrounders (100 steers and 100 heifers).
- Click on the **Heifers** group to activate it and to enter management data for this group.
 - For "Management period #1", enter "**October 1, 2021**" as the '**Start date**' and "**January 18, 2022**" as the '**End date**' (110 days).
 - **General Tab:**

- Enter 100 as the number of animals.
- Enter 1.1 kg/day as the daily gain.
- Enter 240 kg as the start weight.

■ **Diet Tab:**

- Select **Medium Growth** as the diet type.
- Select **None** as the diet additive.

■ **Housing Tab:**

- Select **Confined no barn** as the housing type.

■ **Manure Tab:**

- Select **Deep bedding** as the manure handling system.

- The management data for the **Steers** group is the same as **Heifers**. Right click on the Steers group to activate the right-click menu and select **Copy Management From → Heifers**

Holos 2022

File Settings Tools View Help

Component Selection

Select components for your farm by dragging and dropping items from the "Available Components" column to the "My Farm" column. Click on an item under the "My Farm" column to enter details.

Available Components

Land management

Beef production

Dairy cattle

Swine

Sheep

Other livestock

Poultry

Infrastructure

My Farm

Beef production

Beef Cow-Calf

Beef Finisher

Beef Stockers & Backgrounders

Land management

Crop rotation #1

Native Grassland

Wheat & Hairy vetch

Beef Stockers & Backgrounders

Step 1: Define the animal groups for this backgrounding component

Add Group

Group name	
Heifers	X
Steers	X

Step 2: Define the management periods for the selected animal group (heifers)

Add Management Period

Management period name	Start date	End date	Number of days	
Management period #1	10-01-2021	01-18-2022	110	X

Step 3: Define the management of heifers during this selected period of time (2021-10-01 - 2022-01-18) Show Additional Information: Yes No

General Diet Housing Manure

General Management

Basic information about this group of animals can be entered here. Values entered here will apply to the animals during the selected management period.

Number of animals:

100

Daily gain:

1.09

(kg day⁻¹)

Start weight:

240

(kg)

End weight:

360

(kg)

Please review the selected diet information, body weight, and daily gain to ensure animals are being provided with their daily energy requirements.

Back

Next

Canada

851003 (Portage, Manitoba) | Total emissions: 1,074.93 (Mg CO2e) | Loss: \$ 88,849.06 | Multi-Year Mode

Finishing Feedlot Operation

We will now **repeat the steps used for Beef Stockers & Backgrounders** to enter the beef feedlot management data. Drag a new **Beef Finisher** component from **All components** to your list of components.

- The beef farm manages 200 backgrounders (100 steers and 100 heifers) in a feedlot operation for **170 days**.
- Click on the **Heifers** group to activate it and to enter management data for this group.
 - For "Management period #1", enter "**January 19, 2022**" as the **Start date** and "**July 7th, 2022**" as the **End date**.
 - **General Tab:**
 - Enter **100** as the number of animals.
 - Enter **1.2 kg/day** as the daily gain.
 - Enter **350 kg** as the start weight.
 - **Diet Tab:**
 - Select **Barley grain based diet** as the diet type.
 - Select **None** as the diet additive.
 - **Housing Tab:**
 - Select **Confined no barn** as the housing type.
 - **Manure Tab:**
 - Select **Deep bedding** as the manure handling system.
- The management data for the **Steers** group is the same as **Heifers**. Right click on the Steers group to activate the right-click menu and select **Copy Management From → Heifers**

File
Settings
Tools
View
Help

Component Selection

Select components for your farm by dragging and dropping items from the "Available Components" column to the "My Farm" column. Click on an item under the "My Farm" column to enter details.

Available Components

Land management

Crop Rotation
Shelterbelt
Field

Beef production

Beef Cow-Calf
Beef Stockers & Backgrounders
Beef Finisher

Dairy cattle
Swine
Sheep
Other livestock
Poultry
Infrastructure

My Farm

Beef production

Beef Finisher
Beef Stockers & Backgrounders
Beef Cow-Calf

Land management

Crop rotation #1
Native Grassland
Wheat & Hairy Vetch

Beef Finisher

Step 1: Define the animal groups for this finishing component

Add Group

Group name	
Heifers	X
Steers	X

Step 2: Define the management periods for the selected animal group (heifers)

Add Management Period

Management period name	Start date	End date	Number of days	
Management period #1	01-19-2022	07-07-2022	170	X

Step 3: Define the management of heifers during this selected period of time (2022-01-19 - 2022-07-07)

Show Additional Information: Yes No

General
Diet
Housing
Manure

Manure Management

Manure handling system details can be entered on this tab. To adjust optional manure handling system information, click the 'Show Additional Information' button.

Manure handling system : Deep bedding

Back
Next

Canada

851003 (Portage, Manitoba)
Total emissions: 1,011.40 (Mg CO2e)
Loss: \$ 164,504.54
Multi-Year Mode

Figure 19: Beef Finisher component

Adding a Manure Application to the Wheat Field

Holos has the ability to add manure applications from manure that is sourced from livestock on the current farm or from imported manure (off-site). Since we have now defined our animal components, we can apply manure to any field on our farm.

1. Select the **Wheat & hairy vetch** field from the list of components added to our farm.
2. Click on the **Manure** tab and then click the **Add Manure Application** button.
 - Select **Beef cattle** as the **Manure type**.
 - Select **Livestock** as the **Origin of manure**.
 - Select **Solid storage** as the **Manure handling system**.
 - Enter **1,000 kg/ha** as the amount of manure applied to this field.
3. Note that both chemical fertilizers and manure applications can be made on the same field

Adding supplemental hay/forage for grazing animals

We can also add additional hay/forage for animals that are grazing on a particular field. Since we have now placed a group of animals on the "Native Grassland" field component, and we have also provided harvest information for our mixed hay crops on the crop rotation component, we can then add an additional forage supplement for these grazing animals.

1. Select the **Native Grassland** field component we created earlier.
2. Click on the **Grazing** tab.
 - Click the **Add Supplemental Hay** button to add additional forage for the animals on this field.
 - Enter "On-farm" as the **Sources of bales**.
 - Choose **Crop rotation #1 [Field #2] - Tame Mixed (grass/legume)** under **Field** to select the source of the supplemental hay.
 - Change the **Number of bales** to 1
 - Enter **500** as the wet bale weight.
 - Keep the moisture content as the default value.

Component Selection
Select components for your farm by dragging and dropping items from the "Available Components" column to the "My Farm" column. Click on an item under the "My Farm" column to enter details.

My Farm

- Beef production
 - Beef Finisher
 - Beef Stockers & Backgrounders
 - Beef Cow-Calf
- Land management
 - Crop rotation #1
 - Native Grassland
 - Wheat & Hairy Vetch

Total area of this field : 100 (ha)
Start year of field history : 1985
End year : 2022

entry to populate the field history.

Add Crop

Year	Crop	Winter/Cover/Undersown Crop
2022	Seeded grassland	Seeded grassland

Step 3: Adjust details of the crop (seeded grassland) grown on this field

Show Additional Information : Yes No

General Fertilizer Manure Harvest Grazing Winter & Cover Crops Residue Economics

2022 Seeded grassland - Grazing Details

Grazing Animal Groups

All animal groups grazing on this pasture are displayed below. Modifications to grazing period dates can be made by adjusting the management period of the animal group(s). All manure from animals grazing on this field is accounted for. Additional manure applications from other sources can be added to the field from the manure tab.

- 4 bulls grazing from 1 May to 31 October (184 days), with a high energy protein feed [Edit Group](#)
- 4 bulls grazing from 1 November to 31 December (61 days), with a medium energy protein feed [Edit Group](#)
- 150 cows grazing from 1 May to 31 October (184 days), with a high energy protein feed [Edit Group](#)
- 150 cows grazing from 1 November to 31 December (61 days), with a medium energy protein feed [Edit Group](#)
- 110 calves grazing from 1 May to 30 September (153 days), with a high energy protein feed [Edit Group](#)

Add Supplemental Hay/Forage for Grazing Animals

Animals grazing on this field can be fed additional hay by clicking the 'Add Supplemental Hay' button. Hay can be imported by setting the 'Source of bales' to 'Off-farm'.

Add Supplemental Hay

Date	Source of bales	Field	Number of bales	Wet bale weight (kg)	Moisture content (%)
03-11-2022	On-farm	Crop rotation #1 [Field #2] - Tame mixed (grass/legume)	1	500	12

4 bales harvested from field 'Crop rotation #1 [Field #2] - Tame mixed (grass/legume)' available for supplemental feed

Back Next

Canada

B51003 (Portage, Manitoba) Total emissions: 12,149.43 (Mg CO2e) Loss : \$ 164,504.54 Multi-Year Mode

Figure 20 - Adding supplemental hay/forage for grazing animals.

Pullet Farm Operation:

We will add one last animal component to our farm. In addition to the beef cattle operations of this farm, we will also be adding a "Chicken Meat Production" component to our farm. If you hover your mouse cursor over the "Chicken Meat Production" component under the "Poultry" category, Holos will display a tooltip that gives a brief description of a chicken meat production operation:

"Chicks arriving in the operation from a multiplier hatchery are raised to market weight (1-4 kg)"

1. Drag one "Chicken Meat Production" component to the farm. For each group (Pullets and Cockerels), each management period for that group will consist of **400** animals. This means numbers of animals will be consistent throughout the management periods / year.

2. Select the **Pullets** group. The start and end dates for each management period will be:
 - **Brooding Stage:** Start January 1st, 2022 - End January 22nd, 2022.
 - **Rearing Stage:** Start January 23rd, 2022 - End June 26th, 2022.
 - **Rearing Stage:** Start June 27th, 2022 - End November 28th, 2022.
3. For each management period, set number of animals to **400**.
4. Leave the entries in Housing and Manure as default.
- The management data for the **Cockerels** group is the same as **Pullets**. Right click on the Cockerels group to activate the right click menu and select **Copy Management From → Pullets**

Timeline Screen

We are now finishing the process of defining our farm. Click the **Next** button to go forward to the timeline screen. The timeline screen provides a visual layout of all the fields from 1985 to the specified end year for each field. This screen also allows the user to add historical and projected production systems.

The **Add Historical Production System** button enables the user to add a different cropping history to individual fields whereas the **Add Projected Production System** button enables the user to add a future (projected) cropping system to individual fields.

Adding a historical production system

We will assume that the barley grain and mixed hay rotation fields were previously in a continuous wheat cropping system between **1985 and 2000**.

1. To add a new historical cropping system, select one of the fields that are in the barley grain and mixed hay rotation. To select an item, click on the timeline bar to activate that field. We will select the first field in this rotation (i.e., the field with the name of "**Crop rotation #1 [Field #1]**")
2. Click on the **Add Historical Production System** button which will add a new row to the table under the "**Step 1**" section in the upper left section of the screen. Notice that this new entry has the words "**Historical management practice**" added.
3. We will set the end year of this historical management practice to the year **2000**. To adjust this we use the numeric up/down buttons within the cell.
4. Select the newly added **Historical management practice** and then click the "**Edit Selected**" button. This will open a new screen that allows us to adjust the crops grown and the management during this period.
5. Click on the "**Barley**" crop under the "**Step 2**" section. Change the crop type to '**Wheat**' and on the '**General**' tab change the yield to **3,500 kg/ha**. We will keep the other settings unchanged.
6. We also need to remove the "**Hay mixed**" crops from this historical period. Click the 'x' icon beside each of the "**Hay mixed**" crops under the "**Step 2**" section. Clicking the 'x' icon will remove these crops from the rotation for this period of time.
7. Click "**Ok**" to save adjustments we just made to this field.
8. Repeat these same steps so that the other fields in this rotation also have continuous wheat from **1985 to 2000** using the same steps we used for the first field.




Figure 21: Customized Timeline Screen




Figure 22: Adjusted start and end year for productions systems on the timeline screen.

Crop rotation #1 [Field #1] - Barley (Historical management practice)

Step 1: Define the general properties for this field

Total area of this field :	70 ▲▼	(ha)
Start year of field history :	1985 ▲▼	
End year :	2000 ▲▼	

Step 2: Click the 'Add Crop' button to describe the crops grown on this field. Enter one crop for continuous fields (i.e. continuous wheat) and the first phase for crop rotations (i.e. wheat, fallow). Holos will use this entry to populate the field history.

	Crop	Winter/Cover/Undersown Crop
>	Wheat	None

Step 3: Adjust details of the crop (wheat) grown on this field

Show Additional Information :

General
Fertilizer
Manure
Winter & Cover Crops
Residue
Economics

General Properties

Amount of irrigation, pesticide usage, and yield for the selected crop can be entered on this tab. Harvested yield can be entered as either wet weight or dry weight. Additional information can be entered by clicking on the 'Show Additional Information' button.

Yield (wet weight) :	3,500 ▲▼		<i>(kg ha⁻¹)</i>
Yield (dry weight) :	3,080 ▲▼		<i>(kg DM ha⁻¹)</i>
Moisture content of crop :	12 ▲▼		(%)
Tillage :	Intensive tillage ▼		
Harvest method :	Cash crop ▼		
Amount of irrigation :	0 ▲▼		(mm)
Number of pesticide passes :	0 ▲▼		

Figure 23: Editing crops in a historical period of the rotation.

Details Screen

Click the **"Next"** button to go forward to the details screen.

To avoid the requirement that a user needs to provide crop yields going back to 1985 for each field on the farm, the model will use Stats Canada reported crop yields as defaults (where available). The model allows the user to calculate how changes in crop type, yield, tillage, residue management, manure, irrigation or fallow will result in changes to soil carbon.

We will adjust this grid so that we can view the above ground and below ground carbon inputs for our wheat field and then we will adjust the crop yield for one specific year.

1. We will set a filter on the first column named **'Field name'** so that we only display information for our **wheat and hairy vetch field**. Beside the column heading, click the **'funnel'** icon to set a filter. Check the box beside **Wheat & hairy vetch**.
2. On the far left of this screen, click the **"Enable Columns"** sidebar (located near the "Field name" column).
3. Place a check beside **Above ground carbon input** to show the column and remove the check beside the **Notes** column to hide it.
4. Click the **Enable Columns** sidebar again to collapse it.
5. We can now (optionally) adjust the yields for our wheat field for any given year if actual measured yields are available.
6. Adjust the yield for **1987** to be **4,100 kg/ha**.
7. Note that Holos has updated the above ground carbon inputs for this.

FileSettingsToolsViewHelp

Details

Use this screen to provide additional details for your farming operations. This screen is optional and may be skipped by clicking "Next"

Cropping System

Reload Data From Previous Screen

Yield Assignment Method:Average Yield

Field name	Year	Crop	Yield (kg ha ⁻¹)	Above ground carbon input (kg ha ⁻¹)	Below ground carbon input (kg ha ⁻¹)
Wheat & Hairy Vetch	1980	Hairy Vetch (Vicia villosa roth)	3000.00	1350.000	670.958
Wheat & Hairy Vetch	1981	Wheat	3000.00	52029.197	1914.686
Wheat & Hairy Vetch	1982	Hairy Vetch (Vicia villosa roth)	3000.00	1350.000	670.958
Wheat & Hairy Vetch	1983	Wheat	3000.00	52029.197	1914.686
Wheat & Hairy Vetch	1984	Hairy Vetch (Vicia villosa roth)	3000.00	1350.000	670.958
Wheat & Hairy Vetch	1985	Wheat	3000.00	54729.197	13256.602
Wheat & Hairy Vetch	1985	Hairy Vetch (Vicia villosa roth)	3000.00	1350.000	670.958
Wheat & Hairy Vetch	1986	Wheat	3000.00	54729.197	13256.602
Wheat & Hairy Vetch	1986	Hairy Vetch (Vicia villosa roth)	3000.00	1350.000	670.958
Wheat & Hairy Vetch	1987	Wheat	4100.00	3548.881	1615.349
Wheat & Hairy Vetch	1987	Hairy Vetch (Vicia villosa roth)	3000.00	1350.000	670.958
Wheat & Hairy Vetch	1988	Wheat	3000.00	54729.197	13256.602
Wheat & Hairy Vetch	1988	Hairy Vetch (Vicia villosa roth)	3000.00	1350.000	670.958
Wheat & Hairy Vetch	1989	Wheat	3000.00	54729.197	13256.602
Wheat & Hairy Vetch	1989	Hairy Vetch (Vicia villosa roth)	3000.00	1350.000	670.958
Wheat & Hairy Vetch	1990	Wheat	3000.00	54729.197	13256.602
Wheat & Hairy Vetch	1990	Hairy Vetch (Vicia villosa roth)	3000.00	1350.000	670.958
Wheat & Hairy Vetch	1991	Wheat	3000.00	54729.197	13256.602
Wheat & Hairy Vetch	1991	Hairy Vetch (Vicia villosa roth)	3000.00	1350.000	670.958
Wheat & Hairy Vetch	1992	Wheat	3000.00	54729.197	13256.602
Wheat & Hairy Vetch	1992	Hairy Vetch (Vicia villosa roth)	3000.00	1350.000	670.958
Wheat & Hairy Vetch	1993	Wheat	3000.00	54729.197	13256.602
Wheat & Hairy Vetch	1993	Hairy Vetch (Vicia villosa roth)	3000.00	1350.000	670.958
Wheat & Hairy Vetch	1994	Wheat	3000.00	54729.197	13256.602
Wheat & Hairy Vetch	1994	Hairy Vetch (Vicia villosa roth)	3000.00	1350.000	670.958
Wheat & Hairy Vetch	1995	Wheat	3000.00	54729.197	13256.602
Wheat & Hairy Vetch	1995	Hairy Vetch (Vicia villosa roth)	3000.00	1350.000	670.958

Back

Next

Canada

851003 (Portage, Manitoba)Total emissions: 34,502.24 (Mg CO2e)Loss: \$ 111,671.08Multi-Year Mode

Figure 24: Details screen

Page 30

Discover results

Click the **"Next"** button to move to the final results report. Results will now be displayed in a variety of reports and charts.

1. Click on the tab named **Emissions Pie Chart**

Starting with the **Emissions pie chart** we can see an overall breakdown of the enteric CH₄, manure CH₄, direct and indirect N₂O. We are also able to see a detailed breakdown of the sources of these emissions.

2. Click the **"Yes"** button beside **Show details**

We can see that the biggest source of emissions from our farm is the cow-calf component. If you hover your mouse pointer over any slice of this chart you can get an isolated look at the different emission sources.

3. Click on the tab named **Detailed Emission Report**

The **Detailed Emission Report** will display a monthly or annual GHG emission report. The detailed emission report will report on enteric methane, manure methane, direct & indirect N₂O, and CO₂ emissions from the farm.

Click the **Report Format (Monthly)** button to switch to a monthly report. Now we can see a monthly breakdown of all emissions from the farm and the emission source.

In the **Unit of measurements** drop-down menu, you can choose to have the results displayed as CO₂ equivalents (CO₂e) or as unconverted greenhouse gas (GHG), and you can also choose the unit of measurement as either tonnes or kilograms.

The **Estimates of Production** report provides total harvest yields, amount of land applied manure, and estimates of milk production for dairy components.

The **Feed Estimate** report provides an estimate of dry matter intake based on energy requirements of the animal and the energy in the feed.




Figure 25: Detailed emissions report.




Figure 26: Estimates of production report.




Figure 27: Feed estimate report.

Soil carbon modelling results

On the results screen we can see the change in soil carbon over time by clicking the "Multiyear Carbon Modelling" tab. This tab displays a graph showing the change in soil carbon over time for each one of our fields.

For each field on the graph, you can hover your mouse over the series to get more information for each historical year of the field.

If we click on one of these points, we can then view a more detailed breakdown of these results. We can also export this data by clicking the "Export to Excel" button.




Figure 28: Carbon report section. Allows switching between graph and table format.

Finally...

Whole-systems approach

An ecosystem consists of not only the organisms and the environment they live in but also the interactions within and between. A whole systems approach seeks to describe and understand the entire system as an integrated whole, rather than as individual components. This holistic approach can be very complex and describing the process can be difficult. One method to conceptualize a whole system is with a mathematical model.

The whole-systems approach ensures the effects of management changes are transferred throughout the entire system to the resulting net farm emissions. In some cases, reducing one GHG will actually increase the emissions of another. The whole-systems approach avoids potentially ill-advised practices based on preoccupation with one individual GHG.

To download Holos, for more information, or to access a recent list of Holos related publications, visit:
www.agr.gc.ca

To contact us, email: aafc.holos.acc@canada.ca