

BlueCarbon R package: Estimation of Organic Carbon Stocks and Sequestration Rates From Soil Core Data

Nerea Piñeiro-Juncal¹, Julen Astigarraga^{2,3}, Valentina Costa^{4,5,6}, Márcio Martins⁷, and Francisco Rodríguez-Sánchez⁸

1 Centro de Investigacions Mariñas da Universidade de Vigo. Departamento de Xeociencias Mariñas e Ordenación do Territorio, Facultade de Ciencias do Mar, Campus Lagoas Marcosende, Universidade de Vigo, Vigo, Spain **2** Universidad de Alcalá, Grupo de Ecología Forestal y Restauración (FORECO), Departamento de Ciencias de la Vida, Spain **3** Department of Physical Geography and Ecosystem Science, Lund University, Lund, Sweden **4** Stazione Zoologica Anton Dohrn – CRIMAC, Calabria Marine Centre, Department of Integrative Marine Ecology, Amendolara (CS), Italy **5** National Institute of Oceanography and Applied Geophysics – OGS, Italy **6** NBFC, National Biodiversity Future Center, Palermo, Italy **7** Centro de Ciências do Mar do Algarve (CCMAR/CIMAR LA), Campus de Gambelas, Universidade do Algarve, 8005-139 Faro, Portugal **8** Departamento de Biología Vegetal y Ecología, Facultad de Biología, Universidad de Sevilla, Spain

DOI:

Software

- [Review](#) ↗
- [Repository](#) ↗
- [Archive](#) ↗

Submitted:

Published:

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License (CC-BY).

Correspondence:

Nerea Piñeiro-Juncal, University of Vigo, Vigo, Spain. Email: np.juncal@gmail.com

Summary

BlueCarbon facilitates the estimation of organic carbon stocks and sequestration rates from soil and sediment cores in depositional environments. It includes seven main functions to (1) estimate core compaction, (2) correct core compaction, (3) estimate sample thickness, (4) estimate organic carbon content from organic matter content, (5) estimate organic carbon stocks and (6) sequestration rates, and (7) visualize the error in stock extrapolation.

Statement of Need

Coastal blue carbon ecosystems have earned significant attention for their role as organic carbon sinks. Over the past decade, publications on blue carbon research have grown exponentially (Quevedo, Uchiyama, & Kohsaka, 2023). While soil samples can be collected by different methods, estimation methodologies remain fairly homogeneous, following the protocols published by the Blue Carbon initiative (Howard, Hoyt, Isensee, Pidgeon, & Telszewski, 2014). Despite the increasing use of R among blue carbon researchers, there are no specialized R packages dedicated to these calculations. *BlueCarbon* aims to standardize and automate the main estimations for calculating soil and sediment blue carbon stocks and sequestration rates from raw field and laboratory data.

Design

BlueCarbon contains seven main functions (Fig. 1) to deal with core compaction (to estimate and mathematically correct core compaction), transform laboratory data (to