

The great acceleration of plant phenological shifts

Authors: Y. Vitasse^{1,2}, F. Baumgarten¹, C. Zohner³, T. Rutishauser², B. Pietragalla⁴, R.
Gehrig⁴, J. Dai^{5,6,7}, H. Wang⁵, Y. Aono⁸, T.H. Sparks^{9,10}

¹WSL Swiss Federal Institute for Forest, Snow and Landscape Research, Birmensdorf, Switzerland

²Swiss Academy of Sciences (SCNAT), Bern, Switzerland.

³Institute of Integrative Biology, ETH Zurich (Swiss Federal Institute of Technology), Switzerland

⁴Federal Office of Meteorology and Climatology MeteoSwiss, Zurich, Switzerland

⁵Key Laboratory of Land Surface Pattern and Simulation, Institute of Geographic Sciences and Natural
Resources Research, Chinese Academy of Sciences, Beijing, China

⁶University of Chinese Academy of Sciences, Beijing, China

⁷China-Pakistan Joint Research Center on Earth Sciences, CAS-HEC, Islamabad 45320, Pakistan

⁸Graduate School of Life and Environmental Sciences, Osaka Prefecture University, Sakai, Japan

⁹Department of Zoology, Poznań University of Life Sciences, 60-625 Poznań, Poland & Museum of Zoology,

¹⁰University of Cambridge, Cambridge CB2 3EN, UK

Running title: The great phenological acceleration

To the Editor — The world’s longest time series of plant blooming and leaf-out in spring
reveal unprecedented shifts since the middle of the 1980s in line with the acceleration of
global warming. These long-term time series provide powerful evidence of the impact of
global warming on life on Earth and can help raising awareness among citizens, decision
makers, and future generations of the urgent need to mitigate greenhouse gas emissions.

Submitted to *Nature Climate Change* as *correspondence* format

With the rise of *Homo sapiens*, the Earth has experienced drastic landscape changes and biodiversity losses. For example, during the prehistoric and preindustrial periods, forest cover and biodiversity declined massively in the world's most populated regions due to the rise of agriculture and the increasing need for firewood and building materials^{1,2}. These abrupt changes have been highlighted by the scientific community through paleoecological and archaeological records and modelling, without being directly visible or understandable to the general public because the sources do not constitute tangible continuous records. More recently, since the Industrial Revolution, human activities have led to massive changes in climatic, biotic and socio-economic indicators, a period now referred to as the "Anthropocene"³. These changes accelerated considerably after the 1950s, a period often referred to as "the great acceleration"⁴, with particularly strong ecological and biogeochemical changes since the 1980s⁵. However, unlike the pre-Anthropocene changes, recent climatic changes and seasonal plant activity have been recorded continuously for several centuries in a few locations and can be used to demonstrate the impact of human activities on global warming and its consequences for living organisms. In fact, the phenological cycle of plants, is strongly temperature-dependent, and spring phenological events (e.g. flowering or egg hatching in birds) in boreal, temperate and subtropical regions have progressively advanced over recent warmer decades^{6,7}. However, phenological observations are very limited before 1950, and we thus lack important information on whether current trends during the Anthropocene's great acceleration are unique over longer time scales. Here we present the five world's longest phenological time series known to date to provide evidence for an abrupt change linked to the current acceleration of global warming. The longest phenological series ever recorded is the blooming of cherry trees (*Prunus jamasakura*) in Kyoto, Japan, with observations extracted from old diaries and chronicles stretching back to the year 812 AD⁸. The longest European time series was recorded by the Marham family in the southeast of the UK and includes leaf-out dates of several common tree species from 1736 to 1958⁹. Fortunately, also in lowland southeast UK, a unique series of oaks budburst (*Quercus robur*) has been recorded since 1950 by a single observer, J. Combes, allowing the Marham's series to be extended to the present day. The three other series presented here are reconstructed flowering dates of three Chinese woody species (*Amygdalus davidiana*, *Cercis chinensis* and *Paeonia suffruticosa*) based on phenological records from old diaries since 1834¹⁰, budburst dates of a horse chestnut (*Aesculus hippocastanum*) observed since 1808 in Geneva, Switzerland, by the Grand Conseil de la République et canton de Genève, and flowering dates of a cherry tree (*Prunus avium*) recorded in the north of Switzerland since

1894 by the Landwirtschaftliches Zentrum Ebenrain, Sissach and the Federal Office of Meteorology and Climatology MeteoSwiss.

Remarkably, the five world's longest plant phenological series showed fairly stable flowering and leaf-out dates throughout the 19th century (Fig. 1). Spring phenological events started to advance in the first half of the 20th century in Switzerland and Japan, in parallel with the temperature increase. No major shift was observed during the period 1950–1985, except for the blooming of cherry tree in Kyoto which continued to advance, matching the increase in spring temperature observed in this location (Fig. 1). In contrast, the strongest advance of spring phenology was detected in the middle of the 1980s for most of the series, again consistent with the accelerated warming trend observed across the Northern Hemisphere¹¹. On average, spring phenology occurred 6 (China) to 30 days (Switzerland) earlier in the last 36 years (1985–2020) compared to the period before 1950 (Fig. 1). Notably, the blooming of cherry trees in Kyoto in spring 2021 was the earliest date ever recorded in more than 1,200 years, further underscoring the ongoing, rapid phenological shifts.

The acceleration of global warming since 1985 (Fig. 1A) led to historically unprecedented advances in spring phenology in all series that spanned over two centuries (or- for the Japan series- even over a millenia). Because of the strong relationship between spring phenology and temperature (Fig. 1B), we argue that plant phenological shifts are an important bioindicator of the current acceleration of global warming. Strikingly, the series presented here are not situated in those parts of the world where warming trends have been the most pronounced, and we can therefore expect even stronger phenological shifts towards the circumarctic and central Asian regions where more than 2.5°C warming has occurred over 150 years¹². To the best of our knowledge, no continuous long-term phenology series exist for North and South America, Africa and Australia. However, recent and promising efforts have been made to combine historical information from herbarium specimens with contemporary records, especially in the northeastern US¹³.

Unlike the rise of CO₂ in the atmosphere, seasonal biological events, such as leaf-out or blooming, are easily identifiable by everyone including children, and represent concrete and visible evidence of how climate change is affecting the ecosystems around us. We therefore believe that these long-term series are particularly useful in communication to raise citizen awareness about the urgency to mitigate climate change and to foster public engagement in citizen science, in education to further sensitize the coming generations, and in politics to

promote sustainable environmental decisions. Such abrupt changes in the timing of key phenological events can have multiple implications for ecosystem functioning, species interactions via the food-web¹⁴, and global carbon balance¹⁵, and should therefore attract greater attention from society.

Acknowledgment

We are grateful to all the recorders and institutions who provided phenological observations for the different long-term series for decades and centuries. YV is supported by the Swiss National Science Foundation SNF (research grant 315230_192712) and the SwissForestLab (research grant SFL-17 P2). CMZ is funded by the Ambizione grant PZ00P3_193646.

Competing interests

The author declares no competing interests.

References

1. Kaplan JO, Krumhardt KM, Zimmermann N. The prehistoric and preindustrial deforestation of Europe. *Quaternary Science Reviews* 2009, **28**(27-28): 3016-3034.
2. Zheng Z, Ma T, Roberts P, Li Z, Yue Y, Peng H, *et al.* Anthropogenic impacts on Late Holocene land-cover change and floristic biodiversity loss in tropical southeastern Asia. *Proceedings of the National Academy of Sciences* 2021, **118**(40).
3. Lewis SL, Maslin MA. Defining the Anthropocene. *Nature* 2015, **519**(7542): 171–180.
4. Steffen W, Broadgate W, Deutsch L, Gaffney O, Ludwig C. The trajectory of the Anthropocene: the great acceleration. *The Anthropocene Review* 2015, **2**(1): 81–98.
5. Ripple WJ, Wolf C, Newsome TM, Barnard P, Moomaw WR. World scientists' warning of a climate emergency. *Bioscience* 2019.
6. Cohen JM, Lajeunesse MJ, Rohr JR. A global synthesis of animal phenological responses to climate change. *Nature Climate Change* 2018, **8**(3): 224–228.
7. Vitasse Y, Ursenbacher S, Klein G, Bohnenstengel T, Chittaro Y, Delestrade A, *et al.* Phenological and elevational shifts of plants, animals and fungi under climate change in the European Alps. *Biological Reviews* 2021, **96**(5): 1816-1835.
8. Aono Y, Kazui K. Phenological data series of cherry tree flowering in Kyoto, Japan, and its application to reconstruction of springtime temperatures since the 9th century. *International Journal of Climatology: A Journal of the Royal Meteorological Society* 2008, **28**(7): 905–914.
9. Sparks TH, Carey PD. The responses of species to climate over 2 centuries - an analysis of the Marsham phenological record, 1736-1947. *Journal of Ecology* 1995, **83**(2): 321–329.

10. Ge Q, Wang H, Zheng J, Rutishauser T, Dai J. A 170 year spring phenology index of plants in eastern China. *Journal of Geophysical Research-Biogeosciences* 2014, **119**(3): 301-311.
11. IPCC. *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press: Cambridge, United Kingdom and New York, NY, USA, 2021.
12. Post E, Steinman BA, Mann ME. Acceleration of phenological advance and warming with latitude over the past century. *Scientific Reports* 2018, **8**(1): 3927.
13. Primack RB, Miller-Rushing AJ. Uncovering, Collecting, and Analyzing Records to Investigate the Ecological Impacts of Climate Change: A Template from Thoreau's Concord. *Bioscience* 2012, **62**(2): 170-181.
14. Kharouba HM, Ehrlén J, Gelman A, Bolmgren K, Allen JM, Travers SE, *et al.* Global shifts in the phenological synchrony of species interactions over recent decades. *Proceedings of the National Academy of Sciences* 2018, **115**(20): 5211–5216.
15. Richardson AD, Keenan TF, Migliavacca M, Ryu Y, Sonnentag O, Toomey M. Climate change, phenology, and phenological control of vegetation feedbacks to the climate system. *Agricultural and Forest Meteorology* 2013, **169**: 156–173.
16. Parker DE, Legg TP, Folland CK. A new daily central England temperature series, 1772–1991. *International Journal of Climatology* 1992, **12**(4): 317-342.

Figure caption

Figure 1. The world's longest phenological series with the associated spring temperatures. A, mean air temperature from January to March in the different sites or areas where the phenological series have been conducted; B, correlation between spring temperature (January to March) and phenology for each series. Pearson correlation coefficients are indicated and phenological observations before and after 1985 are distinguished with two distinct symbols; C, flowering dates of the wild cherry tree (*Prunus avium* L.) in Liestal, Switzerland (1894–2020); D, spring flowering index derived from three woody species (flowering of *Amygdalus davidiana* (Carrière) Franch., *Cercis chinensis* Bunge and *Paeonia suffruticosa* Andrews) in China (1834–2020); E, budburst dates of horse chestnut (*Aesculus hippocastanum*) in Geneva, Switzerland (1808–2020); F, budburst dates of pedunculate oak (*Quercus robur* L.) in SE UK from the Marsham series (dark orange, 1736–1958) and J. Combes series (light orange, 1950–2020). The mean difference between the two series of observations (~3 days) indicates that the latter represent a reliable continuation of the Marsham's series; G, blooming of the yamazakura cherry tree (*Prunus jamasakura* Lindl.) in Kyoto in Japan (1500–2020, note that the series goes as far back as the year 812 AD but is shown here since the year 1500 only). In all graphs, the grey thin lines represent the annual phenology or temperatures and the thick lines represent the ten-years moving averages. In the phenological series, horizontal lines represent the periods detected by the breakpoint analysis allowing a minimum of breaks that significantly decrease the penalty value using the R-package 'changepoint' with the function 'cpt.mean' and the 'BinSeg' method and 'BIC' penalty. On the right side of each phenological series, graphs show the data distribution (violin plots and mean \pm 1 SD) for three periods, i.e. before the great acceleration (<1950), during 1950–1984 and during 1985–2020. Different letters among the different time periods indicate significant differences in spring phenology (Tukey's honestly significant difference (HSD) test). Source of temperature data: Beijing [1881–2020], cleaned and homogenized data by the NASA to account for urban effects, downloaded from <https://data.giss.nasa.gov>; Basel [1755–2020] and Geneva [1753–2020], homogenized data available from the Federal Office of Meteorology and Climatology MeteoSwiss; Central England [1659–2020], longest available instrumental record of temperature in the world¹⁶, representative of a roughly triangular area of the United Kingdom enclosed by Lancashire, London and Bristol, made available by the Hadley Centre Central England Temperature (HadCET); Kyoto [1881–2020], data available from the Japan Meteorological agency.

Figure 1.

