

















































Mid-Atlantic Benthic Habitats

| | Code | Description | Selected Species |
|--|------|---|---|
|  | 1 | Depressions and mid-position flats, shallow to moderate depth (0 - 45 m) on coarse to fine sand | Shimmy worm , Astarte, Lancelet |
|  | 2 | Flat depressions at shallow to moderate depth (0 - 45 m) in medium sand | Chevron worm , Burrowing anemone , Dog welk |
|  | 4 | Mid-position flats in shallow water (25 - 45 m) on coarse to medium sand | Bamboo worm , Thread worm , Chestnut astarte |
|  | 7 | Mid-position flats and depressions in shallow water (25 - 45 m) on medium to coarse substrate | Fringe worm , Frilled anemone , Common sea star |
|  | 25 | Depressions at moderate depths (15 - 82 m) on fine to coarse sand | Bristle worm , Moon snail, Lined anemone |
|  | 32 | Mid-position flats at shallow to moderate depths (22 - 45 m) on medium sand | Atlantic rock crab , Longnose spider crab, Common northern moon snail |
|  | 38 | Depressions in water shallow (15 - 22 m) on medium to coarse sand | Bamboo worm , Olivepit porcelain crab , Arctic paper-bubble |
|  | 44 | Depressions and mid-position flats mostly very shallow (0 - 22m), but occasionally very deep on fine to coarse sand | Jonah crab , Amethyst gemclam , Northern sea star |
|  | 64 | Depressions and mid-position flats in shallow water (15 and 22 m) on medium sand | Blood worm , Sharp-tailed cumacean , Blue mussel |
|  | 84 | All types of flats at moderate depth (22 - 82 m) on fine to medium sand | Beardworm, Sea urchin , Nutclam |
|  | 87 | Depressions and high flats in shallow water (15 - 22 m) on medium sand | Burrowing scale worm , Glass shrimp , Sand dollar |
|  | 216 | High slopes in deep water (95 - 592 m) on medium to fine sand | American lobster , Sea feather , Bobtail squid |
|  | 218 | Depressions at moderate depths (45 - 82 m) on medium to coarse sand | Feather duster worm , Bean mussel , Conrad's thracia |
|  | 219 | High flats at moderate depths (45 - 82 m) on coarse to fine sand | Prawn, Sea scallop , Mud star |
|  | 229 | High flats and depressions at shallow to deep depths (22- 592 m) on a fine to medium sand | Jonah crab , Green sea urchin , Anemone |
|  | 301 | Any seabed form at moderate to deep depths (45 -592) on any substrate | Gould's pandora , Calcareous coral , Foraminiferida |
|  | 306 | All types of flats at medium depth (45 - 82 m) on medium sand | Acadian hermit crab , Daisy brittle star |
|  | 384 | High slopes and canyons in deep water (95 - 592 m) on any substrate | Scale worm , Florida lobsterette , Royal red shrimp |
|  | 395 | Depressions and high flats at moderate depths (45 - 82 m) on fine to medium sand | Spionid mud worm, Paper clam , Sea star |
|  | 505 | Slopes and canyons in very deep water (>592 m) on silt and mud | Beardworms, Hairy sea cucumber , Stony corals |
|  | 520 | Mid position flats and depressions at moderate depths (45 - 82) on mostly coarse to occasional fine sand | Long-horned skeleton shrimp , Hatchet shell , Peanut worm |
|  | 592 | Mid-position flats at moderate depth (45 - 82 m) on medium sand | Paddle worm , Cumacea, Arctic rock borer |
|  | 768 | Depressions in very shallow water (0 - 15 m) on silt to fine sand | Mysid shrimp , Elongated macoma , Burrowing brittle star |
|  | 1223 | High flats in moderately deep water (82 - 95 m) on medium sand | Eunice worm , Striate scallop , Margined sea stars |

Southern New England Benthic Habitats

| | Code | Description | Selected Species |
|--|------|---|--|
|  | 1 | Variable settings in a wide range of depths on fine to coarse sand | n/a |
|  | 3 | Flats and slopes at moderate to very deep depths (average 128 m, min 44 m) on fine to very fine sand | n/a |
|  | 6 | High slopes and flats at moderate to deep depths (44 - 139 m) on coarse to fine sand | Aesop shrimp, Arctic rock borer, Daisy brittle star |
|  | 11 | High slopes, canyons, flats in deep water (60 – 485 m) on medium to fine sand | Parrot shrimp, Bobtail squid, Margined sea star |
|  | 25 | Flats and side slopes in very shallow to shallow water (0 - 23 m) on fine to coarse sand | Blood worm, Glass shrimp, Surfclam |
|  | 36 | Depressions and high flats in very shallow to moderate depths (0 – 75 m) on medium to coarse sand | Green crab, Bittium snail, Egg cockle |
|  | 66 | High flats and slopes in moderately deep water (75 - 139 m) on very fine to fine sand | Fringe worm, Dwarf brittlestar, Slender sea pen |
|  | 82 | All types of flats in moderately deep water (44 – 139 m) on medium to coarse sand | Sea scallop, Limpet, Green sea urchin |
|  | 109 | Depressions in very shallow water (0 - 23 m) mostly on medium to coarse sand but occasionally on silt | Bamboo worm, Bay barnicle, Common razor clam |
|  | 113 | Depressions and mid-position flats at moderate depths (23 - 44 m) on very fine sand | Paddle worm, Paranoid worm, Spaghetti-mouth worm |
|  | 200 | Depressions at very shallow to moderate depths (0 – 44 m) on very fine to medium sand | Sludge worm, Pitted baby-bubble |
|  | 223 | Mid-position flats and depressions at moderate depths (44 - 75 m) on fine to medium sand | Terebellid worm, Hatchet shell, Tube worm |
|  | 229 | Depressions in shallow depths (8.4 to 44 meter) on very fine sand | Clam worm, Cephalocarid, Lined anemone |
|  | 230 | Depressions in shallow depths (23 - 44 m) on very fine sand | Burrowing scale worm, Fan worm, Pyramid snail |
|  | 316 | Flats in shallow water (8-44 m) on very fine to medium sand | Bamboo worm, Mageloni worm, Other polychaetes |
|  | 317 | Mid-position flats at moderate depths (31 - 75 m) on fine to medium sand | Sandbar worm, Acadian hermit crab, Northern moon shell |
|  | 372 | Depressions and low slopes at moderate depths (44 – 75 m) on very fine sand | Thread worm, Nutclam, Burrowing anemone |
|  | 381 | Mid and high position flats in moderate depths (44 - 79 m) on fine to very fine sand | Fan worm, Long-horned skeleton shrimp, Sea cucumber |
|  | 387 | High slopes and flats in very deep water (>139 m) on fine sand | Beard worm, Ark shell, Sea butterfly |
|  | 390 | Depressions in shallow water (23 - 44 m) in very fine to fine sand | Feather duster worm, Fringeworm, Oval Yoldia |
|  | 437 | High flats and slopes in deep to very deep water (75 - 200 m) on fine sand | American Lobster, Jonah Crab, Northern shortfin squid |
|  | 873 | Flats and side slopes in shallow water (8 - 31 m) on very fine to medium sand | Blood worm, Bristle worm, Atlantic razor |
|  | 949 | Mid and low flats in deep water (75-139 m) on medium to fine sand | Longfin squid |
|  | 2537 | Depressions and high flats in shallow water (23 - 31 m) on very fine to fine sand | Hesion worm, Paddle worm, Atlantic rock crab |