



Fakultät für Physik

Institut für Theorie der Kondensierten Materie

Quantentransport in Spindichtesystemen mit dem Memory-Matrix-Formalismus

Masterthesis

von

Martin Lietz

27. Mai 2017 bis 27. Mai 2018

Referent: Prof. Dr. Jörg Schmalian
Korreferent: Prof. Dr. Alexander Shnirman

Ich erkläre hiermit, dass die Arbeit selbstständig angefertigt, alle benutzten Quellen und Hilfsmittel vollständig und genau angegeben und alles kenntlich gemacht wurde, das aus Arbeiten anderer unverändert oder mit Abänderungen entnommen ist.

Karlsruhe, den 22. April 2018

.....
(**Martin Lietz**)

Acknowledgments

I would like to express my thanks to my advisor Jörg Schmalian who encouraged and supported me and gave the right hints at the right time during the last year. His essential input lead to the results presented in this work.

Furthermore, I would like to thank Una Karahasanovic for all the discussions and information which became decisive for one topic of this work.

Then, I have to thank especially Jian Kang who crucially contributed to the discussion of bond currents and Rafael Fernandes for the fruitful discussions on the general topic.

Special thanks to all the colleagues of the condensed matter group at the KIT. Especially to Bhilahari Jeevanesan for all the discussions and help with several problems and to Pablo Schad for finalizing this work.

Last but not least, I have to thank my family, my mother and sister and especially Anja for all the support and encouraging words.

Contents

1	Motivation	1
2	Spin-Fermion-Model	3
3	Memory-Matrix-Formalism	5
4	Calculation	7
4.1	Infinite conductivity in systems with unbroken translation symmetry .	7
5	Conclusion	9
A	Appendix	11

1 Motivation

2 Spin-Fermion-Model

3 Memory-Matrix-Formalsim

4 Calculation

In the last chapter the memory-matrix-formalism was introduced, which give us an exact formula to calculate correlation functions. Now this formalism is used to determine the static conductivity of the spin-fermion-model, introduced in chapter (), perturbate umklapp-scattering.

make link to chapter spin-fermion-model

4.1 Infinite conductivity in systems with unbroken translation symmetry

After Drude published his theory about the electrical transport in metals [Dru00] in the beginning of the last century it is well known that a broken translation symmetry is needed to get a finite static conductivity. Because of Neother's theorem it is also well known that a unbroken symmetry always implies a conserved quantity. In the case of translation symmetry this quantity is the momentum. Phenomenas breaking the translation symmetry are for example impurity scattering, electron-electron scattering and umklapp scattering. Let us firstly assume the standard spin-fermion-model without a translation symmetry breaking perturbation. In chapter () it is showed that in the used model the momentum is conserved but the current isn't conserved. This property is needed to calculate the static conductivity.

link to chapter spin-fermion-model

In general the static conductivity is given by taking the small frequencie limit of the conductivity and the conductivity itself is given by the current-current correlation function (J-J correlation function), which reslut directly from the linear response theory.

$$\sigma_{dc} = \lim_{z \rightarrow 0} \sigma(z) = \lim_{z \rightarrow 0} \beta C_{JJ}(z) \quad (4.1)$$

The memory matrix formalism is used to calculate the J-J correlation function. Before we attend us to the calculation of the correlation function we have to think about the set of operators introduced by defining the projection operator. This set of operators has to be choosen for each calculation seperatly depending of the model and the quantity of interest. In our case we choose only a set of two operators namly the momentum and the current, because we want to figure out the influnece of the momentum on the current. That means the projector \mathcal{P} projects into the two dimensional sub-Hilbertspace, spanned P and J. Lets go back to the correlation function, defined by equation (), where the sum over k und l is implied. Because our interest is focused on the J-J correlation function each index j and l is set to J .

reference to correlation function

$$C_{JJ}(z) = \frac{i}{\beta} \left[z \delta_{iJ} + i \beta (\dot{A}_i | \hat{Q} \frac{i}{z - \hat{Q} \hat{L} \hat{Q}} \hat{Q} | \dot{A}_k) \chi_{kJ}^{-1} \right]^{-1} \chi_{iJ}, \quad (4.2)$$

Now the sum over k is performed explicitly where both contributions for J and P vanish. Let us look separately on J and P starting with the latter case. In our observed model the momentum is conserved and so the time derivative of P is zero. No more words are needed to see that the expectation value doesn't contribute. In the case of J the time derivative doesn't vanish so the two dimensional sub-Hilbertspace and the action of J in this Hilbertspace has to be considered. In the investigated system the whole current lives in the J - P Hilbertspace to any time, so no single part of J is transported out of the Hilbertspace. The appearing operator \mathcal{Q} is the inverse of \mathcal{P} and therefore projected out of the J - P Hilbertspace. Combining both statements it is clear that $\mathcal{Q}|\dot{J}\rangle = 0$. The only resulting term choosing $i = J$ is

$$\mathcal{C}_{JJ}(z) = \frac{i}{\beta} z^{-1} \chi_{JJ}(\omega = 0) = \frac{i}{z} \mathcal{C}_{JJ}(t = 0), \quad (4.3)$$

where the correlation function at $t = 0$ is given by the scalar product $(J(0)|J(0))$ defined in equation (4.1). During the motivation of the previous chapter we explained that each observable can be split in one secular and one non-secular part. This is equivalent with splitting a vector in a parallel and a perpendicular component, respectively.

$$|J\rangle = |J_{\parallel}\rangle + |J_{\perp}\rangle \quad (4.4)$$

What does this mean in physical language? In the investigated system the current isn't conserved, but a part of it is. This part is represented by the secular part and has to be parallel with the momentum. Therefore the projection from J at P yields the parallel component of J .

$$|J_{\parallel}\rangle = \mathcal{P}|J\rangle = \frac{|P\rangle\langle P|}{\langle P|P\rangle}|J\rangle = \frac{\chi_{PJ}}{\chi_{PP}}|P\rangle \quad (4.5)$$

$$\mathcal{C}_{JJ}(t = 0) = (J(0)|J(0)) = (J_{\parallel}|J_{\parallel}) + (J_{\perp}|J_{\perp}) \quad (4.6)$$

$$\mathcal{C}_{JJ}(t = 0) = \frac{|\chi_{PJ}|^2}{|\chi_{PP}|^2} \mathcal{C}_{PP}(t = 0) + (J_{\perp}|J_{\perp}) \quad (4.7)$$

$$\mathcal{C}_{JJ}(z) = \frac{i}{z\beta} \frac{|\chi_{PJ}|^2}{|\chi_{PP}|} + \frac{i}{z} (J_{\perp}|J_{\perp}) \quad (4.8)$$

$$\sigma(z) = \frac{|\chi_{PJ}|^2}{|\chi_{PP}|} \frac{i}{z} + \sigma_{\text{reg}}(z) \quad (4.9)$$

where the regular conductivity $\sigma_{\text{reg}}(z) = \frac{i\beta}{z} (J_{\perp}|J_{\perp})$ is introduced. Setting $z = \omega + i\eta$, where the limit $\eta \rightarrow 0$ is implied.

$$\sigma_{\text{dc}} = \frac{|\chi_{PJ}|^2}{|\chi_{PP}|} \left(\mathcal{P} \frac{i}{\omega} + \pi \delta(\omega) \right) \quad (4.10)$$

where \mathcal{P} symbolized that the principal value is taken.

5 Conclusion

A Appendix

Todo list

make link to chapter spin-fermion-model	7
link to chapter spin-fermion-model	7
reference to correlation function	7
reference to scalar product	8

Bibliography

- [Dru00] P. Drude. „Zur Elektronentheorie der Metalle“. In: *Annalen der Physik* 306.3 (1900), pp. 566–613. DOI: 10.1002/andp.19003060312.

