



Fakultät für Physik

Institut für Theorie der Kondensierten Materie

Quantentransport in Spindichtesystemen mit dem Memory-Matrix-Formalismus

Masterthesis

von

Martin Lietz

27. Mai 2017 bis 27. Mai 2018

Referent: Prof. Dr. Jörg Schmalian
Korreferent: Prof. Dr. Alexander Shnirman

Ich erkläre hiermit, dass die Arbeit selbstständig angefertigt, alle benutzten Quellen und Hilfsmittel vollständig und genau angegeben und alles kenntlich gemacht wurde, das aus Arbeiten anderer unverändert oder mit Abänderungen entnommen ist.

Karlsruhe, den 20. April 2018

.....
(**Martin Lietz**)

Acknowledgments

I would like to express my thanks to my advisor Jörg Schmalian who encouraged and supported me and gave the right hints at the right time during the last year. His essential input lead to the results presented in this work.

Furthermore, I would like to thank Una Karahasanovic for all the discussions and information which became decisive for one topic of this work.

Then, I have to thank especially Jian Kang who crucially contributed to the discussion of bond currents and Rafael Fernandes for the fruitful discussions on the general topic.

Special thanks to all the colleagues of the condensed matter group at the KIT. Especially to Bhilahari Jeevanesan for all the discussions and help with several problems and to Pablo Schad for finalizing this work.

Last but not least, I have to thank my family, my mother and sister and especially Anja for all the support and encouraging words.

Contents

1	Motivation	1
2	Spin-Fermion-Model	3
3	Memory-Matrix-Formalism	5
4	Calculation	7
4.1	Infinite conductivity in systems with unbroken translation symmetry .	7
5	Conclusion	9
A	Appendix	11

1 Motivation

2 Spin-Fermion-Model

3 Memory-Matrix-Formalsim

4 Calculation

In the last chapter the memory-matrix-formalism was introduced, which give us an exact formula to calculate correlation functions. Now this formalism is used to determine the static conductivity of the spin-fermion-model, introduced in chapter , perturbate umklapp-scattering.

make link to chapter spin-fermion-model

4.1 Infinite conductivity in systems with unbroken translation symmetry

After Drude published his theory about the electrical transport in metals [Dru00] in the beginning of the last century it is well known that a broken translation symmetry is needed to get a finite static conductivity. Because of Neother's theorem it is also well known that a unbroken symmetry always implies a conserved quantity. In the case of translation symmetry this quantity is the momentum. Phenomenas breaking the translation symmetry are for example impurity scattering, electron-electron scattering and umklapp scattering. Let us firstly assume the standard spin-fermion-model without a translation symmetry breaking pertubation. In chapter it is showed that in the used model the momentum is conserved but the currunt isn't conserved. This property is needed to calculate the static conductivity.

link to chapter spin-fermion-model

The static conductivity is given by the conductivity taking the limit for small frequencies and the conductivity itself is given by the current-current correlation function.

$$\sigma_{\text{dc}} = \lim_{z \rightarrow 0} \sigma(z) = \lim_{z \rightarrow 0} \beta \mathcal{C}_{\text{JJ}}(z) \quad (4.1)$$

In the notation of the memory-matrix-formalism the current-current correlation function ist given by

$$\mathcal{C}_{\text{JJ}}(z) = \frac{i}{\beta} \left[z \delta_{iJ} + i\beta \left(\dot{A}_i \left| \hat{Q} \frac{i}{z - \hat{Q} \hat{L} \hat{Q}} \hat{Q} \right| \dot{A}_k \right) \chi_{kJ}^{-1} \right]^{-1} \chi_{iJ}, \quad (4.2)$$

where a sum over k is implied.

$$\mathcal{C}_{\text{JJ}}(z) = \frac{i}{\beta} z^{-1} \chi_{\text{JJ}}(\omega = 0) = \frac{i}{z} \mathcal{C}_{\text{JJ}}(t = 0) \quad (4.3)$$

$$|\text{J}\rangle = |\text{J}_{\parallel}\rangle + |\text{J}_{\perp}\rangle \quad (4.4)$$

$$|J_{\parallel}\rangle = \hat{P}|J\rangle = \frac{|P\rangle\langle P|}{\langle P|P\rangle}|J\rangle = \frac{\chi_{PJ}}{\chi_{PP}}|P\rangle \quad (4.5)$$

$$\mathcal{C}_{JJ}(t=0) = (J(0)|J(0)) = (J_{\parallel}|J_{\parallel}) + (J_{\perp}|J_{\perp}) \quad (4.6)$$

$$\mathcal{C}_{JJ}(t=0) = \frac{|\chi_{PJ}|^2}{|\chi_{PP}|^2}\mathcal{C}_{PP}(t=0) + (J_{\perp}|J_{\perp}) \quad (4.7)$$

$$\mathcal{C}_{JJ}(z) = \frac{i}{z\beta} \frac{|\chi_{PJ}|^2}{|\chi_{PP}|} + \frac{i}{z}(J_{\perp}|J_{\perp}) \quad (4.8)$$

$$\sigma(z) = \frac{|\chi_{PJ}|^2}{|\chi_{PP}|} \frac{i}{z} + \sigma_{\text{reg}}(z) \quad (4.9)$$

where the regular conductivity $\sigma_{\text{reg}}(z) = \frac{i\beta}{z}(J_{\perp}|J_{\perp})$ is introduced. Setting $z = \omega + i\eta$, where the limit $\eta \rightarrow 0$ is implied.

$$\sigma_{\text{dc}} = \frac{|\chi_{PJ}|^2}{|\chi_{PP}|} \left(\mathcal{P} \frac{i}{\omega} + \pi\delta(\omega) \right) \quad (4.10)$$

where \mathcal{P} symbolized that the prinzipal value is taken.

5 Conclusion

A Appendix

Todo list

make link to chapter spin-fermion-model	7
link to chapter spin-fermion-model	7

Bibliography

- [Dru00] P. Drude. „Zur Elektronentheorie der Metalle“. In: *Annalen der Physik* 306.3 (1900), pp. 566–613. DOI: 10.1002/andp.19003060312.

