


Отчёт по протоколу

Набор данных: Файл данных 606be...
Протокол: Первичка — полный отчёт по всем показателям (comparisons vs arm)
ID набора: 606be8ec-47ef-4828-9151-d78c45eef41b

Шаг: cmp_age

Метод: Mann-Whitney U Test
p-value: 0.7595
Статистика: 228.500
Эффект: rbc -0.06


Тип: c



Шаг: cmp_sex

Метод: Chi-Square Test of Independence
p-value: 0.7540
Статистика: 0.098

Тип: c




Вывод: A Chi-Square test of independence was performed to examine the relation between sex and arm. The relation between these variables was not significant ($p = 0.754$). sex appears to be independent of arm.

Шаг: cmp_сад_v1

Тип: c


Метод: Student's t-test (Independent)
p-value: 0.4010
Статистика: 0.848
Эффект: cohen-d 0.26
ДИ эффекта: [-6.02, 14.74]
Мощность: 0.13
BF10: 0.398



Шаг: cmp_дад_v1

Тип: c

Метод: Student's t-test (Independent)
p-value: 0.3440
Статистика: 0.957
Эффект: cohen-d 0.29
ДИ эффекта: [-3.18, 8.90]
Мощность: 0.15
BF10: 0.43



Вывод: An independent Student's t-test (Independent) was conducted to determine if there were differences in ДАД V1 between groups defined by arm. The analysis revealed no statistically significant difference between the groups ($p = 0.344$, Cohen's $d = 0.29$ indicates a small effect).

Шаг: cmp__сад_v2


Тип: c

Метод: Mann-Whitney U Test

p-value: 0.9531

Статистика: 245.000

Эффект: rbc 0.01



Шаг: cmp__дад_v2


Тип: c

Метод: Mann-Whitney U Test

p-value: 0.1694

Статистика: 300.500

Эффект: rbc 0.24




Вывод: An independent Mann-Whitney U Test was conducted to determine if there were differences in ДАД V2 between groups defined by arm. The analysis revealed no statistically significant difference between the groups ($p = 0.169$, $r = 0.24$ indicates a small effect).

Шаг: cmp_ушобп_часть_2_повседневная_активность_баллы_v2

Тип: c

Метод: Student's t-test (Independent)
p-value: 0.1539
Статистика: 1.452
Эффект: cohen-d 0.44
ДИ эффекта: [-0.55, 3.37]
Мощность: 0.29
BF10: 0.69



Вывод: An independent Student's t-test (Independent) was conducted to determine if there were differences in УШОБП часть 2 «Повседневная активность» баллы V2 between groups defined by arm. The analysis revealed no statistically significant difference between the groups ($p = 0.154$, Cohen's $d = 0.44$ indicates a small effect).

Шаг: cmp_ушобп_часть_3_двигательные_функции_баллы_v2

Тип: c

Метод: Welch's T-Test
p-value: 0.9532
Статистика: -0.059
Эффект: cohen-d 0.02
ДИ эффекта: [-3.22, 3.04]
Мощность: 0.05
BF10: 0.298




Вывод: An independent Welch's T-Test was conducted to determine if there were differences in УШОБП часть 3 «Двигательные функции» баллы V2 between groups defined by arm. The analysis revealed no statistically significant difference between the groups ($p = 0.953$, Cohen's $d = 0.02$ indicates a trivial effect).

Шаг: cmp_шкала_депрессии_тревоги_и_стресса_dass_21_баллы_v2

Тип: c

Метод: Student's t-test (Independent)
p-value: 0.0074
Статистика: -2.814
Эффект: cohen-d 0.85
ДИ эффекта: [-8.59, -1.41]
Мощность: 0.78
BF10: 6.2




Вывод: An independent Student's t-test (Independent) was conducted to determine if there were differences in Шкала депрессии тревоги и стресса DASS-21 баллы V2 between groups defined by arm. There was a statistically significant difference between the groups ($p = 0.007$, Cohen's $d = 0.85$ indicates a large effect). Specifically, the Шкала депрессии тревоги и стресса DASS-21 баллы V2 in the active group ($M = 30.73$) was significantly lower than in the placebo group ($M = 35.73$).

Шаг: cmp_шкала_оценки_устоляемости_при_бп_баллы_v2

Тип: c

Метод: Student's t-test (Independent)
p-value: 0.2187
Статистика: -1.249
Эффект: cohen-d 0.38
ДИ эффекта: [-13.44, 3.16]
Мощность: 0.23
BF10: 0.555




Вывод: An independent Student's t-test (Independent) was conducted to determine if there were differences in Шкала оценки утомляемости при БП баллы V2 between groups defined by arm. The analysis revealed no statistically significant difference between the groups ($p = 0.219$, Cohen's $d = 0.38$ indicates a small effect).

Шаг: cmp_шкала_оценки_сонливости_эпуорта_баллы_v2

Тип: c

Метод: Mann-Whitney U Test
p-value: 0.8316
Статистика: 232.500
Эффект: rbc -0.04



Вывод: An independent Mann-Whitney U Test was conducted to determine if there were differences in Шкала оценки сонливости Эпуорта баллы V2 between groups defined by arm. The analysis revealed no statistically significant difference between the groups ($p = 0.832$, $r = -0.04$ indicates a trivial effect).

Шаг: cmp_шкала_апатии_старкстейна_баллы_v2

Тип: c

Метод: Student's t-test (Independent)
p-value: 0.1226
Статистика: 1.576
Эффект: cohen-d 0.48
ДИ эффекта: [-0.59, 4.77]
Мощность: 0.34
BF10: 0.799




Вывод: An independent Student's t-test (Independent) was conducted to determine if there were differences in Шкала апатии Старкстейна баллы V2 between groups defined by arm. The analysis revealed no statistically significant difference between the groups ($p = 0.123$, Cohen's $d = 0.48$ indicates a small effect).

Шаг: cmp_тест_струпа_v2_1

Тип: c


Метод: Mann-Whitney U Test
p-value: 0.4626
Статистика: 237.500
Эффект: rbc 0.14



Шаг: cmp_тест_струпа_v2_2

Тип: c

Метод: Mann-Whitney U Test
p-value: 0.0512
Статистика: 284.000
Эффект: rbc 0.36




Вывод: An independent Mann-Whitney U Test was conducted to determine if there were differences in Тест Струпа V2 2 between groups defined by arm. The analysis revealed no statistically significant difference between the groups ($p = 0.051$, $r = 0.36$ indicates a medium effect).

Шаг: cmp__тест_струпа_v2_3

Тип: c

Метод: Student's t-test (Independent)
p-value: 0.2702
Статистика: 1.118
Эффект: cohen-d 0.35
ДИ эффекта: [-1.73, 5.99]
Мощность: 0.19
BF10: 0.503



Шаг: cmp__индекс_интерференции_v2

Тип: c

Метод: Mann-Whitney U Test
p-value: 0.0330
Статистика: 127.000
Эффект: rbc -0.39




Вывод: An independent Mann-Whitney U Test was conducted to determine if there were differences in Индекс интерференции V2 between groups defined by arm. There was a statistically significant difference between the groups ($p = 0.033$, $r = -0.39$ indicates a medium effect). Specifically, the Индекс интерференции V2 in the active group ($M = 0.65$) was significantly lower than in the placebo group ($M = 0.79$).

Шаг: cmp__тест_прокладывания_пути_trail_making_test_мин_v2_a

Тип: c

Метод: Mann-Whitney U Test
p-value: 0.7871
Статистика: 254.000
Эффект: rbc 0.05




Вывод: An independent Mann-Whitney U Test was conducted to determine if there were differences in «Тест прокладывания пути» Trail Making Test мин V2 A between groups defined by arm. The analysis revealed no statistically significant difference between the groups ($p = 0.787$, $r = 0.05$ indicates a trivial effect).

Шаг: cmp__тест_прокладывания_пути_trail_making_test_мин_v2_б

Тип: c

Метод: Mann-Whitney U Test
p-value: 0.5416
Статистика: 268.500
Эффект: rbc 0.11



Вывод: An independent Mann-Whitney U Test was conducted to determine if there were differences in «Тест прокладывания пути» Trail Making Test мин V2 Б between groups defined by arm. The analysis revealed no statistically significant difference between the groups ($p = 0.542$, $r = 0.11$ indicates a small effect).

Шаг: `cmp__шкала_оценки_качества_жизни_при_бп_баллы_v2`

Тип: c

Метод: Welch's T-Test

p-value: 0.8776


Статистика: -0.155

Эффект: cohen-d 0.05

ДИ эффекта: [-9.61, 8.24]

Мощность: 0.05

BF10: 0.3



Вывод: An independent Welch's T-Test was conducted to determine if there were differences in Шкала оценки качества жизни при БП баллы V2 between groups defined by arm. The analysis revealed no statistically significant difference between the groups ($p = 0.878$, Cohen's $d = 0.05$ indicates a trivial effect).

Шаг: `cmp__ушобп_часть_2_повседневная_активность_баллы_v3`

Тип: c

Метод: Welch's T-Test

p-value: 0.5366

Статистика: 0.624

Эффект: cohen-d 0.19

ДИ эффекта: [-1.33, 2.51]

Мощность: 0.09

BF10: 0.348



Вывод: An independent Welch's T-Test was conducted to determine if there were differences in УШОБП часть 2 «Повседневная активность» баллы V3 between groups defined by arm. The analysis revealed no statistically significant difference between the groups (p = 0.537, Cohen's d = 0.19 indicates a trivial effect).

Шаг: cmp_ушобп_часть_3_двигательные_функции_баллы_v3

Тип: c

Метод: Welch's T-Test

p-value: 0.4230


Статистика: -0.813

Эффект: cohen-d 0.25

ДИ эффекта: [-4.31, 1.86]

Мощность: 0.12

BF10: 0.388



Вывод: An independent Welch's T-Test was conducted to determine if there were differences in УШОБП часть 3 «Двигательные функции» баллы V3 between groups defined by arm. The analysis revealed no statistically significant difference between the groups (p = 0.423, Cohen's d = 0.25 indicates a small effect).

Шаг: cmp_шкала_депрессии_тревоги_и_стресса_dass_21_баллы_v3

Тип: c

Метод: Student's t-test (Independent)

p-value: 0.0025


Статистика: -3.221

Эффект: cohen-d 0.97

ДИ эффекта: [-10.72, -2.46]

Мощность: 0.88

BF10: 14.892



Вывод: An independent Student's t-test (Independent) was conducted to determine if there were differences in Шкала депрессии тревоги и стресса DASS-21 баллы V3 between groups defined by arm. There was a statistically significant difference between the groups ($p = 0.002$, Cohen's $d = 0.97$ indicates a large effect). Specifically, the Шкала депрессии тревоги и стресса DASS-21 баллы V3 in the active group ($M = 27.95$) was significantly lower than in the placebo group ($M = 34.55$).

Шаг: cmp_шкала_оценки_устоляемости_при_бп_баллы_v3


Тип: c

Метод: Mann-Whitney U Test

p-value: 0.3067

Статистика: 198.000

Эффект: rbc -0.18



Вывод: An independent Mann-Whitney U Test was conducted to determine if there were differences in Шкала оценки утомляемости при БП баллы V3 between groups defined by arm. The analysis revealed no statistically significant difference between the groups ($p = 0.307$, $r = -0.18$ indicates a small effect).

Шаг: cmp_шкала_оценки_сонливости_эпуорта_баллы_v3

Тип: c

Метод: Mann-Whitney U Test

p-value: 0.9246

Статистика: 246.500

Эффект: rbc 0.02




Вывод: An independent Mann-Whitney U Test was conducted to determine if there were differences in Шкала оценки сонливости Эпуорта баллы V3 between groups defined by arm. The analysis revealed no statistically significant difference between the groups ($p = 0.925$, $r = 0.02$ indicates a trivial effect).

Шаг: cmp__шкала_апатии_старкстейна_баллы_v3

Тип: c

Метод: Student's t-test (Independent)
p-value: 0.3294
Статистика: -0.987
Эффект: cohen-d 0.30
ДИ эффекта: [-3.32, 1.14]
Мощность: 0.16
BF10: 0.44



Вывод: An independent Student's t-test (Independent) was conducted to determine if there were differences in Шкала апатии Старкстейна баллы V3 between groups defined by arm. The analysis revealed no statistically significant difference between the groups ($p = 0.329$, Cohen's $d = 0.30$ indicates a small effect).

Шаг: cmp__струн_v3_1

Тип: c

Метод: Student's t-test (Independent)
p-value: 0.9964
Статистика: -0.005
Эффект: cohen-d 0.00
ДИ эффекта: [-1.91, 1.90]
Мощность: 0.05
BF10: 0.3



Вывод: An independent Student's t-test (Independent) was conducted to determine if there were differences in Струн V3 1 between groups defined by arm. The analysis revealed no statistically significant difference between the groups ($p = 0.996$, Cohen's $d = 0.00$ indicates a trivial effect).


Шаг: cmp_струн_v3_2

Метод: Mann-Whitney U Test

p-value: 0.0068

Статистика: 342.500

Эффект: rbc 0.48



Вывод: An independent Mann-Whitney U Test was conducted to determine if there were differences in Струн V3 2 between groups defined by arm. There was a statistically significant difference between the groups ($p = 0.007$, $r = 0.48$ indicates a medium effect). Specifically, the Струн V3 2 in the active group ($M = 30.45$) was significantly higher than in the placebo group ($M = 24.38$).


Шаг: cmp_струн_v3_3

Метод: Mann-Whitney U Test

p-value: 0.0116

Статистика: 349.500

Эффект: rbc 0.44




Вывод: An independent Mann-Whitney U Test was conducted to determine if there were differences in Стрпун V3 3 between groups defined by arm. There was a statistically significant difference between the groups ($p = 0.012$, $r = 0.44$ indicates a medium effect). Specifically, the Стрпун V3 3 in the active group ($M = 20.32$) was significantly higher than in the placebo group ($M = 18.05$).

Шаг: cmp_индекс_интерференции_v4

Тип: c

Метод: Student's t-test (Independent)
p-value: 0.2096
Статистика: -1.275
Эффект: cohen-d 0.39
ДИ эффекта: [-0.14, 0.03]
Мощность: 0.24
BF10: 0.574




Вывод: An independent Student's t-test (Independent) was conducted to determine if there were differences in Индекс интерференции V4 between groups defined by arm. The analysis revealed no statistically significant difference between the groups ($p = 0.210$, Cohen's $d = 0.39$ indicates a small effect).

Шаг: cmp_тест_прокладывания_пути_trail_making_test_мин_v3_a

Тип: c

Метод: Mann-Whitney U Test
p-value: 0.2901
Статистика: 196.500
Эффект: rbc -0.19




Вывод: An independent Mann-Whitney U Test was conducted to determine if there were differences in «Тест прокладывания пути» Trail Making Test мин V3 А between groups defined by arm. The analysis revealed no statistically significant difference between the groups ($p = 0.290$, $r = -0.19$ indicates a small effect).

Шаг: cmp_тест_прокладывания_пути_trail_making_test_мин_v3_б

Тип: c

Метод: Mann-Whitney U Test
p-value: 0.1624
Статистика: 302.000
Эффект: rbc 0.25




Вывод: An independent Mann-Whitney U Test was conducted to determine if there were differences in «Тест прокладывания пути» Trail Making Test мин V3 Б between groups defined by arm. The analysis revealed no statistically significant difference between the groups ($p = 0.162$, $r = 0.25$ indicates a small effect).

Шаг: cmp_шкала_оценки_качества_жизни_при_бп_баллы_v3

Тип: c

Метод: Welch's T-Test
p-value: 0.1144
Статистика: -1.621
Эффект: cohen-d 0.49
ДИ эффекта: [-14.24, 1.61]
Мощность: 0.35
BF10: 0.845




Вывод: An independent Welch's T-Test was conducted to determine if there were differences in Шкала оценки качества жизни при БП баллы V3 between groups defined by arm. The analysis revealed no statistically significant difference between the groups ($p = 0.114$, Cohen's $d = 0.49$ indicates a small effect).

Шаг: cmp__сад_v4

Тип: c


Метод: Student's t-test (Independent)
p-value: 0.8353
Статистика: -0.209
Эффект: cohen-d 0.06
ДИ эффекта: [-7.26, 5.89]
Мощность: 0.05
BF10: 0.303



Шаг: cmp__дад_v4

Тип: c

Метод: Mann-Whitney U Test
p-value: 0.1909
Статистика: 298.000
Эффект: rbc 0.23



Вывод: An independent Mann-Whitney U Test was conducted to determine if there were differences in ДАД V4 between groups defined by arm. The analysis revealed no statistically significant difference between the groups ($p = 0.191$, $r = 0.23$ indicates a small effect).

Шаг: cmp_ушобп_часть_2_повседневная_активность_баллы_v4

Тип: c

Метод: Welch's T-Test

p-value: 0.7560


Статистика: 0.313

Эффект: cohen-d 0.09

ДИ эффекта: [-1.50, 2.04]

Мощность: 0.06

BF10: 0.31



Вывод: An independent Welch's T-Test was conducted to determine if there were differences in УШОБП часть 2 «Повседневная активность» баллы V4 between groups defined by arm. The analysis revealed no statistically significant difference between the groups ($p = 0.756$, Cohen's $d = 0.09$ indicates a trivial effect).

Шаг: cmp_ушобп_часть_3_двигательные_функции_баллы_v4

Тип: c

Метод: Welch's T-Test

p-value: 0.1690

Статистика: -1.414

Эффект: cohen-d 0.43

ДИ эффекта: [-4.90, 0.90]

Мощность: 0.28

BF10: 0.661



Вывод: An independent Welch's T-Test was conducted to determine if there were differences in УШОБП часть 3 «Двигательные функции» баллы V4 between groups defined by arm. The analysis revealed no statistically significant difference between the groups (p = 0.169, Cohen's d = 0.43 indicates a small effect).

Шаг: cmp_шкала_депрессии_тревоги_и_стресса_dass_21_баллы_v4

Тип: c

Метод: Welch's T-Test

p-value: < 0.001


Статистика: -3.876

Эффект: cohen-d 1.17

ДИ эффекта: [-12.05, -3.76]

Мощность: 0.97

BF10: 72.108



Вывод: An independent Welch's T-Test was conducted to determine if there were differences in Шкала депрессии тревоги и стресса DASS-21 баллы V4 between groups defined by arm. There was a statistically significant difference between the groups (p < 0.001, Cohen's d = 1.17 indicates a large effect). Specifically, the Шкала депрессии тревоги и стресса DASS-21 баллы V4 in the active group (M = 25.95) was significantly lower than in the placebo group (M = 33.86).

Шаг: cmp_шкала_оценки_устоляемости_при_бп_баллы_v4

Тип: c

Метод: Welch's T-Test

p-value: 0.2386

Статистика: -1.199

Эффект: cohen-d 0.36

ДИ эффекта: [-11.51, 2.96]

Мощность: 0.22

BF10: 0.53




Вывод: An independent Welch's T-Test was conducted to determine if there were differences in Шкала оценки утомляемости при БП баллы V4 between groups defined by arm. The analysis revealed no statistically significant difference between the groups ($p = 0.239$, Cohen's $d = 0.36$ indicates a small effect).

Шаг: cmp_шкала_оценки_сонливости_эпуорта_баллы_v4

Тип: c

Метод: Mann-Whitney U Test
p-value: 0.5223
Статистика: 214.500
Эффект: rbc -0.11




Вывод: An independent Mann-Whitney U Test was conducted to determine if there were differences in Шкала оценки сонливости Эпуорта баллы V4 between groups defined by arm. The analysis revealed no statistically significant difference between the groups ($p = 0.522$, $r = -0.11$ indicates a small effect).

Шаг: cmp_шкала_апатии_старкстейна_баллы_v4

Тип: c

Метод: Mann-Whitney U Test
p-value: 0.2322
Статистика: 191.000
Эффект: rbc -0.21




Вывод: An independent Mann-Whitney U Test was conducted to determine if there were differences in Шкала апатии Старкстейна баллы V4 between groups defined by arm. The analysis revealed no statistically significant difference between the groups ($p = 0.232$, $r = -0.21$ indicates a small effect).

Шаг: cmp_тест_прокладывания_пути_trail_making_test_мин_v4_a

Тип: c

Метод: Mann-Whitney U Test
p-value: 0.0325
Статистика: 150.500
Эффект: rbc -0.38




Вывод: An independent Mann-Whitney U Test was conducted to determine if there were differences in «Тест прокладывания пути» Trail Making Test мин V4 A between groups defined by arm. There was a statistically significant difference between the groups ($p = 0.032$, $r = -0.38$ indicates a medium effect). Specifically, the «Тест прокладывания пути» Trail Making Test мин V4 A in the active group ($M = 37.91$) was significantly lower than in the placebo group ($M = 49.45$).

Шаг: cmp_тест_прокладывания_пути_trail_making_test_мин_v4_б

Тип: c

Метод: Mann-Whitney U Test
p-value: 0.8880
Статистика: 235.500
Эффект: rbc -0.03



Вывод: An independent Mann-Whitney U Test was conducted to determine if there were differences in «Тест прокладывания пути» Trail Making Test мин V4 Б between groups defined by arm. The analysis revealed no statistically significant difference between the groups ($p = 0.888$, $r = -0.03$ indicates a trivial effect).

Шаг: `cmp__шкала_оценки_качества_жизни_при_бп_баллы`

Тип: c

Метод: Welch's T-Test

p-value: 0.1395


Статистика: -1.516

Эффект: cohen-d 0.46

ДИ эффекта: [-13.00, 1.91]

Мощность: 0.32

BF10: 0.744



Шаг: `cmp__ушобп_часть_2_повседневная_активность_баллы_v5`

Тип: c

Метод: Welch's T-Test

p-value: 0.7882

Статистика: -0.271

Эффект: cohen-d 0.08

ДИ эффекта: [-1.94, 1.49]

Мощность: 0.06

BF10: 0.306



Вывод: An independent Welch's T-Test was conducted to determine if there were differences in УШОБП часть 2 «Повседневная активность» баллы V5 between groups defined by arm. The analysis revealed no statistically significant difference between the groups (p = 0.788, Cohen's d = 0.08 indicates a trivial effect).

Шаг: cmp_ушобп_часть_3_двигательные_функции_баллы_v5

Тип: c

Метод: Welch's T-Test

p-value: 0.0296


Статистика: -2.309

Эффект: cohen-d 0.70

ДИ эффекта: [-6.11, -0.35]

Мощность: 0.62

BF10: 2.386



Вывод: An independent Welch's T-Test was conducted to determine if there were differences in УШОБП часть 3 «Двигательные функции» баллы V5 between groups defined by arm. There was a statistically significant difference between the groups (p = 0.030, Cohen's d = 0.70 indicates a medium effect). Specifically, the УШОБП часть 3 «Двигательные функции» баллы V5 in the active group (M = 8.45) was significantly lower than in the placebo group (M = 11.68).

Шаг: cmp_шкала_депрессии_тревоги_и_стресса_dass_21_баллы_v5


Тип: c

Метод: Mann-Whitney U Test

p-value: < 0.001

Статистика: 81.500

Эффект: rbc -0.66



Вывод: An independent Mann-Whitney U Test was conducted to determine if there were differences in Шкала депрессии тревоги и стресса DASS-21 баллы V5 between groups defined by arm. There was a statistically significant difference between the groups ($p < 0.001$, $r = -0.66$ indicates a large effect). Specifically, the Шкала депрессии тревоги и стресса DASS-21 баллы V5 in the active group ($M = 24.50$) was significantly lower than in the placebo group ($M = 33.32$).

Шаг: cmp_шкала_оценки_устоляемости_при_бп_баллы_v5


Тип: C

Метод: Mann-Whitney U Test

p-value: 0.2002

Статистика: 187.000

Эффект: rbc -0.23



Вывод: An independent Mann-Whitney U Test was conducted to determine if there were differences in Шкала оценки утомляемости при БП баллы V5 between groups defined by arm. The analysis revealed no statistically significant difference between the groups ($p = 0.200$, $r = -0.23$ indicates a small effect).

Шаг: cmp_шкала_оценки_сонливости_эпуорта_баллы_v5


Тип: C

Метод: Mann-Whitney U Test

p-value: 0.4322

Статистика: 208.500

Эффект: rbc -0.14




Вывод: An independent Mann-Whitney U Test was conducted to determine if there were differences in Шкала оценки сонливости Эпуорта баллы V5 between groups defined by arm. The analysis revealed no statistically significant difference between the groups ($p = 0.432$, $r = -0.14$ indicates a small effect).

Шаг: cmp_шкала_апатии_старкстейна_баллы_v5

Тип: c

Метод: Mann-Whitney U Test
p-value: 0.0949
Статистика: 171.000
Эффект: rbc -0.29



Вывод: An independent Mann-Whitney U Test was conducted to determine if there were differences in Шкала апатии Старкстейна баллы V5 between groups defined by arm. The analysis revealed no statistically significant difference between the groups ($p = 0.095$, $r = -0.29$ indicates a small effect).

Шаг: cmp_струп_v5_1

Тип: c

Метод: Student's t-test (Independent)
p-value: 0.0044
Статистика: -3.008
Эффект: cohen-d 0.91
ДИ эффекта: [-2.96, -0.58]
Мощность: 0.84
BF10: 9.322




Вывод: An independent Student's t-test (Independent) was conducted to determine if there were differences in Струн V5 1 between groups defined by arm. There was a statistically significant difference between the groups ($p = 0.004$, Cohen's $d = 0.91$ indicates a large effect). Specifically, the Струн V5 1 in the active group ($M = 14.14$) was significantly lower than in the placebo group ($M = 15.91$).

Шаг: cmp_струн_v5_2

Тип: c

Метод: Student's t-test (Independent)
p-value: 0.0205
Статистика: 2.408
Эффект: cohen-d 0.73
ДИ эффекта: [0.38, 4.26]
Мощность: 0.65
BF10: 2.842




Вывод: An independent Student's t-test (Independent) was conducted to determine if there were differences in Струн V5 2 between groups defined by arm. There was a statistically significant difference between the groups ($p = 0.020$, Cohen's $d = 0.73$ indicates a medium effect). Specifically, the Струн V5 2 in the active group ($M = 23.95$) was significantly higher than in the placebo group ($M = 21.64$).

Шаг: cmp_струн_v5_3

Тип: c

Метод: Student's t-test (Independent)
p-value: 0.2717
Статистика: 1.114
Эффект: cohen-d 0.34
ДИ эффекта: [-0.63, 2.17]
Мощность: 0.19
BF10: 0.489




Вывод: An independent Student's t-test (Independent) was conducted to determine if there were differences in Струп V5 3 between groups defined by arm. The analysis revealed no statistically significant difference between the groups ($p = 0.272$, Cohen's $d = 0.34$ indicates a small effect).

Шаг: cmp_индекс_интерференции_v5

Тип: c

Метод: Student's t-test (Independent)
p-value: 0.1934
Статистика: -1.322
Эффект: cohen-d 0.40
ДИ эффекта: [-0.14, 0.03]
Мощность: 0.25
BF10: 0.598




Вывод: An independent Student's t-test (Independent) was conducted to determine if there were differences in Индекс интерференции V5 between groups defined by arm. The analysis revealed no statistically significant difference between the groups ($p = 0.193$, Cohen's $d = 0.40$ indicates a small effect).

Шаг: cmp_тест_прокладывания_пути_trail_making_test_мин_v5_a

Тип: c

Метод: Mann-Whitney U Test
p-value: 0.0021
Статистика: 110.500
Эффект: rbc -0.54




Вывод: An independent Mann-Whitney U Test was conducted to determine if there were differences in «Тест прокладывания пути» Trail Making Test мин V5 A between groups defined by arm. There was a statistically significant difference between the groups ($p = 0.002$, $r = -0.54$ indicates a large effect). Specifically, the «Тест прокладывания пути» Trail Making Test мин V5 A in the active group ($M = 36.18$) was significantly lower than in the placebo group ($M = 49.50$).

Шаг: cmp_тест_прокладывания_пути_trail_making_test_мин_v5_б

Тип: c

Метод: Mann-Whitney U Test
p-value: 0.7512
Статистика: 228.000
Эффект: rbc -0.06




Вывод: An independent Mann-Whitney U Test was conducted to determine if there were differences in «Тест прокладывания пути» Trail Making Test мин V5 Б between groups defined by arm. The analysis revealed no statistically significant difference between the groups ($p = 0.751$, $r = -0.06$ indicates a trivial effect).

Шаг: cmp_шкала_оценки_качества_жизни_при_бп_баллы_v5

Тип: c

Метод: Welch's T-Test
p-value: 0.0300
Статистика: -2.277
Эффект: cohen-d 0.69
ДИ эффекта: [-14.66, -0.80]
Мощность: 0.60
BF10: 2.256




Вывод: An independent Welch's T-Test was conducted to determine if there were differences in Шкала оценки качества жизни при БП баллы V5 between groups defined by arm. There was a statistically significant difference between the groups ($p = 0.030$, Cohen's $d = 0.69$ indicates a medium effect). Specifically, the Шкала оценки качества жизни при БП баллы V5 in the active group ($M = 52.64$) was significantly lower than in the placebo group ($M = 60.36$).

Шаг: cmp__сад_v6

Тип: С

Метод: Student's t-test (Independent)
p-value: 0.6646
Статистика: -0.437
Эффект: cohen-d 0.13
ДИ эффекта: [-7.41, 4.77]
Мощность: 0.07
BF10: 0.321



Шаг: cmp__дад_v6

Тип: С

Метод: Student's t-test (Independent)
p-value: 0.0065
Статистика: 2.862
Эффект: cohen-d 0.86
ДИ эффекта: [1.93, 11.16]
Мощность: 0.80
BF10: 6.843



Вывод: An independent Student's t-test (Independent) was conducted to determine if there were differences in ДАД V6 between groups defined by arm. There was a statistically significant difference between the groups ($p = 0.007$, Cohen's $d = 0.86$ indicates a large effect). Specifically, the ДАД V6 in the active group ($M = 84.27$) was significantly higher than in the placebo group ($M = 77.73$).

Шаг: cmp_ушобп_часть_2_повседневная_активность_баллы_v6

Тип: C

Метод: Welch's T-Test

p-value: 0.7010


Статистика: -0.388

Эффект: cohen-d 0.12

ДИ эффекта: [-2.00, 1.36]

Мощность: 0.07

BF10: 0.316



Вывод: An independent Welch's T-Test was conducted to determine if there were differences in УШОБП часть 2 «Повседневная активность» баллы V6 between groups defined by arm. The analysis revealed no statistically significant difference between the groups ($p = 0.701$, Cohen's $d = 0.12$ indicates a trivial effect).

Шаг: cmp_ушобп_часть_3_двигательные_функции_баллы_v6


Тип: C

Метод: Mann-Whitney U Test

p-value: 0.0605

Статистика: 162.000

Эффект: rbc -0.33




Вывод: An independent Mann-Whitney U Test was conducted to determine if there were differences in УШОБП часть 3 «Двигательные функции» баллы V6 between groups defined by arm. The analysis revealed no statistically significant difference between the groups ($p = 0.061$, $r = -0.33$ indicates a medium effect).

Шаг: cmp_шкала_депрессии_тревоги_и_стресса_dass_21баллы_v6

Тип: c

Метод: Mann-Whitney U Test
p-value: < 0.001
Статистика: 66.500
Эффект: rbc -0.73




Вывод: An independent Mann-Whitney U Test was conducted to determine if there were differences in Шкала депрессии, тревоги и стресса DASS-21баллы V6 between groups defined by arm. There was a statistically significant difference between the groups ($p < 0.001$, $r = -0.73$ indicates a large effect). Specifically, the Шкала депрессии, тревоги и стресса DASS-21баллы V6 in the active group ($M = 24.00$) was significantly lower than in the placebo group ($M = 32.73$).

Шаг: cmp_шкала_оценки_устоляемости_при_бп_баллы_v6

Тип: c

Метод: Mann-Whitney U Test
p-value: 0.1922
Статистика: 186.000
Эффект: rbc -0.23




Вывод: An independent Mann-Whitney U Test was conducted to determine if there were differences in Шкала оценки утомляемости при БП баллы V6 between groups defined by arm. The analysis revealed no statistically significant difference between the groups ($p = 0.192$, $r = -0.23$ indicates a small effect).

Шаг: cmp_шкала_оценки_сонливости_эпуорта_баллы_v6

Тип: c

Метод: Mann-Whitney U Test
p-value: 0.4983
Статистика: 213.000
Эффект: rbc -0.12




Вывод: An independent Mann-Whitney U Test was conducted to determine if there were differences in Шкала оценки сонливости Эпуорта баллы V6 between groups defined by arm. The analysis revealed no statistically significant difference between the groups ($p = 0.498$, $r = -0.12$ indicates a small effect).

Шаг: cmp_шкала_апатии_старкстейна_баллы_v6

Тип: c

Метод: Mann-Whitney U Test
p-value: 0.0298
Статистика: 150.000
Эффект: rbc -0.38




Вывод: An independent Mann-Whitney U Test was conducted to determine if there were differences in Шкала апатии Старкстейна баллы V6 between groups defined by arm. There was a statistically significant difference between the groups ($p = 0.030$, $r = -0.38$ indicates a medium effect). Specifically, the Шкала апатии Старкстейна баллы V6 in the active group ($M = 4.82$) was significantly lower than in the placebo group ($M = 7.64$).

Шаг: cmp_струп_v6_1

Тип: c

Метод: Mann-Whitney U Test
p-value: < 0.001
Статистика: 98.500
Эффект: rbc -0.59




Вывод: An independent Mann-Whitney U Test was conducted to determine if there were differences in Струп V6 1 between groups defined by arm. There was a statistically significant difference between the groups ($p < 0.001$, $r = -0.59$ indicates a large effect). Specifically, the Струп V6 1 in the active group ($M = 13.91$) was significantly lower than in the placebo group ($M = 16.14$).

Шаг: cmp_струп_v6_2

Тип: c

Метод: Mann-Whitney U Test
p-value: 0.0299
Статистика: 334.500
Эффект: rbc 0.38




Вывод: An independent Mann-Whitney U Test was conducted to determine if there were differences in Струн V6 2 between groups defined by arm. There was a statistically significant difference between the groups ($p = 0.030$, $r = 0.38$ indicates a medium effect). Specifically, the Струн V6 2 in the active group ($M = 22.77$) was significantly higher than in the placebo group ($M = 21.27$).

Шаг: cmp_струн_v6_3

Тип: С

Метод: Mann-Whitney U Test
p-value: 0.4882
Статистика: 271.500
Эффект: rbc 0.12



Вывод: An independent Mann-Whitney U Test was conducted to determine if there were differences in Струн V6 3 between groups defined by arm. The analysis revealed no statistically significant difference between the groups ($p = 0.488$, $r = 0.12$ indicates a small effect).

Шаг: cmp_индекс_интерференции_v6

Тип: С

Метод: Student's t-test (Independent)
p-value: 0.1819
Статистика: -1.357
Эффект: cohen-d 0.41
ДИ эффекта: [-0.13, 0.02]
Мощность: 0.26
BF10: 0.621



Вывод: An independent Student's t-test (Independent) was conducted to determine if there were differences in Индекс интерференции V6 between groups defined by arm. The analysis revealed no statistically significant difference between the groups ($p = 0.182$, Cohen's $d = 0.41$ indicates a small effect).

Шаг: cmp_тест_прокладывания_пути_trail_making_test_мин_v6_a


Тип: c

Метод: Mann-Whitney U Test

p-value: < 0.001

Статистика: 92.000

Эффект: rbc -0.62



Вывод: An independent Mann-Whitney U Test was conducted to determine if there were differences in «Тест прокладывания пути» Trail Making Test мин V6 A between groups defined by arm. There was a statistically significant difference between the groups ($p < 0.001$, $r = -0.62$ indicates a large effect). Specifically, the «Тест прокладывания пути» Trail Making Test мин V6 A in the active group ($M = 35.64$) was significantly lower than in the placebo group ($M = 49.64$).

Шаг: cmp_тест_прокладывания_пути_trail_making_test_мин_v6_б


Тип: c

Метод: Mann-Whitney U Test

p-value: 0.2311

Статистика: 190.500

Эффект: rbc -0.21



Вывод: An independent Mann-Whitney U Test was conducted to determine if there were differences in «Тест прокладывания пути» Trail Making Test мин V6 Б between groups defined by arm. The analysis revealed no statistically significant difference between the groups ($p = 0.231$, $r = -0.21$ indicates a small effect).

Шаг: cmp__шкала_оценки_качества_жизни_при_бп_баллы_v6

Тип: c

Метод: Welch's T-Test

p-value: 0.0038


Статистика: -3.146

Эффект: cohen-d 0.95

ДИ эффекта: [-16.28, -3.45]

Мощность: 0.87

BF10: 12.582



Вывод: An independent Welch's T-Test was conducted to determine if there were differences in Шкала оценки качества жизни при БП баллы V6 between groups defined by arm. There was a statistically significant difference between the groups ($p = 0.004$, Cohen's $d = 0.95$ indicates a large effect). Specifically, the Шкала оценки качества жизни при БП баллы V6 in the active group ($M = 50.95$) was significantly lower than in the placebo group ($M = 60.82$).

Шаг: cmp__stroop_sum_v2_sec

Тип: c

Метод: Student's t-test (Independent)

p-value: 0.0522

Статистика: 2.003

Эффект: cohen-d 0.63

ДИ эффекта: [-0.14, 28.12]

Мощность: 0.50

BF10: 1.458



Вывод: An independent Student's t-test (Independent) was conducted to determine if there were differences in Stroop sum V2 (sec) between groups defined by arm. The analysis revealed no statistically significant difference between the groups ($p = 0.052$, Cohen's $d = 0.63$ indicates a medium effect).

Шаг: cmp_stroop_sum_v3_sec


Тип: c

Метод: Mann-Whitney U Test

p-value: 0.0076

Статистика: 356.000

Эффект: rbc 0.47



Вывод: An independent Mann-Whitney U Test was conducted to determine if there were differences in Stroop sum V3 (sec) between groups defined by arm. There was a statistically significant difference between the groups ($p = 0.008$, $r = 0.47$ indicates a medium effect). Specifically, the Stroop sum V3 (sec) in the active group ($M = 66.86$) was significantly higher than in the placebo group ($M = 56.68$).

Шаг: cmp_stroop_sum_v5_sec

Тип: c

Метод: Student's t-test (Independent)

p-value: 0.4061

Статистика: 0.839

Эффект: cohen-d 0.25

ДИ эффекта: [-1.85, 4.49]

Мощность: 0.13

BF10: 0.395




Вывод: An independent Student's t-test (Independent) was conducted to determine if there were differences in Stroop sum V5 (sec) between groups defined by arm. The analysis revealed no statistically significant difference between the groups ($p = 0.406$, Cohen's $d = 0.25$ indicates a small effect).

Шаг: cmp__stroop_sum_v6_sec

Тип: c

Метод: Student's t-test (Independent)
p-value: 0.8709
Статистика: -0.163
Эффект: cohen-d 0.05
ДИ эффекта: [-3.64, 3.09]
Мощность: 0.05
BF10: 0.301




Вывод: An independent Student's t-test (Independent) was conducted to determine if there were differences in Stroop sum V6 (sec) between groups defined by arm. The analysis revealed no statistically significant difference between the groups ($p = 0.871$, Cohen's $d = 0.05$ indicates a trivial effect).

Шаг: cmp__trail_a_b_v2_sec

Тип: c

Метод: Mann-Whitney U Test
p-value: 0.5186
Статистика: 270.000
Эффект: rbc 0.12




Вывод: An independent Mann-Whitney U Test was conducted to determine if there were differences in Trail A+B V2 (sec) between groups defined by arm. The analysis revealed no statistically significant difference between the groups ($p = 0.519$, $r = 0.12$ indicates a small effect).

Шаг: cmp__trail_a_b_v6_sec

Тип: c

Метод: Student's t-test (Independent)
p-value: 0.0317
Статистика: -2.222
Эффект: cohen-d 0.67
ДИ эффекта: [-40.68, -1.96]
Мощность: 0.58
BF10: 2.056




Вывод: An independent Student's t-test (Independent) was conducted to determine if there were differences in Trail A+B V6 (sec) between groups defined by arm. There was a statistically significant difference between the groups ($p = 0.032$, Cohen's $d = 0.67$ indicates a medium effect). Specifically, the Trail A+B V6 (sec) in the active group ($M = 125.86$) was significantly lower than in the placebo group ($M = 147.18$).

Шаг: cmp__ушобп_часть_3_двигательные_функции_баллы_v2_v6_улучшение

Тип: c

Метод: Mann-Whitney U Test
p-value: < 0.001
Статистика: 433.000
Эффект: rbc 0.79




Вывод: An independent Mann-Whitney U Test was conducted to determine if there were differences in Δ УШОБП часть 3 «Двигательные функции» баллы (V2→V6, улучшение) between groups defined by arm. There was a statistically significant difference between the groups ($p < 0.001$, $r = 0.79$ indicates a large effect). Specifically, the Δ УШОБП часть 3 «Двигательные функции» баллы (V2→V6, улучшение) in the active group ($M = 3.77$) was significantly higher than in the placebo group ($M = 0.32$).

Шаг: cmp_ушобп_часть_2_повседневная_активность_баллы_v2_v6_улучшение

Тип: c

Метод: Mann-Whitney U Test
p-value: < 0.001
Статистика: 389.500
Эффект: rbc 0.61




Вывод: An independent Mann-Whitney U Test was conducted to determine if there were differences in Δ УШОБП часть 2 «Повседневная активность» баллы (V2→V6, улучшение) between groups defined by arm. There was a statistically significant difference between the groups ($p < 0.001$, $r = 0.61$ indicates a large effect). Specifically, the Δ УШОБП часть 2 «Повседневная активность» баллы (V2→V6, улучшение) in the active group ($M = 2.14$) was significantly higher than in the placebo group ($M = 0.41$).

Шаг: cmp_шкала_оценки_сонливости_эпуорта_баллы_v2_v6_улучшение

Тип: c

Метод: Mann-Whitney U Test
p-value: 0.3013
Статистика: 286.000
Эффект: rbc 0.18



Вывод: An independent Mann-Whitney U Test was conducted to determine if there were differences in Δ Шкала оценки сонливости Эпурта баллы (V2→V6, улучшение) between groups defined by arm. The analysis revealed no statistically significant difference between the groups ($p = 0.301$, $r = 0.18$ indicates a small effect).

Шаг: `cmp_шкала_оценки_устоляемости_при_бп_баллы_v2_v6_улучшение`


Тип: c

Метод: Mann-Whitney U Test

p-value: 0.1953

Статистика: 297.500

Эффект: rbc 0.23



Вывод: An independent Mann-Whitney U Test was conducted to determine if there were differences in Δ Шкала оценки устольяемости при БП баллы (V2→V6, улучшение) between groups defined by arm. The analysis revealed no statistically significant difference between the groups ($p = 0.195$, $r = 0.23$ indicates a small effect).

Шаг: `cmp_шкала_апатии_старкстейна_баллы_v2_v6_улучшение`


Тип: c

Метод: Mann-Whitney U Test

p-value: < 0.001

Статистика: 446.000

Эффект: rbc 0.84




Вывод: An independent Mann-Whitney U Test was conducted to determine if there were differences in Δ Шкала апатии Старкстейна баллы (V2→V6, улучшение) between groups defined by arm. There was a statistically significant difference between the groups ($p < 0.001$, $r = 0.84$ indicates a large effect). Specifically, the Δ Шкала апатии Старкстейна баллы (V2→V6, улучшение) in the active group ($M = 5.00$) was significantly higher than in the placebo group ($M = 0.09$).

Шаг: cmp__шкала_оценки_качества_жизни_при_бп_баллы_v2_v6_улучшение

Тип: c

Метод: Mann-Whitney U Test
p-value: < 0.001
Статистика: 416.500
Эффект: rbc 0.72



Вывод: An independent Mann-Whitney U Test was conducted to determine if there were differences in Δ Шкала оценки качества жизни при БП баллы (V2→V6, улучшение) between groups defined by arm. There was a statistically significant difference between the groups ($p < 0.001$, $r = 0.72$ indicates a large effect). Specifically, the Δ Шкала оценки качества жизни при БП баллы (V2→V6, улучшение) in the active group ($M = 11.27$) was significantly higher than in the placebo group ($M = 2.09$).

Шаг: cmp__stroop_sum_v2_v6_улучшение

Тип: c

Метод: Student's t-test (Independent)
p-value: 0.0374
Статистика: 2.155
Эффект: cohen-d 0.67
ДИ эффекта: [0.85, 26.82]
Мощность: 0.56
BF10: 1.849



Вывод: An independent Student's t-test (Independent) was conducted to determine if there were differences in Δ Stroop sum (V2→V6, улучшение) between groups defined by arm. There was a statistically significant difference between the groups ($p = 0.037$, Cohen's $d = 0.67$ indicates a medium effect). Specifically, the Δ Stroop sum (V2→V6, улучшение) in the active group ($M = 25.79$) was significantly higher than in the placebo group ($M = 11.95$).

Шаг: cmp__trail_a_b_v2_v6_улучшение


Тип: c

Метод: Mann-Whitney U Test

p-value: 0.0542

Статистика: 324.500

Эффект: rbc 0.34



Вывод: An independent Mann-Whitney U Test was conducted to determine if there were differences in Δ Trail A+B (V2→V6, улучшение) between groups defined by arm. The analysis revealed no statistically significant difference between the groups ($p = 0.054$, $r = 0.34$ indicates a medium effect).

Шаг: cmp__сад_v6_v1

Тип: c

Метод: Student's t-test (Independent)

p-value: 0.2159

Статистика: -1.256

Эффект: cohen-d 0.38

ДИ эффекта: [-14.81, 3.44]

Мощность: 0.23

BF10: 0.56



Вывод: An independent Student's t-test (Independent) was conducted to determine if there were differences in Δ САД (V6–V1) between groups defined by arm. The analysis revealed no statistically significant difference between the groups ($p = 0.216$, Cohen's $d = 0.38$ indicates a small effect).

Шаг: cmp_дад_v6_v1

Тип: c

Метод: Student's t-test (Independent)
p-value: 0.1685
Статистика: 1.401
Эффект: cohen-d 0.42
ДИ эффекта: [-1.62, 8.98]
Мощность: 0.28
BF10: 0.652

