

## 1. THE $\mathbb{Q}G$ RIGID SINGULARITIES

We wish to study groups  $G$  that are finite subgroups  $\mathrm{GL}_2(\mathbb{C})$  with the property  $G \cap \mathrm{SL}_2(\mathbb{C}) = \mathrm{BD}_{4n}$ . Consider the three elements

$$a = \begin{pmatrix} \zeta_m & 0 \\ 0 & \zeta_m \end{pmatrix}, \quad b = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad c_m = \begin{pmatrix} \sigma_{2n} & 0 \\ 0 & \sigma_{2n}^{-1} \end{pmatrix}$$

Here  $\zeta$  We split this into two cases the case where  $m$  is odd or even. In the odd case  $G = \langle a, b, c_m \rangle$ , otherwise it is  $\langle c, b \circ a_{2m} \rangle$ . Similar analysis can be done for the groups  $E_7$  and  $E_8$ . We now discuss when their  $\mathbb{Q}g$  smoothings.

### Case 1

Here  $m$  is odd so  $G = \langle a, b, c \rangle$ . Looking at the index one cover we have a  $x^2 + y^2z + z^{n+1}$ . With  $x = (u^{2n} - v^{2n})uv$ ,  $y = u^{2n} + v^{2n}$ ,  $z = u^2v^2$ . Looking at the derivatives we see that  $\mathcal{T}^1$  is generated by  $1, y, z, \dots, z^{n-1}$ . As this is a rational surface singularity we see that  $\mathcal{T}^2 = 0$ , so these deformations are unobstructed. We see that  $c$  acts on  $x, y, z$  with weights  $(2n+2, 2n, 4)$ . This correspond to  $\frac{1}{m}(n+1, n, 2)$  action. Our equation has weight  $2n+2$ . To check rigidity we first need  $n \not\equiv_m -2$ . In this case we have  $y$  cannot be in the  $\mathbb{Q}G$  smoothing. Other than that we just need  $z^i$  cannot have weight  $2n+2$ . This corresponds to  $2n+2 \not\equiv_m 2i$  for  $i$  in  $0 \dots n-1$ . These will be the only  $\mathbb{Q}G$  rigid singularities. In particular this means that if  $m > 2n+2$  then it is rigid.

### Case 2

Here  $m = 2m'$  is even so  $\langle a, b \circ c_{2m} \rangle$ . So we have the same equation as above and we now get a  $\frac{1}{2m'}(2n+m', 2n, 4)$ , so to be rigid once again we need the  $y$  term not to be in the smoothing  $2n+4 \not\equiv_{2m'} 0$ . Once again we need no  $z^i$  terms this means  $4n+4 \not\equiv_{2m'} 4i$  for  $i$  in  $0 \dots n-1$ .

It is easy to see that if a partial smoothing exists then it either stay non cyclic quotient or, if you can deform equivariantly by  $y$ , it becomes a cyclic quotient singularity which is of the form  $\frac{1}{m}(n+1, 2)$  or  $\frac{1}{2m'}(2n_m, 4)$ . From this it is easy to classify the  $\mathbb{Q}g$  smoothable singularities. We note for a given  $n$  there exists only a finite amount of  $\mathbb{Q}g$  smoothable singularities with the  $D_n$  singularity as its index one cover. This is in contrast to the  $A_n$  case.

## 2. THE $E_i$ SINGULARITY

We start with the  $E_6$  singularity as this behaves differently more similarly to the  $D_n$  singularities. Once again we aim to classify the  $\mathbb{Q}G$  rigid singularities. We split it into two cases again.

### Case 1

This is a  $\mathbb{Z}_n$  action with  $n$  coprime to 6. This will act with weights  $(6, 4, 3)$  on the equation  $x^2 + y^3 + z^4$ . This lies in the eigenspace of degree 12. Once again by computing partial derivatives we get that the smoothing parameters are  $1, y, z, z^2$ . We wish for none of these two be invariant. This means that  $12 \not\equiv_n 0, 4, 3, 6$ . This implies that  $n \neq 2, 3, 4, 9, 12$ .

However none of these cases can occur via the coprimality condition stated at the beginning.

### Case 2

This is a  $\mathbb{Z}_n$  action with  $(n, 6) = 3$ . Writing  $n = 3n'$  we have the action with weights  $(6 + n', 4, 3 + n')$ . Once again this is non rigid if  $n \neq 2, 3, 4, 9, 12$ . Via the coprimality condition we get this is non rigid if  $n = 3$  or  $n = 9$ .

The  $E_7$  and  $E_8$  cases are not divide by case. The  $E_7$  has a single possibility of  $\mathbb{Z}_n$  with  $(n, 6) = 1$ . This acts on  $x^2 + y^3 + yz^3$  with the weights  $(9, 6, 4)$ . The partials give us a basis for the deformations of  $1, y, y^2, z, z^2, yz$ . This means that  $18 \not\equiv_n 0, 6, 12, 4, 8, 10$ . All of these would correspond to  $n$  being even so there are no non rigid singularities.

### 3. SINGULARITY CONTENT

We wish to generalise the notion of singularity content from cyclic quotients. We have that given a non cyclic quotient there exists a unique residual singularity. We wish to have other invariants indicating what it has smoothed from. The natural thing is to consider the invariant  $h_{\text{top}}^2(X^0)$  where  $X^0$  is  $X - \{\text{singular locus}\}$ . First considering the case where there is no smoothing in terms of  $y$  so the only terms we can put in are of the form  $z^i$  giving  $x^2 + y^2z + z^j \prod_{k=1}^p (z^i - a_k)$  then it is clear that we have exactly by projecting  $(x, y, z) \mapsto z$  we get a map to  $\mathbb{C}$  with degenerate fibers of 2 curves transversely intersecting over the finitely many points corresponding to  $\lfloor \frac{n+1}{p} \rfloor$ . **Ignoring the central fiber over the origin.** We can apply Mayer-Vietoris repeatedly to get that  $h_{\text{top}}^2(X^0)$  is a deformation invariant. In the case where we have  $y$  in the deformation we have