

A network modelling approach to assess non-pharmaceutical disease controls against SARS-CoV-2 in a worker population

Edward Hill, Ben Atkins, Matt Keeling, Louise Dyson, Mike Tildesley.

Zeeman Institute: SBIDER, School of Life Sciences & Mathematics Institute, University of Warwick, UK.
Joint UNiversities Pandemic and Epidemiological Research, <https://maths.org/juniper/>.

Demonstrates the utility of **network model frameworks** to capture **heterogeneity of demographic attributes** across worker roles and the **individual nature** of non-pharmaceutical interventions.



Click the QR code to download the paper



Edward.Hill@warwick.ac.uk
@EdMHill
<https://edmhill.github.io>

1. Motivation & aims

Globally, many countries have employed **social distancing measures** and **non-pharmaceutical interventions (NPIs)** to curb the spread of SARS-CoV-2 [1]. As part of a collective effort to protect public health by disrupting viral transmission, businesses also need to act appropriately by taking all reasonable measures to **minimise exposure** to coronavirus in workplaces and premises open to the public [2]. Adjustments in working practices can result in changes to the amount, duration, and/or proximity of interactions, thereby **altering the dynamics of viral spread**.

Study objectives:

- Parameterise an **individual-based** network model of workers, stratified into work sectors, using a **data-driven** approach;
- Study **epidemic spread of SARS-CoV-2** amongst a population of workers and analyse the impact of **NPIs** targeted towards working practices.


2. Network model description

- Network model construction with **nodes representing workers** and **connections representing contacts** that can result in disease transmission in distinct settings: workplace, household and social (Fig. 1).
- Ran a SARS-CoV-2 outbreak upon the network, with **transmission scaled** according to setting and symptomatic status.

Simulation overview:

- Population & time horizon:** 10,000 workers, 365 days.
- Simulation count:** 1000 runs
 - 50 distinct network realisations
 - 20 runs per network realisation
- Initial disease state conditions:**
 - Ten individuals in an infectious state.
 - All other individuals began in a susceptible state.
- Default worker pattern:** Assumed all workers had the same working pattern of five days at the workplace (Monday-Friday) and two days off (Saturday and Sunday).
- Intervention implementation:** Assumed that all NPIs implemented from day 15, including isolation and test-and-trace (default assumption of 70% adherent).


Fig. 1: Illustration of the structured layers of the worker network model.



3. Contact parameterisation


- Contact distributions and risk parameterised by the Warwick contact survey [3-5].
- We generated static contacts using a "configuration model" style algorithm, specifying a desired degree distribution for each of 41 work sectors.
- Lognormal distributions consistently provided stronger correspondence to the data, across different occupations, than alternative distribution choices (Fig. 2).

Fig. 2: Parametric distribution fits to empirical data on number of: (a) work contacts (office worker industry sector as an example), (b) social contacts.




4. Results: Workplace targeted interventions


- A large proportion of the workforce working from home **stunts outbreaks** (Fig. 3).



- Asynchronous work patterns** reduces infections compared with scenarios where all workers work on the same days, particularly for longer working weeks (Fig. 4).




- Smaller work teams** and a greater reduction in transmission risk lessened the probability of large, prolonged outbreaks (Fig. 5).



5. Results: Adherence to test, trace & isolation

- Absence of sufficient adherence** to non-pharmaceutical interventions **increases** the chance of SARS-CoV-2 spreading widely in the population (Fig. 6).

Fig. 6: Case and isolation summary statistics under differing levels of adherence to test, trace and isolate measures.



6. Possible model developments

- Augment model with **age structure**.
- Allow for **clustering of individuals** within individual workplaces.
- Inclusion of **part time workers**.
- Explore **sensitivity** to alternative epidemiological and intervention assumptions, e.g. presence of other respiratory infections and impact on test capacity when levels of cough and fever are high due to non-COVID-19 causes.
- Application to **other countries**, given availability of associated data to parameterise the model framework.

Acknowledgements

The Social Contact Survey data are available from <https://wrap.warwick.ac.uk/54273/>. The code repository for the study is available at: https://github.com/EdMHill/covid19_worker_network_model. EH, BA, MK, LD and MT were supported by the Medical Research Council through the COVID-19 Rapid Response Rolling Call [grant number MR/V009761/1]; MK, LD and MT were supported by the Engineering and Physical Sciences Research Council through the MathSys CDT [grant number EP/S022244/1] and by UKRI through the JUNIPER modelling consortium [grant number MR/V038613/1]. The funders had no role in the study design, data collection or analysis.

References

- [1] Hale T, Angrist N, Cameron-Blake E, Hallas L, Kira B, Majumdar S, et al. Oxford COVID-19 Government Response Tracker, Blavatnik School of Government.; 2020. URL: www.bsg.ox.ac.uk/covidtracker.
- [2] UK Government. Working safely during coronavirus (COVID-19); 2020. URL: <https://bit.ly/38hCjwG>
- [3] Danon L, Read JM, Keeling MJ, House TA, Vernon MC. Social contact study [Dataset]; 2009. Available from: <https://wrap.warwick.ac.uk/54273/>.
- [4] Danon L, House TA, Read JM, Keeling MJ. Social encounter networks: collective properties and disease transmission. J R Soc Interface. 2012;9(76):2826–2833.
- [5] Danon L, Read JM, House TA, Vernon MC, Keeling MJ. Social encounter networks: characterizing Great Britain. Proc R Soc B Biol Sci. 2013;280(1765):20131037.