Dear Valued Customers:

Your gene construct(s) has special growth protocols.

Re-plating:

A) Re-suspend lyophilized polynucleotide (4ug) in 40ul of 10mM Tris (pH 8.5). Final concentration is ~100ng/ul. This is the original stock. Please note, 4ug is an approximate amount. To accurately determine the quantity of DNA present, please measure by optical density at OD260nm.

B) In a new micro centrifuge tube, make a 1: 10 dilution of original stock in 10mM Tris (pH 8.5). This is 1:10 diluted stock.

C) To plate DNA, transform 2ul of original stock DNA and 2ul 1:10 diluted stock DNA into the appropriate competent cell. This is with a minimum transformation competency at 10^8-10^9 cfu/ug. Incubate mixture on LB agar plates (with desired antibiotic selection) at**22oC or room temperature**for**24hr to 48 hr**.

D) Select a well separated, **SINGLE** colony and inoculate in LB medium with desired antibiotic selection for large culture. Grow again at 22°C or room temperature for 24hr to 48 hr.

It is important to select only a single colony

E) Purify DNA from large culture. Verify sequences and continue with downstream applications.

If you encounter any problems, please do not hesitate to contact us. We will be more than happy to assist you.

Gene Synthesis Department
Bio Basic Inc.

20 Konrad Cres., Markham Ontario L3R 8T4 Canada
Tel: (905) 474 4493, (800) 313 7224 Fax: (905) 474 5794
Email: order@biobasic.com Web: www.biobasic.com