

Physics-Informed Neural Differential Equations and Multimodal Spectral Sensing for Autonomous Quality and Freshness Forecasting in Colombian Specialty Coffee

Autor(es): Prof. Dr. Isabel Suárez — Deggendorf Institute of Technology (DIT), AI Campus, September 17, 2026
Prof. Dr. Ricardo Vivas — Universidad de Cartagena (UdC), Faculty of Exact and Natural Sciences

Objective

To formulate, benchmark, and validate a physics-regularized deep learning framework coupling Attenuated Total Reflection Fourier-Transform Infrared (ATR-FTIR) spectroscopy with standardized physicochemical reference assays in Colombian Specialty Coffee (**Coffea arabica**). The investigation centers on systematically benchmarking classical chemometrics (PLSR, SVR) against black-box deep learning (1D-CNN) and physics-informed neural networks (PINNs regularized by Arrhenius degradation kinetics for chlorogenic acids and lipid peroxidation) under accelerated storage conditions, complemented by the exploratory evaluation of emerging sequence models (Mamba SSM, Spectral Transformers) and bilateral capacity building between the Deggendorf Institute of Technology (DIT) and the Universidad de Cartagena (UdC).

1. Executive Summary and Project Framing

1.1 Bioeconomic Motivation and Quality Challenges in Specialty Coffee

The Republic of Colombia is renowned globally for its production of high-altitude Arabica coffee (**Coffea arabica**). The coffee agro-industry serves as the socio-economic foundation of rural Colombia, sustaining over 550,000 smallholder farming families across Andean mountain ranges (Huila, Nariño, Cauca, Eje Cafetero, Santander, and Sierra Nevada de Santa Marta). Specialty coffees commanding cup scores above 85 points according to Specialty Coffee Association (SCA) standards achieve substantial market price premiums in European and particularly Bavarian specialty markets.

However, preserving this sensory excellence through long-term storage, sea freight logistics (20 to 45 days transit from Cartagena de Indias to European ports), and tropical warehouse conditions remains an acute challenge. Environmental fluctuations (20 – 40 °C, 50 – 85% RH) trigger complex degradation cascades:

- 1. Oxidative Degradation of Coffee Lipids:** Coffee contains 12 – 17% lipids (predominantly unsaturated linoleic fatty acids and diterpene cafestol/kahweol esters) that undergo free-radical peroxidation, forming hydroperoxides, conjugated dienes, and staling aldehydes (hexanal).

2. **Hydrolysis and Lactonization of Chlorogenic Acids (CGA):** Bioactive caffeoylquinic acids (5-CQA, 3-CQA, 4-CQA) undergo hydrolytic cleavage into quinic and caffeic acids, or lactonize into astringent quinides, impairing cup acidity and generating metallic off-notes.
3. **Moisture Adsorption and Water Activity (a_w):** Fluctuations in water activity accelerate non-enzymatic browning and enzymatic decay.

Standard quality monitoring relies either on subjective sensory cupping or expensive, destructive chromatography (HPLC, GC-MS costing 150 – 300 USD per sample), rendering routine longitudinal monitoring unfeasible for rural cooperatives.

Central Scientific Research Question

Core Research Question: To what extent can deep learning and physics/chemistry-informed neural models (PINNs incorporating Arrhenius degradation kinetics) reliably forecast coffee quality alterations from longitudinal ATR-FTIR spectra, and how do they benchmark against traditional chemometric methods (PLSR, SVR) in terms of predictive accuracy, out-of-distribution generalization, and causal spectral interpretability?

1.2 Seed Grant Scope and Strategic Scalability Roadmap

This 12-month seed project (8,610 EUR) focuses exclusively on **Colombian Specialty Coffee (*Coffea arabica*)** to maximize experimental depth and analytical rigor.

The validated methodology—coupling ATR-FTIR, AOAC wet-chemical standards, and physics-informed models—provides the empirical and algorithmic foundation for a subsequent 3-year bilateral grant application under the **DFG-Minciencias Bilateral Research Grants** programme. In that scaled initiative, the framework will be expanded to three additional strategic Colombian agri-food matrices: Fine Cocoa (***Theobroma cacao* L.**), Artisanal Honey (***Apis mellifera***), and Virgin Coconut Oil (***Cocos nucifera***), integrating high-throughput chromatography (LC-HRMS, GC-MS).

2. Botanical, Phytochemical, and Degradation Fundamentals of *Coffea arabica*

The experimental study evaluates certified Colombian cultivars: **Castillo**, **Caturra**, **Colombia**, and **Geisha**. Their phytochemical matrix comprises:

- **Chlorogenic Acids (CGA):** 6 – 9% dry weight, dominated by 5-caffeoylquinic acid (5-CQA).
- **Lipid Fraction:** 12 – 17%, rich in linoleic acid (C18:2) and cafestol/kahweol diterpenes.
- **Alkaloids and Methylxanthines:** Caffeine (1.0 – 1.4%), trigonelline (0.8 – 1.2%).
- **Carbohydrates:** Sucrose (6 – 9%) and structural polysaccharides.

2.1 Kinetic Formulation and Arrhenius Temperature Dependence

The degradation kinetics are governed by n -th order rate equations linked to the Arrhenius relationship:

$$\frac{dC_{i(t)}}{dt} = -k_{i(T)} \cdot [C_{i(t)}]^{n_i}, \quad k_{i(T)} = A_i \cdot \exp\left(-\frac{E_{a,i}}{R \cdot T}\right)$$

where $C_{i(t)}$ represents the concentration of bioactive constituent i (e.g., 5-CQA or unoxidized lipids), n_i is the apparent reaction order, A_i is the pre-exponential frequency factor (s^{-1}), $E_{a,i}$ is the apparent activation energy (kJ/mol), $R = 8.314 \frac{\text{J}}{\text{mol} \cdot \text{K}}$, and T is temperature in Kelvin.

2.2 ATR-FTIR Vibrational Spectroscopy of Coffee

Diamond Attenuated Total Reflection Fourier-Transform Infrared (ATR-FTIR) spectroscopy captures the 4000 – 600 cm^{-1} mid-infrared spectral region non-destructively:

Wavenumber (cm ⁻¹)	Vibrational Mode	Functional Group	Kinetic Trajectory During Storage Aging
3400 – 3250	$\nu(\text{O-H})$ stretching	Water (a_w) and phenolic OH (5-CQA)	Intensity decreases upon desiccation; broadens during moisture sorption
3008	$\nu(= \text{C-H})$ <i>cis</i> -alkene	Unsaturated fatty acid acyl chains	Progressive signal loss driven by radical lipid peroxidation
2925, 2854	$\nu_{\text{as}}, \nu_{\text{s}}(\text{C-H})$ $\frac{\text{CH}_2}{\text{CH}_3}$	Aliphatic hydrocarbon chains in lipids	Thermostable reference bands utilized for internal standard normalization
1745	$\nu(\text{C=O})$ ester carbonyl	Triacylglycerols and diterpene esters	Band broadening and emergence of free acid shoulder at 1715 cm ⁻¹
1715 – 1695	$\nu(\text{C=O})$ carboxylic acid	Liberated quinic acid and free fatty acids	Emergence and intensity rise tracking coffee bean staling
1650 – 1600	$\nu(\text{C=C})$ aromatic / $\nu(\text{C=O})$	Aromatic ring in caffeic acid moieties	Signal decline due to polyphenol oxidation and quinone polymerization
1380 – 1150	$\nu(\text{C-O}), \delta(\text{O-H})$	Caffeoylquinic ester linkages	Significant decrease correlated with HPLC chlorogenic acid depletion
1050 – 1030	$\nu(\text{C-O})$ carbohydrate	Sucrose and structural polysaccharides	Gradual signal decay associated with non-enzymatic browning

3. Standardized Physicochemical Reference Assays (AOAC / ISO)

- The Universidad de Cartagena conducts established standard analytical assays to calibrate spectral predictions:
- 1. Titratable Acidity (AOAC 920.92):** Aqueous extraction (10 g coffee in 100 mL degassed water) followed by potentiometric titration with 0.1 N NaOH to *pH* 8.20.
 - 2. Moisture Content & Water Activity (AOAC 968.11 / ISO 6673):** Thermogravimetric loss-on-drying at 105 °C; water activity (a_w) measured via chilled-mirror dewpoint sensor.
 - 3. Peroxide Value of Coffee Lipids (AOAC 965.33):** Cold-solvent lipid extraction and iodometric titration with sodium thiosulfate ($\text{Na}_2\text{S}_2\text{O}_3$) against starch indicator ($\text{meq } \frac{\text{O}_2}{\text{kg lipid}}$).
 - 4. Total Chlorogenic Acids (Folin-Ciocalteu / UV Spectrophotometry after AOAC 980.23):** Spectrophotometric quantification at 765 nm and 325 nm calibrated against pure 5-CQA standards.

4. Critical Literature Review, Research Gaps, and Central Hypothesis

- The synergy between vibrational spectroscopy and artificial intelligence represents a high-growth frontier in food analytics:
- **Classical Chemometrics:** Conventional studies rely almost exclusively on Principal Component Analysis (PCA) and Partial Least Squares Regression (PLSR) for static classification and adulteration testing ([1], [2], [3]). These models operate on static snapshots, failing to capture dynamic degradation kinetics.
 - **Convolutional Architectures:** Passos ([4]) demonstrated that 1D-CNNs with multiscale kernels effectively capture overlapping vibrational modes, but cautioned against severe overfitting when trained on limited spectral samples without inductive bias.
 - **Attention Mechanisms:** Upadhyay & Chaudhary ([5]) pioneered Spectral Attention Mechanisms (SAM), proving that self-attention layers successfully align neural weights with specific vibrational frequencies.
 - **State-Space Sequence Models:** Li et al. ([6]) introduced ASGMamba, showcasing linear $\mathcal{O}(L)$ computational complexity for multivariate time-series, which is ideally suited for long continuous aging trajectories.
 - **Physics-Informed Machine Learning (PIML):** Perré ([7]) established PINN regularizers for moisture diffusion. Pronk & Anthony ([8]) demonstrated that placing neural networks as residual corrections in coffee roasting models prevents thermodynamic violations. Felizzato et al. ([9]) and Grundy et al. ([10]) further highlighted multi-sensor data fusion imperatives.

Critical Research Gap: Prior literature lacks a rigorous benchmark comparing classical chemometrics against physics-regularized deep learning on real longitudinal FTIR coffee aging spectra under tropical conditions.

5. Exploratory Metamodel and 3-Tier Architecture Benchmark

The project implements a structured 3-tier comparative benchmark:

Benchmark Tier	Model Family	Configuration and Analytical Role
Tier 1: Baseline	Classical Chemometrics (Linear & Kernel ML)	PLSR (5–12 latent variables), Support Vector Regression (SVR, RBF kernel), and Random Forest on SNV + Savitzky-Golay preprocessed spectra.
Tier 2: Deep Learning	Empirical Deep Learning (Data-Driven Networks)	1D-CNN with multiscale receptive fields (3, 7, 15, 31 spectral channels) and MLP regression heads without physical constraints.
Tier 3: PIML	Physics-Informed DL (PINN with Kinetic Loss)	Hybrid PINN minimizing $\mathcal{L}_{\text{total}} = \mathcal{L}_{\text{data}} + \lambda \mathcal{L}_{\text{kinetic}}$, embedding pseudo-first order Arrhenius rate equations as soft-penalties.
Exploratory Screening	Advanced Sequence Models & NODEs (LSTM / Mamba / Transformers / Neural ODEs)	Modular evaluation of recurrent networks (LSTM/GRU), Mamba State-Space Models ([6]), Spectral Transformers with SAM attention maps ([5]), and continuous Neural ODEs (NODE).

5.1 PINN Loss Function Formulation

The PINN framework optimizes:

$$\mathcal{L}_{\text{total}} = \frac{1}{N} \sum_{i=1}^N (y_i - \hat{y}_i)^2 + \lambda_1 \frac{1}{M} \sum_{j=1}^M \left(\frac{d\hat{C}_j}{dt} + k(T)\hat{C}_j \right)^2 + \lambda_2 \mathcal{R}_{\text{monotonic}}$$

where $\mathcal{L}_{\text{kinetic}}$ penalizes deviations from theoretical Arrhenius degradation, and $\mathcal{L}_{\text{monotonic}}$ enforces non-increasing trajectories for chlorogenic acids in closed storage.

5.2 Multi-Objective Pareto Selection and Edge AI Deployment

Model selection optimizes three Pareto axes:

- Generalization Accuracy:** Cross-validation RMSE and $R^2 \geq 0.90$ on **Leave-One-Batch-Out** splits.
- Causal Interpretability:** Grad-CAM and SHAP attribution maps verifying focus on diagnostic bands (1745, 1715, 1600 – 1650 cm^{-1}).
- Edge Deployment:** FP16/INT8 quantized execution on a Raspberry Pi 4B/5 single-board computer achieving latency < 2.0 seconds and RAM usage < 500 MB.

6. Bilateral Cooperation, Workplan, and Budget Allocation

6.1 Institutional Synergies and Research Leadership

- Technische Hochschule Deggendorf (DIT) — Prof. Dr. Isabel Suárez:** Mathematical formulation, GPU cluster training, PINN and Mamba architecture implementation, XAI spectral attribution, and delivery of the on-site workshop in Cartagena.
- Universidad de Cartagena (UdC) — Prof. Dr. Ricardo Vivas:** Sourcing and botanical verification of specialty coffee cultivars, accelerated aging chambers (25, 40, 60 °C), longitudinal ATR-FTIR scanning, and AOAC analytical assays.

6.2 Transatlantic Mobility and Researcher Training

- Mission 1 (DIT -> UdC, Month 2, 10 days):** Research visit by Prof. Dr. Isabel Suárez to Cartagena. Delivery of a 1-week intensive workshop on «Applied AI and Physics-Informed Deep Learning for Spectroscopic Agro-Food Analytics» for UdC researchers and graduate students; standardization of laboratory measurement protocols.

2. **Mission 2 (UdC -> DIT, Months 8–9, 21 days):** 3-week research stay of an early-career researcher from Prof. Vivas's team at DIT's AI Campus. Hands-on GPU neural network training, XAI attribution validation, and co-drafting the DFG-Minciencias grant application.

6.3 12-Month Workplan and Key Milestones

- **WP1 (Months 1–4, UdC/DIT):** Protocol harmonization, coffee sampling, Workshop 1 in Cartagena with Prof. Suárez (M2). **Milestone M1 (M4):** Standardized protocol and pilot dataset established.
- **WP2 (Months 3–8, UdC):** Accelerated storage trials (180 days), periodic FTIR scans, and AOAC wet chemistry. **Milestone M2 (M8):** Longitudinal coffee spectral database completed.
- **WP3 (Months 6–11, DIT/UdC):** 3-tier benchmark execution, research stay of UdC junior researcher at DIT (M8–9). **Milestone M3 (M9):** Optimal Pareto architecture identified.
- **WP4 (Months 10–12, DIT/UdC):** Bilateral synthesis, two open-access manuscripts submitted, DFG-Minciencias grant drafted. **Milestone M4 (M12):** Submission of follow-up grant and final BAYLAT report.

6.4 Detailed Budget Plan (Total Requested: 8,610 €)

The budget strictly follows BAYLAT regulations and matches
06_Reiseverwaltung_Kostenkalkulation_Anschubfinanzierung.xlsx:

Budget Category	Mission 1 (DIT->UdC)	Mission 2 (UdC->DIT)	Total	Basis of Calculation
Airfare (Round-trip)	1,600 €	1,700 €	3,300 €	Transatlantic economy airfare (Munich-Cartagena / Cartagena-Munich)
Accommodation Costs	1,100 €	1,365 €	2,465 €	10 nights @ 110 € in Cartagena / 21 nights @ 65 € in Deggendorf
Per Diem (Subsistence)	500 €	945 €	1,445 €	10 days @ 50 € (Colombia) / 21 days @ 45 € (Germany) per BayRKG
Hospitality / Workshop	400 €	-	400 €	Catering and organization for on-site AI workshop at UdC
Travel Insurance	100 €	100 €	200 €	International health insurance (maximum 200 € per BAYLAT rules)
Laboratory Materials	-	-	600 €	Reference standards and reagents for coffee aging (maximum 600 €)
Digital Conference Costs	-	-	200 €	Streaming accessories for hybrid workshop delivery and collaboration
Total BAYLAT Funding	3,700 €	4,110 €	8,610 €	Total funding requested from BAYLAT (Deckblatt & Belegliste)

6.5 Pathway to DFG-Minciencias Follow-Up Funding

The models and experimental databases matured in this seed project will serve as the indispensable preliminary empirical evidence for a 3-year bilateral grant application under the **DFG-Minciencias Bilateral Research Grants** programme. The follow-up project will transfer the validated PIML methodology to **Fine Cocoa, Artisanal Honey, and Virgin Coconut Oil**, coupling low-cost spectroscopy with high-resolution chromatographic metabolomics (LC-HRMS, GC-MS).

References

- [1] V. Boadu, «Developing a novel onsite detection technology by using chemometrical analysis of hand-held near-infrared sensor technique for assessing coffee quality», *Unknown journal*, 2024.
- [2] L. Feng, B. Wu, S. Zhu, Y. He, y C. Zhang, «Application of Visible/Infrared Spectroscopy and Hyperspectral Imaging With Machine Learning Techniques for Identifying Food Varieties and Geographical Origins», *Frontiers in Nutrition*, vol. 8, jun. 2021, doi: 10.3389/fnut.2021.680357.
- [3] S. Widiarto, L. Zakaria, S. H. Jamal, D. S. Pratama, y I. Saputra, «Detection of Adulteration in Coffee Products Using FTIR Spectroscopy and Multivariate Analysis», *Jurnal Kimia Sains dan Aplikasi*, vol. 29, n.º 4, pp. 296-305, abr. 2026, doi: 10.14710/jksa.29.4.296-305.

-
- [4] D. Passos, «Convolutional neural networks in Vis–NIR chemometrics: From contradiction to conditional design», *TrAC Trends in Analytical Chemistry*, vol. 203, p. 119015, oct. 2025, doi: 10.1016/j.trac.2026.119015.
- [5] A. Upadhyay y N. Chaudhary, «Enhancing spectral analysis of ghee adulteration via a deep learning-based multimodal attention mechanism», *Scientific Reports*, vol. 16, n.º 1, may 2026, doi: 10.1038/s41598-026-48661-6.
- [6] Q.-Y. Li, X. Zhang, S.-X. Wang, J. Wei, y Y. Xing, «ASGMamba: Adaptive Spectral Gating Mamba for Multivariate Time Series Forecasting», *ArXiv*, vol. abs/2602.01668, feb. 2026, doi: 10.48550/arxiv.2602.01668.
- [7] P. Perré, «Toward mechanistic models “augmented” by machine learning. Example of a drying simulation data set exploited by a Physics Informed Neural Network», *Drying Technology*, vol. 43, n.º 1–2, pp. 147-161, oct. 2025, doi: 10.1080/07373937.2024.2411582.
- [8] M. Pronk y B. W. Anthony, «Neural-network placement in physics-informed machine learning for mechanistic process-model repair: a case study in industrial coffee roasting», *Scientific Reports*, sep. 2026, doi: 10.1038/s41598-026-67034-7.
- [9] G. Felizzato, E. Bagnulo, A. Guglielmetti, y C. Bortolini, «Data Fusion for the Food Industry: Leveraging Machine Learning and Deep Learning Algorithms», *Unknown journal*, 2025.
- [10] H. H. Grundy, H. J. Hird, V. Bailey-Horne, M. Sykes, y A. Charlton, «Review of Current and Emerging Analytical Methods for the Testing of Edible Oil for Authenticity», *FSA Research and Evidence*, oct. 2025, doi: 10.46756/001c.145026.