


Copyright for test papers and marking guides remains with *West Australian Test Papers*.
Test papers may only be reproduced within the purchasing school according to the advertised Conditions of Sale.
Test papers should be withdrawn after use and stored securely in the school until 15th October 2020.



SEMESTER TWO

MATHEMATICS SPECIALIST UNITS 3 & 4

2020

SOLUTIONS

Calculator-Free Solutions

$$1. \quad u = \sec(\theta) \rightarrow \frac{du}{d\theta} = \frac{-\sin \theta}{-\cos^2 \theta} \rightarrow du = \sec \theta \tan \theta d\theta \quad \checkmark$$

$$u(\theta) = \frac{1}{\cos \theta} \rightarrow u(0) = 1 \wedge u\left(\frac{\pi}{3}\right) = \frac{1}{0.5} = 2 \quad \checkmark$$

$$\therefore \int_0^{\frac{\pi}{3}} \sec \theta \tan \theta d\theta = \int_1^2 du = [u]_1^2 = 2 - 1 = 1 [4]$$

$$2. \quad P(z) \text{ has real coefficients} \Rightarrow \bar{z} = 2 - i \text{ is also a solution.} \quad \checkmark$$

$$\therefore (z - 2 - i)(z - 2 + i) = z^2 - 4z + 5 \text{ is a factor of } P(z) \quad \checkmark$$

$$\begin{array}{r}
 z^2 - 4z + 5 \quad \begin{array}{r}
 \begin{array}{rrrrr}
 z^2 & -2z & 5 & & \\
 \hline
 z^4 & -6z^3 & 18z^2 & -30z & 25 \\
 z^4 & -4z^3 & 5z^2 & & \\
 \hline
 0 & -2z^3 & 13z^2 & -30z & 25 \\
 & -2z^3 & 8z^2 & -10z & \\
 \hline
 & 0 & 5z^2 & -20z & 25 \\
 & & 5z^2 & -20z & 25 \\
 \hline
 & & 0 & 0 & 0
 \end{array}
 \end{array}
 \end{array}
 \quad \checkmark \checkmark$$

$$\therefore z^2 - 2z + 5 = 0$$

$$z = \frac{2 \pm \sqrt{4 - 20}}{2} = 1 \pm 2i$$

$$\text{Solutions are } z = 1 \pm 2i, 2 \pm i \quad \checkmark$$

[5]

$$3. \quad \frac{2x^2 + 1}{x(x^2 + 1)} = \frac{Ax + B}{x^2 + 1} + \frac{C}{x} \rightarrow 2x^2 + 1 = x(Ax + B) + C(x^2 + 1) \quad \checkmark$$

$$x = 0 \rightarrow 1 = 0 + C \rightarrow C = 1 \quad \checkmark$$

$$x = 1 \rightarrow 3 = A + B + 2$$

$$x = -1 \rightarrow 3 = A - B + 2$$

$$\therefore A = 1, B = 0 \quad \checkmark$$

$$\therefore \int_1^2 \frac{2x^2 + 1}{x^3 + x} dx = \int_1^2 \left(\frac{x}{x^2 + 1} + \frac{1}{x} \right) dx = \left[\frac{1}{2} \ln(x^2 + 1) + \ln|x| \right]_1^2$$

$$\therefore \frac{1}{2} \ln(5) + \ln(2) - \frac{1}{2} \ln(2) = \frac{1}{2} \ln 10 = \ln \sqrt{10} \quad \text{as required} \quad \checkmark \quad [5]$$


4. (a) $f(g(x)) = f(-|x|) = |x|^2 - 2|x| - 3 = x^2 - 2|x| - 3$ ✓

For $x \geq 0$, $f(g(x)) = x^2 - 2x - 3 = (x-1)^2 - 4$

which has a turning point at $(1, -4)$

$\therefore D_x = R$ and $R_y = \{y \in R : y \geq -4\}$ ✓✓

(b)



✓
Parabolic shape

✓
Mirror image over the y-axis

✓
Local min at $(\pm 1, -4)$, and roots at $x = \pm 3$

(c) $f(x)$ has its turning point at $(-1, -4)$, $\therefore k = -1$ ✓

$\therefore f^{-1}(x) = \sqrt{x+4} - 1$ ✓

(d) (i) $g(f^{-1}(0)) = g(1) = -|1| = -1$ ✓

(ii) $g(f^{-1}(f(x+1))) = g(x+1) = -|x+1|$ ✓ [10]

5. (a) R is a solid sphere centred at $(1, 1, 0)$ with radius 3 units. ✓✓

$|r|_{\max} = \left| \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} \right| + 3 = \sqrt{2} + 3$ units ✓

(b) xz plane $\rightarrow y=0$, therefore the intersection is given by

$(x-1)^2 + (0-1)^2 + (z)^2 = 9$

$(x-1)^2 + z^2 = 8$ ✓

\therefore circle centred at $(1,0,0)$ and radius $2\sqrt{2}$ units.

✓

5. (c) The line is given by $r = \begin{pmatrix} 5+\lambda \\ 2 \\ -\lambda \end{pmatrix}$, hence:

$$\left| \begin{pmatrix} 5+\lambda \\ 2 \\ -\lambda \end{pmatrix} - \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} \right| = 3$$

$$\therefore (4+\lambda)^2 + 1^2 + (-\lambda)^2 = 3^2 \quad \checkmark$$

$$2\lambda^2 + 8\lambda + 8 = 0$$

$$2(\lambda+2)^2 = 0 \rightarrow \therefore \lambda = -2 \quad \checkmark$$


Unique solution implies the line is tangent to the sphere R \checkmark

$$r(-2) = \begin{pmatrix} 5-2 \\ 2 \\ -2 \end{pmatrix} = \begin{pmatrix} 3 \\ 2 \\ 2 \end{pmatrix}$$

Point of tangency is $(3, 2, 2)$. \checkmark

[9]

6. (a)




\checkmark

Square root function with correct orientation, root at $x=3$ and y-intercept at $y=2$

\checkmark

Open boundary at $x=-5$
closed boundary at $x=4$




\checkmark

Mirror image over the y-axis, completely below the x-axis

\checkmark

y-intercept at $y=-2$,
open boundaries at ± 5 .

6. (b)



✓

Poles at $x = \pm 3$


✓

Closed boundaries at $x = \pm 4$ Y-intercept at $y = -0.5$

✓

Behaviour on either side of both poles

(c) $f(x)$ and $f^{-1}(x)$ are shown below, hence, for $f^{-1}(x)$:



✓

$$D_x = \{x \in \mathbb{R} : -4 < x \leq 2\}$$

✓

$$R_y = \{y \in \mathbb{R} : -5 < y \leq 4\}$$

[9]

7. (a) (i) The internal parabolic outline is given by $y = \frac{x^2}{4} - 4$

$$\therefore x^2 = 4(y + 4)$$

Hence, the volume is given by:

$$v_y = \pi \int_0^{21} x^2 dy = \pi \int_0^{21} 4(y + 4) dy = 4\pi \int_0^{21} (y + 4) dy$$

$$\therefore f(y) = y + 4 \text{ as required}$$

7. (a) (ii)

$$V_y = 4\pi \int_0^{21} (y+4) dy = 4\pi \left[\frac{y^2}{2} + 4y \right]_0^{21}$$

$$\therefore V_y = 4\pi \left[y \left(\frac{y}{2} + 4 \right) \right]_0^{21} = 4\pi \times 21 \times \left(\frac{21}{2} + 4 \right)$$

$$\therefore 2\pi \times 21 \times 29 = 1218\pi \text{ cm}^3 \quad \checkmark$$

(b) $V = V_{\text{curved side}} + V_{\text{base}}$

$$= (V_{\text{outer shell}} - V_{\text{inner shell}}) + V_{\text{base}}$$

$$\therefore V = \pi \int_0^{21} 5(y+5) dy - \pi \int_0^{21} 4(y+4) dy + \pi \int_{-1}^0 5(y+5) dy$$

[7]

8. (a) $a = \frac{dv}{dt} \times \frac{dx}{dx}$

$$\therefore \frac{dx}{dt} \times \frac{dv}{dx} = v \frac{dv}{dx} \quad \text{as required} \quad \checkmark$$


(b) $\frac{d}{dx} \rightarrow 4x^2 - v^2 = 1$

$$8x - 2v \frac{dv}{dx} = 0 \quad \checkmark \checkmark$$

$$\therefore v \frac{dv}{dx} = a = 4x \quad \checkmark \quad [4]$$


Calculator-assumed Solutions

9. (a)



- ✓
Semicircle
- ✓
Centred at (2,0) and radius 1
- ✓
Discontinuous at $x=3$
(or $x=-1$ for FT marks if drawn on LHS)

(b)



From diagram, $\arg(z-2)_{\max} = \frac{\pi}{6}$ and $|z-2| = \sqrt{3}$

✓✓

OR


$\arg(z-2)_{\max} = \pi$ and $|z-2| = \sqrt{3}$ for follow through marks if the circle is drawn on the LHS of the y-axis.

[5]

10. (a) $\left. \frac{dy}{dx} \right|_{\left(\frac{\pi}{2}, 2\right)} = 2 \times \sin \frac{\pi}{2} = 2$

✓

(b)



✓

Follows the isoclines to show a cyclic curve (must be wider along the y-axis, it cannot look like a sinusoid)

✓

Passes through the point P

(c) $\frac{dy}{dx} = y \sin x \rightarrow \frac{dy}{y} = \sin x \, dx$

$$\therefore \int \frac{1}{y} dy = \int \sin x \, dx$$

$$\therefore \ln|y| = -\cos x + C_1$$

✓

$$y = e^{-\cos x + C_1} = e^{C_1} \times e^{-\cos x} = C_2 e^{-\cos x}$$

$$\left(\frac{\pi}{2}, 2\right) \rightarrow 2 = C_2 \times e^0 \rightarrow C_2 = 2$$

$$\therefore y = 2e^{-\cos x}$$

✓

11. (a) (i) $z^4 = -1 = \text{cis}(\pi + 2k\pi), k = 0, \pm 1, -2$

✓


$$\therefore z = \text{cis}\left(\frac{\pi + 2k\pi}{4}\right), k = 0, \pm 1, -2$$

✓

$$\therefore z = \text{cis}\left(\frac{\pi}{4}\right), \text{cis}\left(\frac{3\pi}{4}\right), \text{cis}\left(\frac{-\pi}{4}\right), \text{cis}\left(\frac{-3\pi}{4}\right)$$

✓✓

11. (a) (ii) Solutions from (a)(i) move two units up:



✓
4 solutions equidistant
from $2i$

✓✓
equally spaced at $\frac{\pi}{2}$
radians from each other
within the circle centred
at $2i$

$$(b) \quad (i) \quad z = \frac{\sqrt{3}+i}{1-i} \times \frac{1+i}{1+i} = \left(\frac{\sqrt{3}-1}{2}\right) + \left(\frac{\sqrt{3}+1}{2}\right)i$$

$$\therefore \Re(z) = \frac{\sqrt{3}-1}{2} \wedge \Im(z) = \frac{\sqrt{3}+1}{2}$$

(or CAS)

✓✓

$$(ii) \quad z = \frac{\sqrt{3}+i}{1-i} = \frac{2 \operatorname{cis}\left(\frac{\pi}{6}\right)}{\sqrt{2} \operatorname{cis}\left(\frac{-\pi}{4}\right)} = \frac{2}{\sqrt{2}} \operatorname{cis}\left(\frac{\pi}{6} + \frac{\pi}{4}\right) = \sqrt{2} \operatorname{cis}\left(\frac{5\pi}{12}\right)$$

✓✓

$$\therefore \sqrt{2} \cos\left(\frac{5\pi}{12}\right) = \Re(z) = \frac{\sqrt{3}-1}{2}$$

✓

$$\cos\left(\frac{5\pi}{12}\right) = \frac{\sqrt{3}-1}{2\sqrt{2}} = \frac{\sqrt{2}}{4}(\sqrt{3}-1) \text{ as required}$$

✓

$$12. \quad (a) \quad r(t) = \int v(t) dt = \left(\int 2t dt\right)i + \left(\int \sin t dt\right)j$$

$$i(t^2 + C_1)i + (-\cos t + C_2)j$$

✓

$$r(0) = \begin{pmatrix} C_1 \\ -1 + C_2 \end{pmatrix} = \begin{pmatrix} 2 \\ 1 \end{pmatrix} \rightarrow \begin{matrix} C_1 = 2 \\ C_2 = 2 \end{matrix}$$

✓

$$\therefore r(t) = (t^2 + 2)i + (2 - \cos t)j$$

✓

- (b) $a(t) = \frac{d}{dt} v(t) = 2i + \cos t j$ ✓
- $\therefore a(0) = 2i + \cos 0 j = 2i + j$ ✓
- $\therefore |a(0)| = \left| \begin{pmatrix} 2 \\ 1 \end{pmatrix} \right| = \sqrt{5} \approx 2.24 \text{ m/s}^2$ ✓
12. (c) $y_{\max} = 2 - (-1) = 3$ for $\cos t = -1 \rightarrow t = \pi$ ✓
- $\therefore a = \pi^2 + 2$ and $b = 3$ ✓✓
- (d) $\int_0^{10} \sqrt{4t^2 + \sin^2 t} dt \approx 100.43 \text{ m}$ ✓✓✓
13. (a) \bar{T} is approximately a normal distribution by the Central Limit Theorem since $n = 100 > 30$ ✓
- $\therefore \bar{T} \sim N\left(37, \frac{0.25^2}{100}\right) = N(37, 6.25 \times 10^{-4})$ ✓
- with $\sigma(\bar{T}) = \frac{1}{40} = 0.025$ ✓
- (b) $P(\bar{T} > 37.005) = P\left(z > \frac{37.005 - 37}{0.0025}\right) = P(z > 0.2)$ ✓
- $\therefore P(z > 0.2) \approx 0.42074$ ✓
- (c) Since $n > 100 \Rightarrow \sigma(\bar{T}) < 0.025$
- \therefore the answer in (b) would be lower/smaller ✓
- because the lower standard deviation of the sample mean ✓
- (d) Require $P(37 < \bar{T} < 37.005) > 0.45$
- $\therefore P(0 < z < k) > 0.45$ ✓
- CAS $\rightarrow k > 1.6449$
- $\frac{37.005 - 37}{\left(\frac{0.25}{\sqrt{n}}\right)} > 1.6449$ ✓
- $\rightarrow n > 6764.2 \approx n > 6764$ patients ✓
- [Accept 6764 and 6765 depending on accuracy used]

$$14. \quad (a) \quad \vec{OA} = \begin{pmatrix} 10 \\ -10 \\ 0 \end{pmatrix} + 2 \begin{pmatrix} -2 \\ 1 \\ 3 \end{pmatrix} = \begin{pmatrix} 16 \\ -8 \\ 6 \end{pmatrix} \quad \checkmark$$

$$\vec{OB} = \begin{pmatrix} -10 \\ 30 \\ 0 \end{pmatrix} + 2 \begin{pmatrix} 3 \\ -2 \\ 4 \end{pmatrix} = \begin{pmatrix} -4 \\ 26 \\ 8 \end{pmatrix} \quad \checkmark$$

$$\therefore \vec{AB} = \begin{pmatrix} -4 \\ 26 \\ 8 \end{pmatrix} - \begin{pmatrix} 16 \\ -8 \\ 6 \end{pmatrix} = \begin{pmatrix} -20 \\ 34 \\ 2 \end{pmatrix} \rightarrow \sqrt{\frac{-20}{34} + \frac{2}{2}} \approx 39.50 m \quad \checkmark \checkmark$$

$$14. \quad (b) \quad \vec{OA} = \begin{pmatrix} 20 \\ -10 \\ 0 \end{pmatrix} + t \begin{pmatrix} -2 \\ 1 \\ 3 \end{pmatrix} = \begin{pmatrix} 20-2t \\ t-10 \\ 3t \end{pmatrix}$$

$$|\vec{OA}| = \sqrt{\begin{vmatrix} 20-2t \\ t-10 \\ 3t \end{vmatrix}} = \sqrt{(20-2t)^2 + (t-10)^2 + (3t)^2} = 30 \quad \checkmark$$

$\therefore CAS \rightarrow t = 10 \text{ s later}$

$$\therefore \vec{OA} = \begin{pmatrix} 0 \\ 0 \\ 30 \end{pmatrix} \quad \checkmark$$

$$\therefore \vec{OB}(10) = \begin{pmatrix} -10 \\ 30 \\ 0 \end{pmatrix} + 10 \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 30 \end{pmatrix} \quad \checkmark$$

$$\therefore v_B = i - 3j + 3k \text{ m/s} \quad \checkmark$$

15. All coordinates in the first octant are positive. \checkmark

xy-plane intersection ($z=0$):

$$3 + \lambda = 0 \rightarrow \lambda = -3$$

$$\therefore r = \begin{pmatrix} 4+6 \\ 2-3 \\ 3-3 \end{pmatrix} = \begin{pmatrix} 10 \\ -1 \\ 0 \end{pmatrix} \text{ not in the first octant} \quad \checkmark$$

xz-plane intersection ($y=0$):

$$2 + \lambda = 0 \rightarrow \lambda = -2$$

$$\therefore r = \begin{pmatrix} 4+4 \\ 2-2 \\ 3-2 \end{pmatrix} = \begin{pmatrix} 8 \\ 0 \\ 1 \end{pmatrix} \text{ in the first octant} \quad \checkmark$$

yz-plane intersection ($x=0$):

$$4 - 2\lambda = 0 \rightarrow \lambda = 2$$

$$\therefore r = \begin{pmatrix} 4-4 \\ 2+2 \\ 3+2 \end{pmatrix} = \begin{pmatrix} 0 \\ 4 \\ 5 \end{pmatrix} \text{ in the first octant} \quad \checkmark$$

$$\therefore \begin{vmatrix} 8 \\ 0 \\ 1 \end{vmatrix} - \begin{vmatrix} 0 \\ 4 \\ 5 \end{vmatrix} = \begin{vmatrix} 8 \\ -4 \\ -4 \end{vmatrix} \rightarrow \begin{vmatrix} 8 \\ -4 \\ -4 \end{vmatrix} = 4 \begin{vmatrix} 2 \\ -1 \\ -1 \end{vmatrix} = 4\sqrt{6} \text{ units} \quad \checkmark\checkmark$$

$$16. \quad (a) \quad P(0) = \frac{80\,000}{5+3 \times 1} = 10\,000 \text{ cats} \quad \checkmark$$

$$(b) \quad P(t)_{\max} = \frac{80000}{5+3 \times 0} = 16\,000 \text{ cats}$$

✓

which is the carrying capacity of the system

16. (c) $P = \frac{80000}{5+3e^{-0.25t}} \rightarrow 5+3e^{-0.25t} = \frac{80000}{P}$
- $$\frac{dP}{dt} = -80000(5+3e^{-0.25t})^{-2} \times -0.75e^{-0.25t} \quad \checkmark$$
- $$\textcircled{i} \frac{80000}{(5+3e^{-0.25t})} \times \frac{1}{4} \times \frac{1}{(5+3e^{-0.25t})} \times 3e^{-0.25t} \quad \checkmark$$
- $$\textcircled{i} \frac{P}{4} \times \frac{P}{80000} \times \left(\frac{80000}{P} - 5 \right) \quad \checkmark \checkmark$$
- $$\textcircled{i} \frac{P}{4} \left(1 - \frac{P}{16000} \right) \text{ as required}$$
- (d) $\left. \frac{dP}{dt} \right|_{t=0} = \frac{10000}{4} \left(1 - \frac{10000}{16000} \right) = 2500 \times \frac{3}{8} = 937.5 \text{ cats/year} \quad \checkmark \checkmark$
17. (a) 90% confidence interval for μ :
- $$P(-k < z < k) = 0.9 \rightarrow k \approx 1.645 \quad \checkmark$$
- $$\therefore 20 - 1.645(2) < \mu < 20 + 1.645(2) \quad \checkmark$$
- $$16.71 < \mu < 23.29 \text{ min} \quad \checkmark$$
- (b) From July: $\frac{\sigma}{\sqrt{n}} = 2 \rightarrow \sigma = 2\sqrt{n}$
- For August: $\sigma(\bar{X}) = \frac{\sigma}{\sqrt{2n}} = \frac{1}{\sqrt{2}} \left(\frac{\sigma}{\sqrt{n}} \right) = \frac{1}{\sqrt{2}}(2) \approx 1.41 \text{ min} \quad \checkmark \checkmark$
- (c) C is the most accurate. \checkmark
- Because it is based on the smallest standard error \checkmark
- (d) We do not know which interval, if any, \checkmark
- because we cannot determine the true \checkmark
- value of the population mean.
18. (a) $-\omega^2 = -9 \rightarrow \omega = 3 \text{ rad/s} \quad \checkmark$
- $$\therefore T = \frac{2\pi}{\omega} = \frac{2\pi}{3} \approx 2.09 \text{ sec} \quad \checkmark$$

(b) Let $x(t) = A \sin(3t)$ ✓

$$\therefore x'(t) = 3A \cos(3t)$$

$$3A = \frac{3}{2} \rightarrow A = \frac{1}{2} m = 50 \text{ cm} \quad \checkmark$$

18. (c) Use $v^2 = \omega^2(A^2 - x^2)$:

$$6^2 = \omega^2(A^2 - \sqrt{3}^2) \text{ and } (6\sqrt{2})^2 = \omega^2(A^2 - \sqrt{2}^2) \quad \checkmark$$

$$CAS \rightarrow \omega = 3 \rightarrow T = \frac{\pi}{3} \text{ s and } A = 2 \text{ m} \quad \checkmark \checkmark$$

19. (a) Outline of the circle in x/y plane given by:

$$x^2 + y^2 = r^2 \rightarrow y = \pm \sqrt{r^2 - x^2} \quad \checkmark$$

$$\therefore 2x + 2y \frac{dy}{dx} = 0 \rightarrow \frac{dy}{dx} = \frac{-x}{y} \quad \checkmark$$

$$\therefore A = \int_a^b 2\pi y \sqrt{1 + \frac{x^2}{y^2}} dx \quad \checkmark$$

(b) Use $y^2 = r^2 - x^2$:

$$\therefore A = \int_a^b 2\pi y \sqrt{\frac{y^2 + x^2}{y^2}} dx = \int_a^b 2\pi \sqrt{y^2 + x^2} dx \quad \checkmark$$

$$\textcolor{red}{\therefore} \int_a^b 2\pi \sqrt{r^2 - x^2 + x^2} dx = \int_a^b 2\pi r dx \quad \checkmark$$

$$\textcolor{red}{\therefore} 2\pi r [x]_a^b = 2\pi r(b-a) \text{ as required.}$$

(c) $a = -r$ and $b = r$:

$$\therefore A = 2\pi r(r - (-r)) = 2\pi r \times 2r = 4\pi r^2 \text{ as required} \quad \checkmark$$

20. $\sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 + \left(\frac{dz}{dt}\right)^2} = k \text{ for } k \in R^+ \textcolor{red}{\text{ and }} \quad \checkmark$

$$\therefore \left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 + \left(\frac{dz}{dt}\right)^2 = k^2$$

$$\frac{d}{dt} \rightarrow 2\left(\frac{dx}{dt}\right)\frac{d^2x}{dt^2} + 2\left(\frac{dy}{dt}\right)\frac{d^2y}{dt^2} + 2\left(\frac{dz}{dt}\right)\frac{d^2z}{dt^2} = 0 \quad \checkmark \checkmark$$

$$2 \begin{pmatrix} \frac{dx}{dt} \\ \frac{dy}{dt} \\ \frac{dz}{dt} \end{pmatrix} \cdot \begin{pmatrix} \frac{d^2x}{dt^2} \\ \frac{d^2y}{dt^2} \\ \frac{d^2z}{dt^2} \end{pmatrix} = 0 \quad \checkmark$$

$$\therefore v(t) \cdot a(t) = 0 \rightarrow v(t) \perp a(t) \text{ as required} \quad \checkmark$$

[5]