

Copyright for test papers and marking guides remains with *West Australian Test Papers*.
Test papers may only be reproduced within the purchasing school according to the advertised Conditions of Sale.
Test papers should be withdrawn after use and stored securely in the school until 14th October 2019.



MATHEMATICS METHODS UNITS 3 & 4

Semester Two

2019

SOLUTIONS

4. (a) (i) $(\ln x)^2 + 2\ln x - 3 = 0$
 Let $\ln x = m$
 $(m + 3)(m - 1) = 0$ ✓
 $\ln x = -3$ or $\ln x = 1$ ✓
 $x = \frac{1}{e^3}$ or $x = e$ ✓
- (ii) $2^5 = 3x - 4$ ✓
 $36 = 3x$ ✓
 $x = 12$ ✓
- (b) $g(x) = \ln 5 - 2 \ln x$ ✓
 $g'(x) = -\frac{2}{x}$ therefore $m = -\frac{2}{e}$ at $(e, \ln 5 - 2)$ ✓
 $\ln 5 - 2 = -\frac{2}{e}(e) + c$
 $y = -\frac{2}{e}x + \ln 5$ ✓ [8]
5. (a) $\frac{1-a}{1-b}$ ✓✓
- (b) (i) False ✓
 In a very large number of samples, 95% of those confidence intervals would contain p . The endpoints of the confidence interval refer to the confidence and not the probability. p is not a random variable and is either in the CI or not. ✓
- (ii) False ✓
 The confidence interval is about the population proportion and not about the individual Australian households ✓
- (iii) True ✓
- (iv) True ✓ [8]
 (the centre of the interval contains the sample proportion)

6. (a) $2 - 4\cos t = 0 \therefore \cos t = \frac{1}{2}$
- $t = \frac{\pi}{3} \text{ or } \frac{5\pi}{3}$ ✓✓
- (b) $a(t) = 4\sin t = 0$
- $t = 0 \text{ or } \pi \text{ or } 2\pi$ ✓
- $a'(0) > 0 \text{ Min} \quad a'(\pi) < 0 \text{ Max} \quad a'(2\pi) > 0 \text{ Min}$ ✓
- $v(\pi) = 2 - 4(-1) = 6$
- Max velocity is 6m/s ✓
- $$-\int_0^{\frac{\pi}{3}} 2 - 4\cos t \, dt + \int_{\frac{\pi}{3}}^{\pi} 2 - 4\cos t \, dt$$
- (c) ✓
- $$= \left[4\sin t - 2t \right]_0^{\frac{\pi}{3}} + \left[4\sin t - 2t \right]_{\frac{\pi}{3}}^{\pi}$$
 ✓
- $$= -\frac{2\pi}{3} + 2\sqrt{3} + \frac{4\pi}{3} + 2\sqrt{3}$$
 ✓
- $$= \frac{2\pi}{3} + 4\sqrt{3} \text{ m.}$$
 ✓ [9]
7. (a) $\int_{-3}^{-1} f(x)dx = 0 \text{ and } \int_2^3 f(x)dx = 0$
- $\int_{-1}^2 f(x)dx = 3$ ✓
- $\therefore \int_{-p}^{-3} f(x)dx + \int_3^p f(x)dx = -3$ ✓
- $1 \times 1.5 + 1 \times 1.5 = 3 \therefore p = 3 + 1.5$
- $p = 4.5$ ✓
- (b) $\frac{1}{3} \int_0^1 \frac{3x^2}{x^3 + 1} dx = \frac{1}{3} [\ln(x^3 + 1)]_0^1$ ✓
- $= \frac{1}{3} (\ln 2 - \ln 1)$ ✓

$$= \frac{1}{3} \ln 2$$

✓

[6]

8. (a) $\log_m ba^2 = \log_m 5$ ✓
- $b = \frac{5}{a^2}$ ✓
- (b) $\log_3 9 + \log_3 5 + 2\log_7 7 = 2 + \log_3 5 + 2$ ✓
- $= 4 + \log_3 5$ ✓ [4]
9. $4x^3 - 4x^2 + 3x = 2x$
- $x(4x^2 - 4x + 1) = 0$
- $x(2x - 1)^2 = 0$
- $x = 0$ or $\frac{1}{2}$ ✓
- $\int_0^{\frac{1}{2}} 4x^3 - 4x^2 + x \, dx = \left[x^4 - \frac{4x^3}{3} + \frac{x^2}{2} \right]_0^{\frac{1}{2}}$ ✓
- $= \frac{1}{16} - \frac{1}{6} + \frac{1}{8} = \frac{3 - 8 + 6}{48}$ ✓
- $= \frac{1}{48} \text{ units}^2$ ✓ [4]

Calculator-assumed Solutions

10. (a) $\hat{p} - z \sqrt{\left(\frac{\hat{p}(1 - \hat{p})}{500} \right)} = 0.2278$ and $\hat{p} + z \sqrt{\left(\frac{\hat{p}(1 - \hat{p})}{500} \right)} = 0.2922$ ✓
 $\hat{p} = 0.26$ ✓
 0.26 of 500 = 130. ✓
- (b) $z = 1.641$ Therefore 90 % confidence level. ✓✓ [5]
11. (a) (i) $n = 20$ $p = 0.63$
 $P(X \geq 14) = 0.3453$ ✓
- (ii) $P(X = 20 | X \geq 14) = \frac{0.000097}{0.3453} = 0.00028$ ✓✓
- (b) $P(X \geq 1) \geq 0.95$
 $1 - P(X = 0) \geq 0.95$
 $P(X = 0) \leq 0.05$ ✓
 $0.37^n \leq 0.05$ ✓
 $n = 3.01$
 4 people would need to be selected ✓
- (c) $E(X) = np$
 $= 300 \times 0.63$
 $= 189$ ✓
 $\sigma = \sqrt{npq} = \sqrt{300 \times 0.63 \times 0.37} = 8.36$ ✓
- (d) $0.05 = 1.645 \sqrt{\frac{0.63 \cdot (1 - 0.63)}{n}}$ ✓
 $n = 253$ credit card holders ✓ [10]
12. $\int_5^{10} 10 + 10e^{-x} - (5 - 0.02x) dx$ ✓✓
 $= 25.81693$
 The profit earned on sales between 500 and 1000 litres is \$25.82. ✓ [3]

13. (a) (i) $\left(\frac{3}{4}\right)^3 \left(\frac{1}{4}\right) = \frac{27}{256}$ or 0.1055 ✓
- (ii) $P(X = 4) + P(X = 5) + P(X = 6)$ ✓
- $= 0.1055 + 0.079101 + 0.059326$ ✓
- $= 0.243927$ ✓
- (b) (i) Discrete, independent probability ✓
- two outcomes: Success or fail ✓
- (ii) $E(X) = 0.25$ ✓
- Standard deviation = $\frac{\sqrt{3}}{4} = 0.43301$ ✓ [7]
14. (a) $\mu = \frac{6.675 \cdot 10^5}{75} = 8900$ litres ✓
- (b) Variance = $E(X^2) - (E(X))^2 = \frac{5.978 \cdot 10^9}{75} - 8900^2$ ✓
- $= 496\,666.67$ ✓
- Standard deviation = $704.75 \approx 705$ litres ✓
- (c) $P(X \leq 8550 \mid 7000 < X < 10\,000) = \frac{0.306268}{0.937135} = 0.3268$ ✓✓
- $-2.6 = \frac{x - 8900}{705}$
- (d) $x = 7067$ litres ✓
- Garage must sell 7068 litres to be profitable ✓ [7]
15. Underestimate: $1 \times (0 + 10 + 15 + 11 + 4) = 40$ ✓
- Over estimate: $1 \times (10 + 15 + 16 + 16 + 11) = 68$ ✓
- Average = 54
- Distance is therefore approximately 54 m. ✓ [3]
16. (a) Self-selective sample and therefore not representative of the population. ✓
- Only radio listeners were part of the survey, therefore not representative. ✓
- (b) Logical answer for example:
a randomly selected sample from the general public is used ✓
- (c) 0.03 ✓
- (d) $\hat{p} = 0.15$ Therefore point estimate = 0.15 or $\frac{45}{300}$ ✓
- (e) $0.03 = z \sqrt{\frac{0.15(1 - 0.15)}{300}}$ ✓
- $z = 1.45521$ ✓
- $P(-1.45521 < z < 1.45521) = 0.85439$
- 85.4% confidence level ✓ [8]

17. (a) At time t , the distance from B to S is $100 - 50t$. ✓
 At time t , the distance from A to S is $80t$. ✓

$$r^2 = (100 - 50t)^2 + (80t)^2 - 2(100 - 50t)(80t)\cos\frac{\pi}{3}$$
 ✓
 Cosine rule
 Therefore $r^2 = 12900t^2 - 18000t + 10000$

$$\frac{d(r^2)}{dt} = 25800t - 18000 = 0 \text{ or } \frac{dr}{dt} = \frac{5(258t - 180)}{\sqrt{129t^2 - 180t + 100}} = 0$$
 ✓
 (b) $t = \frac{30}{43} \approx 0.698$ ✓
 $r^2(0.698) = 3720.932 \therefore r = 60.999 \approx 61 \text{ km}$ ✓
 $r''(0.698) > 0 \therefore$ minimum ✓
 The minimum distance is 61 km. [7]

18. (a) $p(x) = \frac{1}{15}$ for $25 \leq x \leq 40$ and 0 otherwise ✓
 (b) 32.5 min ✓

$$P(X < 30) = \frac{1}{3}$$
 ✓
 (c)
$$P(X > 30) = \frac{2}{3}$$
 ✓
 (d) (i)
$$\left(\frac{2}{3}\right)^3 \times 3 = \frac{8}{9}$$
 ✓
 (ii) $E(X) = 32.5 \quad \text{Var}(X) = 18.75$ ✓
 $a^2(18.75) = 12.25$ ✓
 $a(32.5) + b = 26$ ✓
 $a = -0.808 \quad b = 52.27 \quad \text{or} \quad a = 0.808 \quad b = -0.27$ ✓
 [9]

19. (a) $f(t) = \frac{1}{20} e^{-\frac{1}{20}t}$ for $t > 0$ ✓✓
- (b) $F(20) = 0.6321$ ✓
- $$P(t \leq 40 | t \geq 25) = \frac{P(25 \leq t \leq 40)}{P(t \geq 25)}$$
- (c) $= \frac{0.151169}{0.2865047} = 0.5276$ ✓✓
- or
- $$P(t < 15) = 1 - e^{-0.75} = 0.5276$$
- (d) $E(t) = \int_0^{\infty} t \times f(t) dt = 20$
20 months is the mean time (or $k = 20 = \mu$) ✓
- (e) $-e^{-\frac{q}{20}} + 1 = 0.096$
 $\therefore q = 2.02$ therefore just over 2 months ✓ [7]
20. (a) $1 = \int_0^x \frac{1}{8} t^3 e^{-t} dt.$ ✓
- $x = 3.0539$ ✓
- (b) $\frac{dP}{dx} = \frac{1}{8} x^3 e^{-x} \mid x = 3$ ✓
- $= 1.2416$ ✓
- (c) $P(x) = \frac{x^4 e^{-x}}{32} + c$
- $P(3) - P(1) = 0.93119 - 0.011496 = 0.9197$
- $$\int_1^3 \frac{1}{8} x^3 e^{-x} dx = 0.9197$$
- or ✓✓
- (d) $\delta x = \frac{1}{365}$ ✓
- $\delta P = \frac{dP}{dx} \delta x$
- $$= 1.2416 \times \frac{1}{365} \approx 0.0034$$
- ✓✓ [9]

21. (a) $0.02275 \times 3000 = 68.25 \quad \therefore 68 \text{ people}$ ✓
 (b) (i) \$75 538.04 ✓
 (ii) $B \sim (3, 0.08) \quad \therefore P(X = 1) = 0.2031$ ✓✓
 (c) New mean = \$66 875 ✓
 Standard deviation = \$7687.50 ✓
 (d) $15 \times \frac{104}{366} = 4.26$ ✓
 Therefore 4 interns ✓ [8]
22. (a) 384 carp ✓
 (b) $385 - e^{0.04t} = 0$
 $t = 148.8 \text{ months}$ ✓
 In the 149th month
 (c) $B(t) = 10e^{0.02t}$ ✓
 $385 - e^{0.04t} = 10e^{0.02t}$ ✓
 $t = 136.2 \text{ months after the intro of bass (the 137th month)}$ ✓ [5]
23. (a) $np = 6$ Therefore the distribution cannot be determined. ✓
 (b) (i) Normal distribution ✓
 $\hat{p} = N(0.06, 0.013711)$ ✓✓
 (ii) 0.9999 ✓ [5]
24. (a) $f(x) = \ln(x + 1) + 2$ ✓✓
 (b) (i) Whisper: $I = 10^{13} \text{ W/m}^2$
 Conversation: $I = 10^{18} \text{ W/m}^2$ ✓
 The intensity of the conversation is 10^5 times more than the whisper. ✓
 (ii) The loudness is an additional 3.0103 dB ($10\log 2$) ✓ [5]