

No other items may be taken into the examination room. It is **your** responsibility to ensure that you do not have any unauthorised material if you have any unauthorised material with you, hand it to the supervisor **before** reading any further.

Important note to candidates

Special items: drawing instruments, templates, notes on two unfolded sheets of A4 paper, and up to three calculators approved for use in this examination.

Standard items: pens (blue/black preferred), pencils (including coloured), sharpener, correction fluid/tape, eraser, ruler, highlighters

To be provided by the candidate

Formula sheet (retained from Section One)

This Question/Answer booklet

To be provided by the supervisor

Materials required/recommended for this section

Working time: one hundred minutes
Reading time before commencing work: ten minutes

Your name

In words

<input type="text"/>							
----------------------	----------------------	----------------------	----------------------	----------------------	----------------------	----------------------	----------------------

Student number: In figures

Calculator-assumed

Section Two:

METHODS

UNIT 3

MATHEMATICS

SOLUTIONS

Question/Answer booklet

Semester One Examination, 2018

COLLEGE
ALL SAINTS'



Structure of this paper

Section	Number of questions available	Number of questions to be answered	Working time (minutes)	Marks available	Percentage of examination
Section One: Calculator-free	8	8	50	52	35
Section Two: Calculator-assumed	13	13	100	98	65
Total					100

Instructions to candidates

1. The rules for the conduct of examinations are detailed in the school handbook. Sitting this examination implies that you agree to abide by these rules.
2. Write your answers in this Question/Answer booklet.
3. You must be careful to confine your response to the specific question asked and to follow any instructions that are specified to a particular question.
4. Supplementary pages for the use of planning/continuing your answer to a question have been provided at the end of this Question/Answer booklet. If you use these pages to continue an answer, indicate at the original answer where the answer is continued, i.e. give the page number.
5. Show all your working clearly. Your working should be in sufficient detail to allow your answers to be checked readily and for marks to be awarded for reasoning. Incorrect answers given without supporting reasoning cannot be allocated any marks. For any question or part question worth more than two marks, valid working or justification is required to receive full marks. If you repeat any question, ensure that you cancel the answer you do not wish to have marked.
6. It is recommended that you do not use pencil, except in diagrams.
7. The Formula sheet is not to be handed in with your Question/Answer booklet.

(2 marks)

(2 marks)

(2 marks)

(6 marks)

This section has **thirteen (13)** questions. Answer **all** questions. Write your answers in the spaces provided.

Section Two: Calculator-assumed

65% (98 Marks)

Working time: 100 minutes.

(a) Describe, with parameters, a suitable probability distribution to model the number of trout in a day's catch. (2 marks)

Solution	Binomial, with $n = 10$ and $p = 0.78$
Specific behaviours	binomial
Probability	parameters

(b) Determine the probability that there are fewer trout than fish of other species in a day's catch. (2 marks)

Solution	$P(X \leq 4) = 0.0104$
Specific behaviours	writes $P(X \leq 4)$ or $P(X < 5)$
Probability	probability, to at least 3dp

(c) Calculate the probability that over two consecutive days, a total of exactly 19 trout are caught. (2 marks)

Solution	$X \sim B(20, 0.78)$
Specific behaviours	defines new distribution
Probability	probability

Question 10**(8 marks)**

The population of a city can be modelled by $P = P_0 e^{kt}$, where P is the number of people living in the city, in millions, t years after the start of the year 2000.

At the start of years 2009 and 2013 there were 1945 000 and 2061 000 people respectively living in the city.

- (a) Determine the value of the constant k .

(2 marks)

Solution
$1.945 = 2.061 e^{4k}$
$k = 0.01448$
Specific behaviours
✓ equation
✓ value of k to at least 3sf

- (b) Determine the value of the constant P_0 .

(2 marks)

Solution
$1.945 = P_0 e^{0.01448 \cdot 9}$
$P_0 = 1.707$
Specific behaviours
✓ equation
✓ value of P_0 (in millions)

- (c) Use the model to determine during which year the population of the city will first exceed 2 500 000.

(2 marks)

Solution
$2.5 = 1.783 e^{0.01448 t}$
$t = 26.3 \Rightarrow$ during 2026
Specific behaviours
✓ value of t
✓ correct year


- (d) Determine the rate of change of the city's population at the start of 2013.

(2 marks)

Solution
$\frac{dP}{dt} = 0.01448 \times 2061 \text{ 000}$
29 800 people per year
Specific behaviours
✓ substitutes into rate of change
✓ correct rate with units

© 2018 WA Exam Papers. All Saints' College has a non-exclusive licence to copy and communicate this document for non-commercial, educational use within the school. No other copying, communication or use is permitted without the express written permission of WA Exam Papers. SN001-115-4.

Question 11
The graph of $f(x) = \frac{x+1}{6x+1}$ is shown below.



(6 marks)

(a)	Complete the missing values in the table below.
Solution (a)	See table
Two rectangles are also shown on the graph, with dotted lines, and they both have corners just touching the curve. The smaller is called the inscribed rectangle.	circumscribed rectangle.
See table	missing values
Specfic behaviours	Specfic behaviours

(b) Complete the table of areas below and use the values to determine a lower and upper bound for $\int_0^2 f(x) dx$.
(4 marks)

x interval	0 to 0.5	0.5 to 1	1 to 1.5	1.5 to 2	Area of circumscribed rectangle
	$\frac{1}{2}$	$\frac{3}{4}$	$\frac{7}{4}$	2	$\frac{13}{6}$
	$\frac{3}{4}$	$\frac{7}{4}$	2	$\frac{13}{6}$	Upper bound: $U = \frac{3}{4} + \frac{4}{7} + 2 + \frac{13}{6} = \frac{29}{4} = 7.25$

$$\text{Lower bound: } L = \frac{1}{2} + \frac{3}{4} + 2 + \frac{6}{7} \approx 5.583$$

Solution	Specfic behaviours
$\int_0^2 f(x) dx$	inscribed areas
$\int_0^2 f(x) dx$	circumscribed areas

(c) Explain how the bounds you found in (b) would change if a larger number of smaller intervals were used.
(1 mark)

Solution	The lower bound would increase and the upper bound decrease.
Specfic behaviours	describes changes to both bounds
Specfic behaviours	see inside page

Question 12

(8 marks)

A fairground shooting range charges customers \$6 to take 9 shots at a target. A prize of \$20 is awarded if a customer hits the target three times and a prize of \$40 is awarded if a customer hits the target more than three times. Otherwise no prize money is paid.

Assume that successive shots made by a customer are independent and hit the target with the probability 0.15.

- (a) Calculate the probability that the next customer to buy 9 shots wins

- (i) a prize of \$20.

Solution	
X	$B(9, 0.15)$
$P(X=3)=0.1069$	
Specific behaviours	
✓ defines distribution	
✓ calculates probability	

(2 marks)

- (ii) a prize of \$40.

Solution	
$P(X \geq 4)=0.0339$	
Specific behaviours	
✓ calculates probability	

(1 mark)

- (b) Calculate the expected profit made by the shooting range from the next 30 customers who pay for 9 shots at the target.

(3 marks)

Solution	
Let Y be the profit per customer	
$P(Y=6)=0.8592$	$P(Y=-14)=0.1069$
$P(Y=-34)=0.0339$	
$E(Y)=2.504$	
Expected profit = $30 \times 2.504 = \$75.13$	
Specific behaviours	
✓ indicates probability distribution	
✓ calculates expected value for one customer	
✓ calculates expected value	

- (c) Determine the probability that more than 6 out of the next 8 customers will not win a prize.

(2 marks)

Solution	
X	$B(8, 0.8592)$
$P(X \geq 7)=0.6862$	
Specific behaviours	
✓ defines distribution	
✓ calculates probability	

Supplementary page

Question number: _____

(Reproducing instruction supplied for guidance only)

Specific behaviours

- ✓ total area
- ✓ evaluates second integral
- ✓ writes second integral
- ✓ evaluates first integral
- ✓ writes first integral

Solution

$$\text{Area} = 7.905 + 10.166 = 18.071 \text{ sq units}$$

$$\int_{15.694}^{6.866} f(x) - h(x) dx = 10.166$$


$$g(x) = h(x) \text{ when } x = 15.694$$

$$\int_{6.866}^0 g(x) - h(x) dx = 7.905$$

$$f(x) = g(x) \text{ when } x = 6.866$$

- (b) Working to three decimal places throughout, determine the area of the region enclosed by all three functions. (5 marks)

- (a) One of the functions is shown on the graph above. Add the graphs of the other two functions. (3 marks)



Three functions are defined by $f(x) = 9e^{-0.07x}$, $g(x) = 3e^{0.09x}$ and $h(x) = 3$.

(8 marks)

Question number: _____

Supplementary page

Question 14

(7 marks)

A fuel storage tank, initially containing 3750 L, is being filled at a rate given by

$$\frac{dV}{dt} = \frac{t^2(240 - 3t)}{500}, \quad 0 \leq t \leq 80$$

where V is the volume of fuel in the tank in litres and t is the time in minutes since filling began.
The tank will be completely full after 80 minutes.

- (a) Calculate the volume of fuel in the tank after half-an-hour. (3 marks)

Solution
$\Delta V = \int_0^{30} V'(t) dt$ <i>3105</i>
$V = 3750 + 3105 = 6855 \text{ L}$
Specific behaviours
✓ indicates use of integral or rate of change ✓ calculates increase ✓ states volume

- (b) Determine the time taken for the tank to fill to three-quarters of its maximum capacity. (4 marks)

Solution
$V = 3750 + \int_0^{80} V'(t) dt$ <i>3750 + 20480 = 24230</i>
$V(T) = \int_0^T V'(t) dt = \frac{4T^3}{25} - \frac{3T^4}{2000} + 3750$
$\frac{4T^3}{25} - \frac{3T^4}{2000} + 3750 = \frac{3 \times 24230}{4}$
$T = 58.4 \text{ minutes}$
Specific behaviours
✓ calculates V_{MAX} ✓ indicates $V(T)$ ✓ indicates equation ✓ solves for time

Question 15

(8 marks)

$P(X=x)$	x	3	4	0.15	a	b	0.2	7

(a) Determine the values of the constants a and b . (3 marks)

The discrete random variable X has a mean of 5.28 and the following probability distribution.

$a+b+0.55=1$
$4a+5b+3.05=5.28$
$a=0.02, b=0.43$

(b) Determine $P(X < 4 | X < 7)$. (2 marks)

(a) Determine the values of the constants a and b . (3 marks)


$a+b+0.55=1$
$4a+5b+3.05=5.28$
$a=0.02, b=0.43$

(b) Determine $P(X < 4 | X < 7)$. (2 marks)Another function, F , is defined as $F(x) = \int_0^x f(t) dt$ over the domain $0 \leq x \leq 16$.

Question 21

(8 marks)

The graph of $y=f(t)$ is shown below, where $f(t)=5 \sin \frac{4t}{\pi}$.




(a) Determine the exact area between the horizontal axis and the curve for $0 \leq t \leq 4$. (2 marks)

(b) Determine the value(s) of x for which $F(x)$ has a maximum and state the value of $F(x)$ at this location. (2 marks)

Solution

$$x=4, x=12, F(4)=F(12)=\frac{40}{\pi}$$

(c) Sketch the graph of $y=F(x)$ on the axes below. (3 marks)



(a) Determine the values of the constants a and b . (3 marks)

Solution

$$P(X < 4 | X < 7) = \frac{0.15}{0.15} = 0.1875 = \frac{3}{16}$$

Specific behaviours

Determine

- ✓ denominator
- ✓ numerator and expresses as decimal or fraction

(c) Determine $\text{Var}(X)$. (3 marks)

Solution

$$\text{Var}(X) = 1.5416 \text{ (using CAS)}$$

Specific behaviours

Determine

- ✓ correct variance

(b) **Solution**

$$E(100 - 15X) = 100 - 15 \times 5.28 = 20.8$$

Specific behaviours

Determine

- ✓ correct mean

(1 mark)

(c) **Solution**

$$\text{Var}(12 - 5X) = (-5)^2 \times 1.5416 = 38.54$$

Specific behaviours

Determine

- ✓ correct variance

(1 mark)

(d) **Solution**

$$\text{Var}(12 - 5X)$$

(1 mark)

(e) **Solution**

$$\text{Var}(12 - 5X)$$

Specific behaviours

Determine

- ✓ correct variance

(1 mark)

(f) **Solution**

$$\text{Var}(12 - 5X)$$

Specific behaviours

Determine

- ✓ correct variance

(1 mark)

(g) **Solution**

$$\text{Var}(12 - 5X)$$

Specific behaviours

Determine

- ✓ correct variance

(1 mark)

(h) **Solution**

$$\text{Var}(12 - 5X)$$

Specific behaviours

Determine

- ✓ correct variance

(1 mark)

(i) **Solution**

$$\text{Var}(12 - 5X)$$

Specific behaviours

Determine

- ✓ correct variance

(1 mark)

(j) **Solution**

$$\text{Var}(12 - 5X)$$

Specific behaviours

Determine

- ✓ correct variance

(1 mark)

(k) **Solution**

$$\text{Var}(12 - 5X)$$

Specific behaviours

Determine

- ✓ correct variance

(1 mark)

(l) **Solution**

$$\text{Var}(12 - 5X)$$

Specific behaviours

Determine

- ✓ correct variance

(1 mark)

(m) **Solution**

$$\text{Var}(12 - 5X)$$

Specific behaviours

Determine

- ✓ correct variance

(1 mark)

(n) **Solution**

$$\text{Var}(12 - 5X)$$

Specific behaviours

Determine

- ✓ correct variance

(1 mark)

(o) **Solution**

$$\text{Var}(12 - 5X)$$

Specific behaviours

Determine

- ✓ correct variance

(1 mark)

(p) **Solution**

$$\text{Var}(12 - 5X)$$

Specific behaviours

Determine

- ✓ correct variance

(1 mark)

(q) **Solution**

$$\text{Var}(12 - 5X)$$

Specific behaviours

Determine

- ✓ correct variance

(1 mark)

Question 16

A particle starts from rest at O and travels in a straight line.

Its velocity $v \text{ ms}^{-1}$, at time $t \text{ s}$, is given by $v=8t-t^2$ for $0 \leq t \leq 3$ and $v=135t^{-2}$ for $t > 3$.

- (a) Determine the initial acceleration of the particle.

(2 marks)

Solution

$$a = \frac{dv}{dt} = 8 - 2t \Rightarrow a|_0 = 8 \text{ ms}^{-2}$$

Specific behaviours

- ✓ differentiates velocity
- ✓ acceleration

- (b) Calculate the change in displacement of the particle during the first three seconds.

(2 marks)

Solution

$$x = \int_0^3 8t - t^2 dt = 27 \text{ m}$$

Specific behaviours

- ✓ integrates velocity
- ✓ change in displacement

- (c) Determine, in terms of t , an expression for the displacement, $x \text{ m}$, of the particle from O for $t > 3$.

(2 marks)

Solution

$$x = \int \frac{135}{t^2} dt = \frac{-135}{t} + c$$

$$x|_3 = 27 = \frac{-135}{3} + c \Rightarrow c = 72$$

$$x = \frac{-135}{t} + 72$$

Specific behaviours

- ✓ integrates velocity
- ✓ evaluates c

- (d) Determine the distance of the particle from O when its acceleration is -1.25 ms^{-2} .

(3 marks)

Solution

$$a = \frac{-270}{t^3}$$

$$\frac{-270}{t^3} = -1.25 \Rightarrow t = 6$$

$$x|_6 = 49.5 \Rightarrow \text{Distance from } O = 49.5 \text{ m}$$

Specific behaviours

- ✓ acceleration for $t > 3$
- ✓ solves for time
- ✓ calculates distance

Question 20

The discrete random variable X is defined by

$$P(X=x) = \begin{cases} \frac{2k}{e^{1-x}} & x=0,1 \\ 0 & \text{elsewhere.} \end{cases}$$

- (a) Show that $k = \frac{e}{2+2e}$.

(3 marks)

Solution

$$\frac{2k}{e} + \frac{2k}{1} = 1$$

$$k \left(\frac{2+2e}{e} \right) = 1$$

$$k = \frac{e}{2+2e}$$

Specific behaviours

- ✓ indicates $P(X=0)$ and $P(X=1)$
- ✓ sums probabilities to 1

- (b) Determine, in simplest form, the exact mean and standard deviation of X .

(3 marks)

Solution

NB Bernoulli distribution.

$$E(X) = 2k = \frac{e}{1+e}$$

$$\text{Var}(X) = \frac{2k}{e} \times 2k = \frac{2^2 k^2}{e}$$

$$SD = \sqrt{\frac{2^2 k^2}{e}} = \frac{2}{\sqrt{e}} \left(\frac{e}{2+2e} \right) = \frac{\sqrt{e}}{1+e}$$

Specific behaviours

- ✓ simplified $E(X)$
- ✓ correct expression for variance

(4 marks)

(a) Determine the values of n and p .

The random variable X is the number of faulty components in the sample. The mean and standard deviation of X are 30.6 and 5.1 respectively.

A random sample of n components are selected at random from a factory production line. The proportion of components that are defective is p and the probability that a component is defective is independent of the condition of any other component.

(7 marks)

Question 17

Show that when the train moves at a steady speed of $x \text{ km/h}$, where $x > 0$, the total cost per kilometre, C , is given by

$$C = \frac{9x}{t} + \frac{144}{x}.$$

(a) When the train moves at a speed of 16 km/h , Other costs amount to \$ 144 per hour, regardless of speed.

The hourly cost of fuel to run a train is proportional to the square of its speed and is \$ 36 per hour when the train moves at a speed of 16 km/h . Other costs amount to \$ 144 per hour, regardless of speed.

METHODS UNIT 3

14

CALCULATOR-ASSUMED

11

CALCULATOR-ASSUMED

Solution	
$X \sim B(n, p)$	
$np(1-p) = 5.1^2$	
$n=204, p=0.15$	
After changes are made to the manufacturing process, the proportion of defective components is now 3 %. Determine the smallest sample size required to ensure that the probability that the sample contains at least one defective component is at least 0.95.	
(b)	

(3 marks)

Use calculus to determine the minimum cost for the train to travel 240 km, assuming that the train travels at a constant speed for the entire journey.

(b) Use calculus to determine the minimum cost for the train to travel 240 km, assuming that the train travels at a constant speed for the entire journey.

Solution	
$X \sim B(n, 0.03)$	
$P(X \geq 1) \leq 0.951 - P(X=0) \leq 0.95$ <th data-kind="ghost"></th>	
$P(X=0) \leq 0.05$ <th data-kind="ghost"></th>	
$n > 98.4 \Leftrightarrow n \geq 99$	
Specific behaviours	
<ul style="list-style-type: none"> ✓ indicates required binomial probability ✓ uses $P(X=0)$ to create inequality ✓ solves and rounds to obtain n 	

After changes are made to the manufacturing process, the proportion of defective components is now 3 %. Determine the smallest sample size required to ensure that the probability that the sample contains at least one defective component is at least 0.95.

Use calculus to determine the minimum cost for the train to travel 240 km, assuming that the train travels at a constant speed for the entire journey.

Solution	
$\frac{dC}{dx} = 0 \Leftrightarrow x = 32(x > 0)$ <th data-kind="ghost"></th>	
$C = \frac{9x^2 - 9216}{64} + \frac{144}{x}$ <th data-kind="ghost"></th>	
Specific behaviours	
<ul style="list-style-type: none"> ✓ indicates first derivative ✓ indicates critical point ✓ indicates optimum cost per km ✓ indicates minimum cost 	

Fuel cost, f , is	
$f = kx^2 \Leftrightarrow k = \frac{16}{9}$	
$t = \frac{9x^2}{64} + 144$	
Total cost per hour, t , is	
$t = \frac{x}{64} + \frac{144}{x}$	
Cost per km, C , is	
$C = \frac{x}{t} = \frac{64}{9x^2} + \frac{144}{x}$	
Specific behaviours	
<ul style="list-style-type: none"> ✓ indicates expression for hourly cost of fuel ✓ indicates expression for total cost per hour ✓ indicates derivation of cost per km 	

Question 19
CALCULATOR-ASSUMED
CALCULATOR-ASSUMED
METHODS UNIT 3

Use calculus to determine the minimum cost for the train to travel 240 km, assuming that the train travels at a constant speed for the entire journey.

Use calculus to determine the minimum cost for the train to travel 240 km, assuming that the train travels at a constant speed for the entire journey.

Question 18

(11 marks)

The air pressure, $P(h)$ in kPa, experienced by a weather balloon varies with its height above sea level h km and is given by

$$P(h) = 101.5 e^{-0.135h}, 0 \leq h \leq 20.$$

- (a) Determine $\frac{dP}{dh}$ when the height of the balloon is 5.5 km.

(2 marks)

Solution
$\frac{dP}{dh} = -0.135 \times 101.5 e^{-0.135(5.5)}$ $\textcolor{red}{\dot{=}} -6.52 \text{ kPa/km}$
Specific behaviours
✓ uses derivative ✓ correct rate of change

- (b) What is the meaning of your answer to (a).

(1 mark)

Solution
The rate of change of pressure with respect to height when the height is 5.5 km.
Specific behaviours

✓ meaning (must include wrt h and refer to height)

The height of the balloon above sea level varies with time t minutes and is given by

$$h(t) = \frac{t^2(120-t)}{12800}, 0 \leq t \leq 80.$$

- (c) Determine the air pressure experienced by the balloon when $t = 48$.

(2 marks)

Solution
$h(48) = 12.96 \text{ km}$
$P(12.96) = 17.65 \text{ kPa}$
Specific behaviours

✓ determines height
✓ determines pressure

- (d) Determine $\frac{dh}{dt}$ when the height of the balloon is 7.04 km.

(3 marks)

Solution
$h(t) = 7.04 \Rightarrow t = 32$
$\frac{dh}{dt} = \frac{240t - 3t^2}{12800}$ $\textcolor{red}{\dot{=}} \frac{240(32) - 3(32)^2}{12800} = \frac{9}{25} = 0.36 \text{ km/m}$
Specific behaviours

✓ determines time
✓ indicates derivative
✓ determines rate of change

- (e) Determine $\frac{dP}{dt}$ when the height of the balloon is 7.04 km.

(3 marks)

Solution
$\frac{dP}{dh} = -0.135 \times 101.5 e^{-0.135(7.04)} \textcolor{red}{\dot{=}} -5.297$
$\frac{dP}{dt} = \frac{dP}{dh} \times \frac{dh}{dt} \textcolor{red}{\dot{=}} -5.297 \times 0.36 \textcolor{red}{\dot{=}} -1.91 \text{ kPa/m}$
Specific behaviours

✓ rate of change of P wrt h
✓ indicates use of chain rule
✓ correct rate of change