

Copyright for test papers and marking guides remains with *West Australian Test Papers*.  
Test papers may only be reproduced within the purchasing school according to the advertised Conditions of Sale.  
Test papers should be withdrawn after use and stored securely in the school until Friday, 1 December 2017.



## **SEMESTER TWO**

### **MATHEMATICS METHODS UNITS 1 and 2**

**2017**

## **SOLUTIONS**

**Calculator-free Solutions**

1. Origin is at (2, 8)  
 Radius  $= \sqrt{(2-6)^2 + (8-8)^2}$   
 $= 4$  ✓  
 $(x-2)^2 + (y-8)^2 = 16$  ✓ [2]

2. (a)  $\sin 2x = \frac{\sqrt{3}}{2}$  ✓  
 $\therefore 2x = \frac{\pi}{3}, \frac{2\pi}{3}, -\frac{4\pi}{3}, -\frac{5\pi}{3}$  ✓  
 $\therefore x = \frac{\pi}{6}, \frac{\pi}{3}, -\frac{2\pi}{3}, -\frac{5\pi}{6}$  ✓  
 (b)  $\cos x$  cannot be more than 1. ✓ [4]

3. (a)  $\frac{dy}{dx} = \frac{9}{5}kx^2$  ✓✓  
 (b)  $y = 3 - \frac{3x^2}{2} + \frac{x^{-\frac{1}{2}}}{2}$  ✓  
 $\frac{dy}{dx} = -3x - \frac{1}{4x^{\frac{3}{2}}}$  ✓✓  
 (c)  $y = 4x^3 - 9x$  ✓  
 $\frac{dy}{dx} = 12x^2 - 9$  ✓ [7]

4.

$y = -2^{-x} - 1$	$y = \left(\frac{1}{2}\right)^x$	$y = 2^x + 3$	$y = 2^x - 1$ ✓✓✓✓ [4]
D	A	B	C

5. (a)

(i)  $(2^4 \times 3^4)^{\frac{1}{4}}$

(ii)  $\frac{(ab^2c^{-3})^2 \sqrt{a^4b^{-2}c^6}}{a^3b^2c}$  ✓  
 $= \frac{ab}{c^4}$  ✓✓  
 $\frac{24^{\frac{2}{3}} \cdot \sqrt{18} \times 3^{\frac{1}{3}}}{(2^3 \cdot 3)^{\frac{2}{3}} \cdot 3 \cdot 2^{\frac{1}{2}} \times 3^{\frac{1}{3}}}$   
 (iii)  $\sqrt[6]{8} \cdot (27)^{\frac{1}{3}}$   
 $= \frac{(2^3 \cdot 3)^{\frac{2}{3}} \cdot 3 \cdot 2^{\frac{1}{2}} \times 3^{\frac{1}{3}}}{(2^3)^{\frac{2}{3}} \cdot 3}$  ✓✓  
 $= 2^2 \times 3$  ✓  
 $= 12$



(b) (i)  $\frac{2^{4x-4}}{2^{-4}} = 2^{-3}$  ✓  
 $2^{4x} = 2^{-3}$  ✓  
 $4x = -3$   
 $x = -\frac{3}{4}$  ✓  
(ii)  $(2^x - 8)(2^x - 1) = 0$   
 $2^x = 8$  or  $2^x = 1$  ✓  
 $x = 3$  or  $x = 0$  ✓ [11]

6. (a) (i) 0.2 ✓  
(ii) 0.4 ✓  
(iii) 0.7 ✓  
(b)  ${}^5C_2 \times 2^3 \times 3^2 = 720$  ✓✓ [5]

7. (a) (i)  $\int x^2 + 1 dx = \frac{x^3}{3} + x + c$  ✓✓  
(ii)  $\int t^{-2} - 2t + \pi t^{-3} dt$  ✓  
 $= -\frac{1}{t} - t^2 - \frac{\pi}{2t^2} + c$  ✓✓  
(b)  $y = 3x + \frac{x^2}{2} - \frac{2x^5}{5} + c$  ✓  
 $2 = 3 + \frac{1}{2} - \frac{2}{5} + c$   
 $\therefore c = -\frac{11}{10}$  ✓  
 $y = -\frac{2x^5}{5} + \frac{x^2}{2} + 3x - \frac{11}{10}$  ✓ [8]

8. (a)  $\frac{64}{2}(4 - b) = 0$   
 $b = 4$  ✓  
(b)  $p(x) = \frac{x^4}{2} - 2x^3$   
 $p'(x) = 2x^3 - 6x^2 = 0$  ✓  
 $2x^2(x - 3) = 0$   
 $x = 0$  or  $x = 3$  ( $x = 0$  is not a minimum) ✓  
When  $x = 3$   $y = \frac{3^4}{2} - 2(3^3)$   
 $\left(3, -\frac{27}{2}\right)$  ✓  
(c) Stationary point at  $x = 0$ ,  $p'(0) = 0$   
 $p''(x) = 6x^2 - 12x = 0$  ✓  
 $6x(x - 2) = 0$   
 $x = 0, 2$   
 $\therefore x = 0$   $p'(x) = p''(x) = 0$  ✓ [6]  
Horizontal point of inflection  
Or sign table:

	$x < 0$	$x = 0$	$0 < x < 3$	$x = 3$	$x > 3$
--	---------	---------	-------------	---------	---------


$p(x)$	↓	↔	↓	↔	↑
$p'(x)$	-	0	-	0	+

9. (a)  $a_1 = 3$   
 $a_2 = 6$   
 $a_3 = 10$  ✓✓  
 (b)  $T_{n+2} = T_n + T_{n+1}$   $T_1 = 1$   $T_2 = 1$  ✓ [3]

**Calculator-assumed Solutions**

10. (a)  $x^2 = 4x^2 + 9 - 2(2x)(3)\cos Q$   
 $\frac{-3(x^2 + 3)}{-12x} = \cos Q$  ✓  
 $\cos Q = \frac{x^2 + 3}{4x}$  ✓  
 (b)  $\cos Q = 0.9125$  ✓  
 $Q = 24.1468^\circ$  ✓  
 $\text{Area} = \frac{1}{2}(3)(4.8)\sin 24.1468$   
 $= 2.945 \text{ units}^2$  ✓ [5]

11. (a)  $-\frac{b}{2a} = -\frac{2}{2a} = -\frac{1}{a}$   
 $y\left(-\frac{1}{a}\right) = a - \frac{1}{a}$  or  $\frac{a^2 - 1}{a}$   
 Turning point  $\left(-\frac{1}{a}, \frac{a^2 - 1}{a}\right)$  ✓✓  
 (b)  $b^2 - 4ac > 0$   
 $4 - 4a^2 > 0$  ✓  
 $a^2 < 1$   
 $-1 < a < 1$  ✓ [4]

12. (a) 

- (b) (i) 0.1 ✓  
 (ii) 0.2 ✓  
 (iii) 0.3 ✓  
 (iv) 0.9 ✓ [5]


13. (a)  $C(t) = \frac{t^3}{3} - \frac{t^2}{2} - 12t + 105$  ✓  
 $C(2) = 81.7 \text{ cents}$  ✓  
 (b)  $(t - 4)(t + 3) = 0$   
 $t = 4 \text{ weeks}$  ✓  
 $C(4) = 70.3 \text{ cents}$  ✓  
 (c) Week 6.8 therefore during week 7 ✓ [5]

14. (a)  $d = k\sqrt{h}$   
 $4665 = k\sqrt{1.7}$   
 $\therefore k = 3577.89 \approx 3578$  ✓  
 $d = 3577.89\sqrt{(1.76 + 85)}$   
 $d = 33\,326.3 \text{ m}$  ✓  
 $d \approx 33 \text{ km}$  ✓
- (b)  $d = 3577.89\sqrt{0.8h}$  ✓  
 $d = 3577.89(0.8944)\sqrt{h}$  ✓  
 $d$  decreases by 10.6%. ✓ [5]
15. (a) (i)  $T_{n+1} = T_n + \sqrt{2}$   $T_1 = 3 - \sqrt{2}$  ✓✓  
(ii)  $T_{n+1} = 3T_n$   $T_1 = -2$  ✓✓
- (b)  $T_n = 17 + (n-1)(3)$  ✓  
 $T_{25} = 17 + (24)(3)$   
 $= 89$  ✓ [6]
16. (a)
- | $h \rightarrow 0$ | Limit =   |
|-------------------|-----------|
| 0.1               | 3.05      |
| 0.01              | 3.005     |
| 0.0001            | 3.00005   |
| 0.000001          | 3.0000005 |
- $\therefore \lim_{h \rightarrow 0} \frac{f(3+h) - f(3)}{h} = 3$  ✓✓
- (b)  $f(x) = 3x^2$  ✓  
 $\therefore f'(x) = 6x$  ✓
- (c)  $4x + k = 0$  ✓  
 $4(3) + k = 0$  ✓  
 $\therefore k = -12$  ✓
- (d)  $y = \int 4x + 1 \, dx$   
 $\therefore y = 2x^2 + x + c$  ✓  
 $2(1)^2 + 1 + c = -2$   
 $c = -5$   
 $y = 2x^2 + x - 5$  ✓ [9]
17. (a)  $\frac{\sqrt{1.44 \times 10^6}}{(2 \times 10^{-2})^4}$   
 $= \frac{1.2 \times 10^3}{1.6 \times 10^{-7}}$  ✓  
 $= 0.75 \times 10^{10}$  ✓  
 $= 7.5 \times 10^9$  ✓
- (b) (i)  $1.5 \times 10^4 = 1200(r)^6$  ✓  
 $P_0 = 1200$  ✓  
 $r = 1.5234$  ✓
- (ii)  $P = 187500$  ✓  
(iii)  $t = 15.987$  ✓  
15 hours and 59 minutes. ✓ [8]





18. (a)



✓✓✓

(b)

A	B	C
(iii)	(ii)	(i)

✓✓✓ [6]

19. (a) (i)  $n = 13$   
 $T_{13} = 26 + (12)(2)$   
 $= 50$  seats
- (ii)  $n = 23$  Row W
- (iii)  $n = 24$   
Sum of seats = 1176

✓

✓

✓

✓

- (b)  $205 = 5(2a + 9d)$   
 $710 = 10(2a + 19d)$   
 $a = 7 \quad d = 3$   
 $S_{30} = 15(14 + 29(3))$   
 $= 1515$

✓

✓

✓

- (c)  $a = 254 \quad d = -3 \quad T_n = 176$   
 $176 = 254 + (n - 1)(-3)$   
 $n = 27$   
 $S_n = \frac{27}{2}(254 + 176)$   
 $= 5805$

✓

✓

✓ [10]

20. (a)  $V(x) = 3x(90 - 3x)\left(\frac{x}{3}\right)$   
 $= x^2(90 - 3x)$

✓

✓

- $= 90x^2 - 3x^3$
- (b)  $V'(x) = 180x - 9x^2$   
 $9x(20 - x) = 0$   
 $x = 0$  or  $x = 20$   
 $V''(x) = 180 - 18x \quad V''(20) < 0 \therefore$  Maximum  
Maximum volume is  $12000 \text{ m}^3$   
When length =  $60 \text{ m}$ , width =  $30 \text{ m}$  and depth =  $6.67 \text{ m}$

✓

✓

✓

✓ [6]

21. (a) (i)  $A_1 = \frac{3}{2}$   $A_2 = \frac{9}{2}$   $A_3 = \frac{27}{2}$  ✓
- (ii)  $A_{n+1} = 3 A_n$   $A_1 = \frac{3}{2}$  ✓✓
- (iii)  $S_{25} = 6.355 \times 10^{11}$  ✓✓
- (b) (i)  $T_n = 2 T_{n-1}$   $T_0 = 6$   
 $T_{12} = 24\,576$  users ✓✓
- (ii) After 18 months ✓✓  
 (Or during the 18<sup>th</sup> month)
- (iii)  $\frac{6000000}{1.15} = 5\,217\,391$  users ✓  
 During the 20<sup>th</sup> month ✓
- (c)  $S_\infty = \frac{1-r}{1-r}$   
 $= \frac{45}{100} \div \left(1 - \frac{1}{100}\right)$  ✓  
 $= \frac{45}{99}$   
 $= \frac{5}{11}$  ✓ [13]
22. (a)  $s = -\frac{t^3}{2} + t^2 + 20t$   $s(3) = 55.5 \text{ m}$  ✓
- (b)  $\frac{55.5}{3} = 18.5 \text{ m/s}$  ✓
- (c)  $v = -\frac{3t^2}{2} + 2t + 20$  ✓  
 $v(2) = 18 \text{ m/s}$  ✓
- (d)  $-\frac{3t^2}{2} + 2t + 20 = 0$  ✓  
 $t = 4.38 \text{ s}$  ✓  
 Max displacement:  $s(4.38) = 64.8 \text{ m}$  ✓
- (e)  $s(8) = -32$  ✓  
 Total distance =  $64.8 + 64.8 + 32 = 161.6 \text{ m}$  ✓ [9]
23. (a)  $g'(x) = 2x - 6$   
 $m = \frac{1}{3} \therefore 2x - 6 = \frac{1}{3}$  ✓  
 $x = \frac{19}{6} = 3\frac{1}{6}$  ✓  
 $y = \frac{1}{36} \left(\frac{19}{6}, \frac{1}{36}\right)$  ✓
- (b) (i)  $f'(x) = x^2 + 2ax + b$   
 $36 - 12a + b = 0$  ✓  
 $-72 + 36a - 6b = 0$  ✓  
 Solving simultaneously  $a = 4$   $b = 12$  ✓
- (ii)  $x^2 + 8x + 12 = 0$   
 $x = -6$  or  $x = -2$  ✓  
 Turning point  $\left(-2, -\frac{32}{3}\right)$  ✓  
 $f''(x) = 2x + 8$   $f''(-2) > 0$  Therefore Local Minimum ✓ [9]

## End of Questions