

Copyright for test papers and marking guides remains with *West Australian Test Papers*.
Test papers may only be reproduced within the purchasing school according to the advertised Conditions of Sale.
Test papers should be withdrawn after use and stored securely in the school until Friday 15th June 2018.

MATHEMATICS METHODS UNIT 3

Semester One

2018

SOLUTIONS

Calculator-free Solutions

1.. $f'(x) = 0$ when $\frac{1-x}{e^x} = 0$ ✓
 $\therefore x = 1$ ✓
 $f''(x) = \frac{(e^x)(-1) - e^x(1-x)}{e^{2x}}$ ✓
 Since $f''(1) < 0$, then maximum ✓

[4]

2. $A = \int_0^k e^{3-x} dx$ ✓
 $\therefore \left[-e^{3-x} \right]_0^k = (-e^{3-k}) - (-e^{3-0})$ ✓
 $= -e^{3-k} + e^3$ ✓
 If $A = e^3(1 - e^{-k}) = e^3$ then $e^{-k} = 0$, which is impossible. ✓
 If $A = e^3(1 - e^{-k}) = e^3 - 1$ ✓
 $\therefore k = 3$ ✓

[6]

3. (a) Stationary point occurs when $3x^2 - kx = 0$
 $\therefore M'(6) = 108 - 6k = 0$ ✓
 $\therefore k = 18$ ✓
 (b) $M(x) = x^3 - 9x^2 + c$ ✓
 Since $M(6) = 1 \rightarrow 216 - 324 + c = 1 \rightarrow c = 109$ ✓
 $\therefore M(x) = x^3 - 9x^2 + 109$ ✓
 \therefore Vertical intercept = 109 ✓

[6]

4. (a) $e^1 = 1 + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \frac{1}{4!}$ ✓
 $= 1 + 1 + \frac{1}{2} + \frac{1}{6} + \frac{1}{24} = 2\frac{17}{24}$ ✓
 (b) $\frac{d}{dx} \left(1 + \frac{x}{1} + \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24} + \dots \right)$
 $= 1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24} + \dots$ ✓
 $= 1 + \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \dots = e^x$ ✓

[4]

5. (a) $\Sigma f(x) = \frac{1}{3} + \frac{1}{3} + \frac{1}{3} = 1$ ✓
 and $0 \leq f(x) \leq 0$ for all x ✓
 \therefore PDF
 (b) $\Sigma f(x) = 1$ ✓
 and $0 \leq f(x) \leq 0$ for all x ✓
 \therefore PDF
 (c) Since $f(0) = -\frac{1}{3}$
 $f(x)$ is not greater than or equal to 0 for all x ✓
 \therefore Not PDF ✓ [6]
6. (a) Since PDF $\Sigma f(x) = 0.8 + a = 1 \rightarrow a = 0.2$ ✓
 (b) $P(X < 2 | X \leq 2) = \frac{P(X < 2)}{P(X \leq 2)}$ ✓

$$= \frac{0.6}{0.7} = \frac{6}{7}$$
 ✓
 (c) $E(X) = 0 + 0.1 + 0.2 + 0.3 + 0.8 = 1.4$ ✓✓
 (d) Variance = $(0 + 1 \times 0.1 + 4 \times 0.1 + 9 \times 0.1 + 16 \times 0.2) - 1.4^2$ ✓✓
 $= 0.1 + 0.4 + 0.9 + 3.2 - 1.96 = 2.64$ ✓ [8]
7. $E(Y) = 2E(X) + 3 = 8$
 $\therefore m = 2.5$ ✓
 $VAR(Y) = 4VAR(X) = 20$
 $\therefore v = 5$ ✓✓ [3]
8. (a) $e^x \sin(e^x)$ ✓✓
 (b) $\frac{d^2 y}{dx^2} = \frac{d}{dx} \left(\frac{dy}{dx} \right) = \frac{d}{dx} (e^x \sin e^x)$ ✓
 $= e^x \sin(e^x) + e^x \cos(e^x)(e^x)$ ✓✓ [5]
9. (a) $y = (\sin x)^{-1}$
 $\frac{dy}{dx} = -1(\sin x)^{-2}(\cos x)$ ✓

$$= \frac{-\cos x}{\sin^2 x}$$
 ✓
 (b) $\int \frac{5 \cos x}{1 - \cos^2 x} dx$
 $= 5 \int \frac{\cos x}{\sin^2 x} dx$ ✓
 $= -5 \int \frac{-\cos x}{\sin^2 x} dx$ ✓
 $= -\frac{5}{\sin x} + c$ ✓ [5]

10. (a) $y = (ex)(e^x)$

$$\therefore \frac{dy}{dx} = e(e^x) + (ex)(e^x)$$

✓

$$\therefore = e^{x+1}(x+1)$$

✓

(b) $y = \frac{\pi \sin x}{\cos x}$

$$\therefore \frac{dy}{dx} = \frac{(\cos x)(\pi \cos x) - (\pi \sin x)(-\sin x)}{\cos^2 x}$$

✓

$$= \frac{\pi(\cos^2 x + \sin^2 x)}{\cos^2 x}$$

✓

$$= \frac{\pi \times 1}{\cos^2 x} = \frac{\pi}{\cos^2 x}$$

✓

[5]

Calculator-assumed Solutions

11. (a) $K = \frac{1}{2} r^2 \sin \theta \rightarrow \frac{dK}{d\theta} = 8 \cos \theta$ when $r = 4$

✓

$$\delta K = \frac{dK}{d\theta} \times \delta \theta \rightarrow \delta K = 8 \cos \theta \times (0.05\pi)$$

✓

$$\therefore \delta K = 8 \cos \frac{\pi}{4} \times 0.05\pi \text{ when } \theta = \frac{\pi}{4}$$

$$\therefore \delta K = 0.889$$

✓

(b) $\Delta K = \frac{1}{2} 16 \left(\sin 0.3\pi - \sin \frac{\pi}{4} \right)$

$$\therefore \Delta K = 0.8153$$

✓

$$\therefore \text{Error} = \frac{0.8886 - 0.8153}{0.8886} \times 100 = 8.3\%$$

✓

[5]

12. $(-3, 0) \rightarrow -27a - 9b - 3c - 9 = 0$

✓

$$\frac{dy}{dx} = 3ax^2 - 2bx + c$$

✓

Stationary point at $x = -3 \rightarrow 0 = 27a + 6b + c$

✓

$$\frac{d^2y}{dx^2} = 6ax - 2b$$

✓

POI at $-\frac{5}{3} \rightarrow -10a - 2b = 0$

✓

$$\therefore \text{Using calc } a = 1, b = -5 \text{ and } c = 3$$

✓✓

[7]

13. (a) $f(x) = -4\cos \frac{x}{2} + c$ ✓
 $f\left(\frac{\pi}{2}\right) = -4\cos \frac{\pi}{4} + c = 4 - 2\sqrt{2} \rightarrow c = 4$ ✓
 $\therefore f(x) = -4\cos \frac{x}{2} + 4$ ✓
 (b) $-4\cos \frac{x}{2} + 4 = 4 - 2\sqrt{2}$
 $\therefore \cos \frac{x}{2} = \frac{\sqrt{2}}{2} \rightarrow \frac{x}{2} = \frac{\pi}{4}, \frac{7\pi}{4}, \dots$ ✓✓
 $\therefore \text{Next time is } x = \frac{7\pi}{2}$ ✓ [6]
14. (a) (i) $0 + 1 + (-5) + 4 = 0$ ✓✓
 (ii) $-3 + 1 + (-5) + 4 = -3$ ✓✓
 (iii) $1 + 5 + 4 = 10$ ✓✓
 $-5 + 4 + \int_2^4 7 \, dx = -1 + (28 - 14) = 13$
 (b) (i) ✓✓
 $2 \int_3^4 f(x) \, dx = 2(4) = 8$
 (ii) ✓✓ [10]
15. (a) (i) $X \sim B(30, 0.75)$ where X = the number graduated ✓
 $\therefore P(X = 25) = 0.1047$ ✓✓
 (ii) $P(X \geq 29 | X \geq 25) = \frac{0.00196}{0.2026}$ ✓✓
 $= 0.00967$ ✓
 (b) $Y \sim B(n, 0.75)$ where Y is the number who graduated out of n
 $\therefore P(Y \geq 10) \geq 0.99 \rightarrow n = 19$ using trial and error ✓✓✓

n	Y
18	0.981
19	0.991
20	0.996

- (c) (i) $\frac{\binom{5}{3}\binom{6}{0} + \binom{5}{2}\binom{6}{1} + \binom{5}{1}\binom{6}{2}}{\binom{11}{3}}$ ✓✓
 $= \frac{10 + 60 + 75}{165} = 0.87879$ ✓
 (ii) $P(S = n) = \begin{cases} \frac{5}{11}, & n = 1 \\ \frac{6}{11}, & n = 0 \end{cases}$ ✓✓✓ [15]

16. (a) $2\cos x \sin x + \cos x = 0$

$$\therefore \cos x = 0 \text{ or } \sin x = -\frac{1}{2}$$

$$\therefore x = \frac{\pi}{2}, \frac{3\pi}{2}, \frac{7\pi}{6}, \frac{11\pi}{6}$$

✓✓

(b) $\left(\frac{\pi}{2}, 0\right), \left(\frac{3\pi}{2}, 0\right), \left(\frac{7\pi}{6}, \frac{\sqrt{3}}{2}\right), \left(\frac{11\pi}{6}, -\frac{\sqrt{3}}{2}\right)$

✓✓

$$\int_{\frac{\pi}{2}}^{\frac{7\pi}{6}} (-\cos x - \sin 2x) dx$$

(c) $A = \frac{\pi}{2}$
 $= 2.25 \text{ units}^2$

✓

✓

[6]

17. (a) $\frac{x+1}{x-1} = 0 \rightarrow x = -1$

✓

(b) (i) $\frac{dy}{dx} = \frac{(x-1) - (x+1)}{(x-1)^2} = \frac{-2}{(x-1)^2}$

✓

(ii) True since $\frac{dy}{dx} \neq 0$ for any value of x .

✓

(c) (i) $\frac{d^2y}{dx^2} = \frac{4}{(x-1)^3}$

✓

(ii) True since $\frac{d^2y}{dx^2} \neq 0$ for any value of x .

✓

[5]

18. (a) $v(0) = 3$

✓

(b) Stationary when $2t^2 - 5t + 3 = 0$

✓

$$\therefore t = 1 \text{ and } 1.5$$

✓✓

(c) (i) $a = 4t - 5$

✓

(ii) v is a minimum when $a = 0 \rightarrow a = 1.25$

✓✓

(d) $x = \frac{2}{3}t^3 - \frac{5}{2}t^2 + 3t + c$

✓

Since $x(0) = 0 \rightarrow c = 0$

$$\therefore x(t) = \frac{2}{3}t^3 - \frac{5}{2}t^2 + 3t$$

✓

$$\therefore x(3) = 4.5$$

✓

$$\text{Distance} = \int_0^3 |v(t)| dt$$

(e) $= 4.58$

✓

✓

(f) Distance travelled \neq Displacement

✓

[13]

19. (a) $A = A_0 e^{0.05t}$ ✓
 $\therefore A = 0.6e^{0.05(10)} = 0.99 \text{ Ha}$ ✓✓
- (b) $5 = 0.6e^{0.05t}$ ✓
 $\therefore t = 42.4 \text{ hours}$ ✓✓ [6]
20. (a) (i) $f'(x) = -\sin x + \cos x$ ✓
(ii) $f''(x) = -\cos x - \sin x$ ✓
- (b) Maximum value occurs when $f'(x) = 0$
 $-\sin x + \cos x = 0 \rightarrow f(x) = \sqrt{2}$ when $x = \frac{\pi}{4}$ ✓✓
 $\therefore f''\left(\frac{\pi}{4}\right) < 0$ then maximum ✓✓
Since
- (c) POI occurs when $f''(x) = 0$ ✓
 $\therefore -\cos x - \sin x = 0 \rightarrow \tan x = -1$ ✓
 $x = \frac{3\pi}{4} \text{ or } \frac{7\pi}{4}$ ✓✓
 $\therefore \left(\frac{3\pi}{4}, 0\right) \text{ and } \left(\frac{7\pi}{4}, 0\right)$ ✓ [10]
21. (a) 10 L/sec ✓
- (b) $10 - \frac{t}{20} = 0 \rightarrow t = 200 \text{ secs}$ ✓
- (c) $F = \int_0^{200} \left(10 - \frac{t}{20}\right) dt$ ✓
 $= 1000 \text{ L}$ ✓ [4]
22. (a) $r' = e^{-t}$ ✓
 $\therefore r'(4) = e^{-4} = 0.018$ ✓
and $r'(5) = e^{-5} = 0.0067$ ✓
- (b) $r''(t) = -e^{-t}$ ✓
Since $-e^{-t} \neq 0$, then the fire is spreading at its greatest rate at the start of the fire,
Or the graph has greatest slope at $t = 0$. ✓
- (c) $A = \pi r^2 \rightarrow \frac{dA}{dt} = 2\pi r$ ✓
 $\therefore \frac{dA}{dt} = \frac{dA}{dr} \times \frac{dr}{dt} = 2\pi r \times e^{-t}$ ✓
When $t = 4$ $\frac{dA}{dt} = 2\pi(-e^{-4} + 4)(0.018) = 0.458 \text{ cm}^2/\text{sec}$ ✓✓ [9]