

SOLUTIONS

2019

**MATHEMATICS
METHODS
UNITS 1 & 2**

SEMESTER TWO




Copyright for test papers and marking guides remains with West Australian Test Papers.
Test papers may only be reproduced within the purchasing school according to the advertised Conditions of Sale.
Test papers should be withdrawn after use and stored securely in the school until 29th November 2019.

Calculator-free Solutions

1. (a) $2^{-2} \times a^{-6} \times b^2$ ✓
 $= \frac{b^2}{4a^6}$ ✓
 $\frac{3^{-3} \times x^9 \times y^{-6}}{9^{-2} \times x^4 \times y^{-4}}$ ✓
(b) $= \frac{81 \times x^5}{27y^2}$ ✓
 $= \frac{3x^5}{y^2}$ ✓ [5]

2. (a) $2^{2-2x} = 2^{3-3x}$ ✓
 $\therefore 2-2x = 3-3x \rightarrow x = 1$ ✓
(b) $3^{x^2+1} = 3^{x+3}$ ✓
 $\therefore x^2+1 = x+3$ ✓
 $\therefore x^2-x-2 = 0 \rightarrow (x-2)(x+1) = 0$ ✓
 $\therefore x = 2 \text{ or } x = -1$ ✓ [6]

3. (a) (i) $m = 4$ ✓
(ii) $f(-2) = 4a^{-2} = \frac{4}{a^2}$ ✓



4. (a) (i) $\frac{dy}{dx} = 3x^3 + 5x^4$ ✓✓
(ii) $f(x) = \frac{x}{3} + \frac{2x^2}{\pi}$
 $\therefore f'(x) = \frac{1}{3} + \frac{4}{\pi}x$ ✓✓ [5]

[6]

(a) $T_{n+1} = T_n + 10$ where $T_1 = 100$

(b) $T_n = 100 + (n-1)(10) = 10n + 90$

(c) $T_8 = 10(8) + 90 = 170 \text{ km}$

(d) $S_{12} = 6(200 + 110) = 6(310) = 1860 \text{ km}$

[5]

(a) $\frac{dy}{dx} = 2x - \pi x^2 + \frac{5}{3}x^2$

(b) $y = x^2 - \pi x^2 + \frac{5}{3}x^3 + c$

(c) $f(n) = \frac{3}{4}n^2 - \frac{3}{4}n$

(d) $f(n) = \frac{9}{4}n^3 - \frac{3}{2}n^2 + c$

[4]

(a) $x(t) = a(t-0.5)^2 + 2.5$

(b) $x(t) = a(t-0.5)^2 + 2.5$

(c) $x(2) \rightarrow 2 = a(0.25) + 2.5 \rightarrow a = -2$

(d) $x(t) = -2(t-0.5)^2 + 2.5$

[9]

(a) $y(2) = -8$

(b) $8y = (3t^2 - 10t) \times 2$

(c) $\frac{dy}{dt} = -2 \rightarrow y = -8t + c$


(d) $8y = (-7) \times 2 = -14$

(e) $y = -8t + 4$

$$\text{Av rate of change} = \frac{-18 - (-4)}{2} = -7$$


5. (a) (i) $y(1) = -4 \text{ and } y(3) = -18$

9. (a)



✓✓✓

(b)



✓✓✓

[6]

21. (a) $g'(x) = 4x^3 - 16x$

✓

$g'(1) = -12$

✓

$\therefore y = -12x + c$

✓

$(1, 9) \rightarrow 9 = -12(1) + c \rightarrow c = 21$

✓

$\therefore y = -12x + 21$

✓

(b) $(1.45, 3.61)$ and $(-3.45, 62.4)$

✓✓

[6]

$$\frac{(\sin a \cos 45^\circ + \sin 45^\circ \cos a)(\cos a \cos 45^\circ - \sin 45^\circ \sin a)}{(\sin a - \cos a)(\sin a + \cos a)}$$

✓

22. (a)

$$= \frac{\left(\frac{\sqrt{2}}{2} \sin a + \frac{\sqrt{2}}{2} \cos a\right)\left(\frac{\sqrt{2}}{2} \cos a - \frac{\sqrt{2}}{2} \sin a\right)}{(\sin a - \cos a)(\sin a + \cos a)}$$

✓


$$= \frac{\frac{\sqrt{2}}{2}(\sin a + \cos a) \frac{\sqrt{2}}{2}(\cos a - \sin a)}{(\sin a - \cos a)(\sin a + \cos a)}$$

✓

$$= \left(\frac{2}{4}\right)(-1) = -\frac{1}{2}$$

✓

(b)



✓

$$180^2 = 100^2 + x^2 - 2x(100)\cos 59^\circ$$

✓✓

$$\therefore x = 210 \text{ km}$$

✓

[8]

23.
$$\left(\sin x + \frac{1}{\sin x}\right)^3$$

✓✓

$$= \sin^3 x + 3(\sin^2 x)\left(\frac{1}{\sin x}\right) + 3(\sin x)\left(\frac{1}{\sin x}\right)^2 + \left(\frac{1}{\sin x}\right)^3$$

✓✓

$$= \frac{1}{\sin^3 x} (\sin^6 x + 3\sin^4 x + 3\sin^2 x + 1)$$

✓

[3]

14. (a) 10L ✓
 (b) $W'(t) = 0.8t^3 - 1.8t^2 - t$ ✓
 $\therefore 0.8t^3 - 1.8t^2 - t = 0$ when $t = 2.7$ ✓
 \therefore During the third minute. ✓
 (c) (i) $W(2.7) = 5.2$ L ✓
 (ii) $W(4) = 14.8$ L ✓ [6]

15. (a) (i) -15 m ✓
 (ii) $v(t) = t^2 - 2t - 4$ ✓✓
 (b) $v(3) = -1$ ✓
 \therefore Speed = 1 m/s ✓
 (c) At rest when $v(t) = 0$
 $\therefore t^2 - 2t - 4 = 0$ ✓
 $\therefore t = 3.2$ s ✓
 (d) $x(0) = -3$
 $x(3.2) = -15.12$
 $x(5) = -6.33$ ✓
 \therefore Distance travelled = $12.12 + 8.79 = 20.91$ m ✓ [10]

16. (a) $c = 5$ ✓
 (b) $(-2, -3) \rightarrow -3 = -8 + 4a - 2b + 5$
 $\therefore 4a - 2b = 0$ ✓
 $\frac{dy}{dx} = 3x^2 + 2ax + b = 0$ when $x = -2$
 $\therefore 4a - b = 12$ ✓
 $\therefore a = 6$ and $b = 12$ ✓✓
 (c) $y = x^3 + 6x^2 + 12x + 5$
 $\therefore x^3 + 6x^2 + 12x + 5 = 0$ when $x = -0.558$ ✓✓ [7]

17. (a) $\frac{dy}{dx} = 15x^4 - 30ax^2 + 15a^2$
 $y = \frac{15x^5}{5} - \frac{30ax^3}{3} + 15a^2x + c$
 $\therefore (0, 0) \rightarrow c = 0$ ✓✓
 $\therefore y = 3x^5 - 10ax^3 + 15a^2x$
 (b) $\frac{dy}{dx} = 15(x^2 - a)^2 = 0$ when stationary ✓
 $\therefore (x^2 - a)^2 = 0$
 $\therefore x^2 = a$ ✓
 Since two stationary points $a > 0$ ✓ [6]