


<div><p>PERTH MODERN SCHOOL</p><p>Exceptional schooling. Exceptional students.</p><p>Independent Public School</p></div>	<div>Year 12 Specialist</div> <div>TEST 2</div> <div>Monday 1 April 2019</div> <div>TIME: 45 minutes working</div> <div>Classpads allowed</div> <div>One page of notes</div> <div>45 marks 7 Questions</div>
---	--

Name: _____

Teacher: _____

Note: All part questions worth more than 2 marks require working to obtain full marks.



- From the diagram, z_1 is a solution to $z^4=k$ for complex k .
- i) Determine k .
- ii) Determine the other three roots and express in the form $a+bi$.

Q2 (2, 3 & 1 = 6 marks)

Let $f(x) = \sqrt{2x-1}$ and $g(x) = \frac{1}{x+5}$.

a) State the natural domain and range of $g(x)$.

b) Does $f \circ g(x)$ exist over the natural domain of g ? If it does not, determine the largest possible domain for the composite to exist.


c) Determine $f \circ f^{-1}(x)$

Q3 (2, 3 & 2 = 7 marks)

Given that $f(x) = 2x^2 - 12x + 19$, $x \leq 3$, determine the following.

a) $f^{-1}(x)$ and its domain.

b) Sketch on the axes below, $f(x)$ & $f^{-1}(x)$




c) On the sketch above show the precise points where $f(x) = f^{-1}(x)$

Q7 (5 marks)

Let $w = 1 + qi$ where q is a real constant. Let $p(z) = z^3 + bz^2 + cz + d$, where b, c & d are real constants. If $p(z) = 0$ for $z = w$ and all roots of $p(z) = 0$ satisfy $|z^3| = 8$, determine all possible values of q, b, c & d .


Now consider the particular triangle OAB with $OA = \begin{pmatrix} 3 \\ 2 \\ \sqrt{3} \end{pmatrix}$ and $OB = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$, where α is a positive constant, chosen so that triangle OAB is isosceles, with $|OB| = |OA|$. Show that $\alpha = 4$.

d) Use a vector method to show that OQ is perpendicular to AB .




Q4 (2 & 3 = 5 marks)
Consider the function $y = f(x)$ for the questions below.

a) Sketch the function $y = |f(x)|$ on the axes below.



b) Sketch the function $y = |f(x) - |x||$ on the axes below.



Q5 (3 & 4 = 7 marks)

- a) Two moving objects have the following position vectors and constant velocities at time, $t = 0$:

$$r_a = \begin{pmatrix} 9 \\ -8 \end{pmatrix} m \quad v_a = \begin{pmatrix} -2 \\ 7 \end{pmatrix} m/s$$

$$r_b = \begin{pmatrix} 11 \\ -3 \end{pmatrix} m \quad v_b = \begin{pmatrix} 5 \\ -3 \end{pmatrix} m/s$$

Determine the closest approach and the time that this will occur.


- b) Let the circle S have a radius 3 units and centre $(1, \beta)$, where β is a constant, and the line

$$r = \begin{pmatrix} -2 \\ 0 \end{pmatrix} + \lambda \begin{pmatrix} 3 \\ -5 \end{pmatrix} \text{ is tangential to this circle. Determine the value(s) of } \beta.$$

c)

Q6 (1, 1, 1, 3, 1 & 3 = 10 marks)

The diagram below shows a triangle with vertices with O, A & B . Let O be the origin, with vectors $OA = a$ and $OB = b$.



- a) Determine the following vectors in terms of a & b .

i) MA , where M is the midpoint of the line segment OA .

ii) BA

iii) AQ , where Q is the midpoint of the line segment AB .

Let N be the midpoint of the line segment OB .

- b) Use a vector method to prove that the quadrilateral $MNQA$ is a parallelogram.