

Probability

For any event A and its complement A'	$P(A) = 1 - P(A')$
$P(A \cup B) = P(A) + P(B) - P(A \cap B)$	$P(A B) = \frac{P(A \cap B)}{P(B)}$

Random variables and probability distributions		
Bernoulli: mean is the sample proportion \hat{p}	$\mu = p$	$\sigma^2 = p(1 - p)$
Binomial distribution: $P(X = x) = \binom{n}{x} p^x (1 - p)^{n-x}$	$\mu = np$	$\sigma^2 = np(1 - p)$
Discrete random variable: $P(X = x) = P(x)$	$\mu = E(x) = \sum xp(x)$	$\sigma^2 = \sum (x - \mu)^2 p(x)$
Continuous random variable: $P(a \leq X \leq b) = \int_a^b p(x) dx$		
Expected value: $E(x) = \int_{-\infty}^{\infty} xp(x) dx$		
Variance: $\sigma^2 = \int_{-\infty}^{\infty} (x - \mu)^2 p(x) dx$		

Sample proportions	
Mean: $E(\hat{p}) = p$	Standard deviation: $s = \sqrt{\frac{p(1 - p)}{n}}$
Margin of error: $E = z \sqrt{\frac{p(1 - p)}{n}}$	Confidence interval: $\hat{p} - z \sqrt{\frac{\hat{p}(1 - \hat{p})}{n}} \leq p \leq \hat{p} + z \sqrt{\frac{\hat{p}(1 - \hat{p})}{n}}$

Note: Any additional formulas identified by the examination panel as necessary will be included in the body of the particular question.



MATHEMATICS METHODS
ATAR COURSE
FORMULA SHEET

2017

Copyright

© School Curriculum and Standards Authority, 2016

This document – apart from any third party copyright material contained in it – may be freely copied, or communicated on an intranet, for non-commercial purposes in educational institutions, provided that it is not changed and that the School Curriculum and Standards Authority is acknowledged as the copyright owner, and that the Authority's moral rights are not infringed.

Copying or communication for any other purpose can be done only within the terms of the *Copyright Act 1968* or with prior written permission of the School Curriculum and Standards Authority. Copying or communication of any third party copyright material can be done only within the terms of the *Copyright Act 1968* or with permission of the copyright owners.

Any content in this document that has been derived from the Australian Curriculum may be used under the terms of the Creative Commons Attribution 4.0 International (CC BY) licence.

This document is valid for teaching and examining until 31 December 2017.

Differentiation and integration

$\frac{d}{dx} (x^n) = nx^{n-1}$	$\int x^n dx = \frac{x^{n+1}}{n+1} + c, \quad n \neq -1$
$\frac{d}{dx} (e^{ax-b}) = ae^{ax-b}$	$\int e^{ax} dx = \frac{1}{a} e^{ax} + c$
$\frac{d}{dx} (\ln x) = \frac{1}{x}$	$\int \frac{1}{x} dx = \ln x + c, \quad x > 0$
$\frac{d}{dx} (\ln f(x)) = \frac{f'(x)}{f(x)}$	$\int \frac{f'(x)}{f(x)} dx = \ln f(x) + c, \quad f(x) > 0$
$\frac{d}{dx} (\sin(ax-b)) = a \cos(ax-b)$	$\int \sin(ax-b) dx = -\frac{1}{a} \cos(ax-b) + c$
$\frac{d}{dx} (\cos(ax-b)) = -a \sin(ax-b)$	$\int \cos(ax-b) dx = \frac{1}{a} \sin(ax-b) + c$
Product rule	<p>If $y = uv$ then $\frac{d}{dx} (uv) = u \frac{dv}{dx} + \frac{du}{dx} v$</p> <p>If $y = f(x) g(x)$ then $y' = f'(x) g(x) + f(x) g'(x)$</p>
Quotient rule	<p>If $y = \frac{u}{v}$ then $\frac{d}{dx} \left(\frac{u}{v} \right) = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$</p> <p>If $y = \frac{f(x)}{g(x)}$ then $y' = \frac{f'(x) g(x) - f(x) g'(x)}{(g(x))^2}$</p>
Chain rule	<p>If $y = f(u)$ and $u = g(x)$ then $\frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx}$</p> <p>If $y = f(g(x))$ then $y' = f'(g(x)) g'(x)$</p>
Fundamental theorem	$\frac{d}{dx} \left(\int_a^x f(t) dt \right) = f(x)$ and $\int_a^b f(x) dx = f(b) - f(a)$
Incremental formula	$\delta y \approx \frac{dy}{dx} \times \delta x$
Exponential growth and decay	$\frac{dP}{dt} = kP \Leftrightarrow P = P_0 e^{kt}$

See next page

Mensuration

Parallelogram	$A = bh$
Triangle	$A = \frac{1}{2} bh$ or $A = \frac{1}{2} ab \sin C$
Trapezium	$A = \frac{1}{2} (a + b)h$
Circle	$A = \pi r^2$ and $C = 2\pi r = \pi d$

Prism	$V = Ah$, where A is the area of the cross section
Pyramid	$V = \frac{1}{3} Ah$, where A is the area of the cross section
Cylinder	$V = \pi r^2 h$ $TSA = 2\pi rh + \pi r^2 h$
Cone	$V = \frac{1}{3} \pi r^2 h$ $TSA = \pi rs + \pi r^2$, where s is the slant height
Sphere	$V = \frac{4}{3} \pi r^3$ $TSA = 4\pi r^2$

Trigonometry

$\sin^2 x + \cos^2 x = 1$	$\tan x = \frac{\sin x}{\cos x}$
---------------------------	----------------------------------

Logarithms

$x = \log_a b \Leftrightarrow a^x = b$	$a^{\log_a b} = b$ and $\log_a (a^b) = b$
$\log_a mn = \log_a m + \log_a n$	$\log_a \frac{m}{n} = \log_a m - \log_a n$
$\log_a (m^k) = k \log_a m$	$\log_e x = \ln x$

See next page