

| | |
|---|--|
|  <p>PERTH MODERN SCHOOL Exceptional schooling. Exceptional students. Independent Public School</p> | <p>Year 12 Specialist TEST 3 Monday 1 July 2019 TIME: 45 minutes working Classspads allowed One page of notes 42 marks 6 Questions</p> |
|---|--|

Name: MARION KEY

Teacher: _____

Note: All part questions worth more than 2 marks require working to obtain full marks.

Q1 (3 & 3 = 6 marks)

a) Solve for the following system of linear equations.

$$x + 2y - z = 3$$

$$2x + 3y + 2z = -1$$

$$3x + 7y - 2z = 6$$

$$\begin{bmatrix} 1 & 2 & -1 & : & 3 \\ 0 & 3 & 2 & : & -1 \\ 0 & 7 & -2 & : & 6 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & -1 \\ 0 & -4 & 3 \end{bmatrix} \begin{matrix} \\ 2R_1 - R_2 \\ 3R_1 - R_3 \end{matrix}$$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & -5 & 10 \end{bmatrix} \begin{matrix} \\ \\ R_2 + R_3 \end{matrix}$$

$$z = -2$$

$$y - 4(-2) = 7$$

$$y = -1$$

$$x + 2(-1) - (-2) = 3$$

$$x = 3$$

soln $(3, -1, -2)$

- ✓ shows working (Gaussian elimination)
- ✓ solves for one variable
- ✓ solves for the other two
- (if no working shown)

Q1 - continued

$$x + 2y - z = 3$$

b) Determine the values of m & p such that $2x + 3y + 2z = m$ such that the system has

$$3x + py - 2z = 6$$

- (i) Infinite solutions
(ii) No solutions

$$\begin{bmatrix} 1 & 2 & -1 & 3 \\ 2 & 3 & 2 & m \\ 3 & p & -2 & 6 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 2 & -1 & 3 \\ 0 & 1 & -4 & 6-m \\ 0 & 6-p & -1 & 3 \end{bmatrix} \begin{matrix} \\ 2R_1 - R_2 \\ 3R_1 - R_3 \end{matrix}$$

$$\begin{bmatrix} 1 & 2 & -1 & 3 \\ 0 & 1 & -4 & 6-m \\ 0 & 1-4(6-p) & 0 & 6-m-12 \end{bmatrix} \begin{matrix} \\ R_2' - 4R_3' \\ \end{matrix}$$

$$\begin{matrix} -23+4p \\ -6-m \end{matrix}$$

✓ determining a line with two zeros

✓ states values for infinite

✓ states values for no soln

i) Infinite $-23+4p=0$ AND $-6-m=0$
 $p = \frac{23}{4}$ AND $m = -6$

ii) No solns $p = \frac{23}{4}$ AND $m \neq -6$

Q6 (4 marks)

Given that x & y are functions of t , show that $\frac{d^2 y}{dx^2} = \frac{\frac{dx}{dt} \frac{d^2 y}{dt^2} - \frac{dy}{dt} \frac{d^2 x}{dt^2}}{\left(\frac{dx}{dt}\right)^3}$

$$\frac{dy}{dx} = \frac{dy}{dt} \frac{dt}{dx} = \frac{\dot{y}}{\dot{x}} \quad \checkmark$$

$$\frac{d^2 y}{dx^2} = \frac{d}{dt} \left(\frac{dy}{dx} \right) \frac{dt}{dx} \quad \checkmark$$


$$= \frac{\dot{x} \ddot{y} - \dot{y} \ddot{x}}{\dot{x}^2} \quad \checkmark$$

$$= \frac{x \ddot{y} - y \ddot{x}}{\dot{x}^3} \quad \checkmark$$

$f(t)$

Q5 (2 & 5 = 7 marks)

Consider two rides at a circus, one is a merry go round and the other is a train on a straight line. Two twins decide to each try one of the two rides. Jane sits on the merry go round with a constant angular speed of $\frac{d\theta}{dt} = 5$ radians/minute moving in a clockwise direction and radius 6 metres and Mia sits on a train moving at 3 metres/minute away from the merry go round. See the diagram below.



a) Determine the distance between Jane and Mia when $\theta = \frac{3}{2}\pi$ and the train is 30 metres from the centre of the merry go round.

✓ uses cosine rule ✓ starts, distance (No need)

$$D^2 = 6^2 + 30^2 - 2(6)(30)\cos\frac{3}{2}\pi$$

$$D = 33.41m$$

b) Determine the time rate of change of this distance at the point defined in (a) above. (metres/minute)

✓

$$D^2 = 6^2 + x^2 - 2(6)(x)\cos\theta$$

$$2D\dot{D} = 2x\dot{x} - 12\dot{x}\cos\theta + 12x\sin\theta\dot{\theta}$$

$$0 = -5$$

$$2(33.41)\dot{D} = 2(30)(3) - 12\cos\frac{3}{2}\pi(3) + 12(30)\sin\frac{3}{2}\pi(-5)$$

$$\dot{D} = -20.37 \text{ m/min}$$

Q2 (2 & 3 = 5 marks)

If $r = \begin{pmatrix} t^2 - 1 \\ 5 - t \end{pmatrix}$, determine

a) $\left| \frac{dr}{dt} \right|$

$$\dot{r} = \begin{pmatrix} 2t \\ -1 \end{pmatrix}$$

$$|\dot{r}| = \sqrt{4t^2 + 1}$$

✓ answer

b) $\frac{d|\dot{r}|}{dt}$

$$|\dot{r}| = \sqrt{(t^2 - 1)^2 + (5 - t)^2}$$

$$\frac{d|\dot{r}|}{dt} = \frac{1}{2} \left[(t^2 - 1)^2 + (5 - t)^2 \right]^{-\frac{1}{2}} (2(t^2 - 1)2t + 2(5 - t)(-1))$$

✓

Q3 (2, 3 & 3 = 8 marks)

An object is initially at the origin with initial speed of $\begin{pmatrix} 7 \\ 3 \end{pmatrix} \text{ m/s}$ and an acceleration given by

$$a = \begin{pmatrix} 3e^t \\ 2 - \sin t \end{pmatrix} \text{ m/s}^2 \text{ at time } t \text{ seconds.}$$

Obtain an expression for the:

a) Velocity at time t . ✓ integrates ✓

$$\dot{x} = \begin{pmatrix} 3e^t + 7 \\ 2t + \cos t + 3 \end{pmatrix} + \begin{pmatrix} 1 \\ 3 \end{pmatrix} = \begin{pmatrix} 7 \\ 3 \end{pmatrix} + \begin{pmatrix} 1 \\ 3 \end{pmatrix} + \begin{pmatrix} 3e^t \\ 2t + \cos t + 6 \end{pmatrix}$$

✓ solves for \dot{x}

b) Position vector r at time t . ✓

$$\dot{r} = \begin{pmatrix} 3e^t + 7 \\ 2t + \cos t + 3 \end{pmatrix} + \begin{pmatrix} 1 \\ 3 \end{pmatrix} = \begin{pmatrix} 8 \\ 6 \end{pmatrix} + \begin{pmatrix} 3e^t \\ 2t + \cos t + 6 \end{pmatrix}$$

$$r = \begin{pmatrix} 3e^t + 8t \\ t^2 + \sin t + 6t \end{pmatrix} + \begin{pmatrix} 8t \\ 6t \end{pmatrix} = \begin{pmatrix} 3e^t + 8t \\ t^2 + \sin t + 6t \end{pmatrix}$$

c) Is the velocity ever perpendicular to the acceleration? Explain and if necessary solve for t values (if any).


always grad then zero
will never be perpendicular

$$\dot{a} \cdot \dot{v} = 9e^{2t} + (2 - \sin t)(2t + \cos t) > 0$$

uses dot product
gives exp for grad in term of t
states that dot product > 0

Q4 (3, 3, 3 & 3 = 12 marks)

Consider a cannon ball that is projected from the top of a building with speed V at an angle θ to the surface of the roof. There is a constant cross wind of 3 metres per second acting against the ball and the acceleration due to gravity is 9.8 m/s^2 down as shown in the diagram below. (Note- let the origin be at the top of the building on the edge)



- a) Given that the acceleration is given by $\ddot{r} = \begin{pmatrix} 0 \\ -9.8 \end{pmatrix} \text{ m/s}^2$ show using vector calculus that the

velocity $\dot{r} = \begin{pmatrix} V \cos \theta - 3 \\ V \sin \theta - 9.8t \end{pmatrix} \text{ m/s}.$

$$\dot{r}(0) = \begin{pmatrix} V \cos \theta - 3 \\ V \sin \theta \end{pmatrix}$$

$$\dot{r} = \begin{pmatrix} 0 \\ -9.8t \end{pmatrix} + \underline{c} \quad \begin{array}{l} \checkmark \text{ integrates} \\ \checkmark \text{ solves for } \underline{c} \\ \checkmark \text{ states } \underline{c} \end{array}$$

$$\begin{pmatrix} V \cos \theta - 3 \\ V \sin \theta \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} + \underline{c}$$

$$\dot{r} = \begin{pmatrix} V \cos \theta - 3 \\ V \sin \theta - 9.8t \end{pmatrix}$$

- b) Determine the cartesian equation of the path of the cannon ball in terms of V & θ . Show your working.

$$\underline{r} = \begin{bmatrix} (V \cos \theta - 3)t \\ V \sin \theta - 4.9t^2 \end{bmatrix} + \underline{c}$$

$$\underline{r}(0) = 0 \quad \therefore \underline{c} = 0$$

\checkmark states $\underline{r}(0) = 0$ or $\underline{c} = 0$

$$x = (V \cos \theta - 3)t \quad t = \frac{x}{V \cos \theta - 3}$$

\checkmark obtains exp for time in terms of x

$$y = V \sin \theta - 4.9t^2$$

\checkmark obtains cartesian eqn

$$y = \frac{x V \sin \theta}{V \cos \theta - 3} - \frac{4.9 x^2}{(V \cos \theta - 3)^2}$$

(-1 if origin changed)

Question states origin at top of building

Q4 continued

- c) Given that a point on the cartesian path has been measured as (7.4, 1.1) metres and the initial speed V of the ball from the cannon is 12 m/s , determine the initial angle θ of the ball when projected into the air.

$$1.1 = \frac{7.4(12) \sin \theta}{12 \cos \theta - 3} - \frac{4.9(7.4)^2}{(12 \cos \theta - 3)^2} \quad | 0 < \theta < \frac{\pi}{2}$$

$$\theta = 30.1^\circ \text{ or } 54.7^\circ$$

$$(0.524^R) \quad (0.984^R)$$

\checkmark subs known x & y values into cartesian

\checkmark solves for θ

\checkmark states 2 solutions (acute)

- d) If $V = 25 \text{ m/s}$ and $\theta = 45^\circ$ and a cross wind of 3 m/s as in the diagram on last page, determine how far from the foot of the building that the cannon ball lands on the road.

$$-50 = 25t \sin 45^\circ - 4.9t^2$$

\checkmark sets up eqn for t

$$t = -1.86, 5.47 \text{ s}$$

\checkmark subs t value into x

$$t \geq 0 \quad \therefore t = 5.47 \text{ s}$$

\checkmark states distance (No need for units)

$$x = (25 \cos 45^\circ - 3) 5.47$$

$$= 80.286 \text{ m}$$