

# MATHEMATICS: UNITS 3A AND 3B FORMULA SHEET 2012



School Curriculum and Standards Authority  
Government of Western Australia



Copyright  
© School Curriculum and Standards Authority, 2012  
This document—apart from any third party copyright material contained in it—may be freely copied, or communicated on an intranet, for non-commercial purposes by educational institutions, provided that it is not changed in any way and that the School Curriculum and Standards Authority is acknowledged as the copyright owner.  
Copying or communication for any other purpose can be done only within the terms of the Copyright Act or by permission of the Authority.  
Copying or communication of any third party copyright material contained in this document can be done only within the terms of the Copyright Act or by permission of the copyright owners.  
This document is valid for teaching and examining until 31 December 2012.

Mathematics 3A and 3B Formula Sheet updated July 2012

Numbers and algebra

Index laws:

For  $a, b > 0$  and  $m, n$  real,

$$a^m b^m = (a b)^m$$

$$a^m a^n = a^{m+n}$$

$$(a^m)^n = a^{mn}$$

$$\frac{1}{a^m} = a^{-m}$$

$$\frac{a^m}{a^n} = a^{m-n}$$

$$a^0 = 1$$

For  $a > 0$  and  $m$  an integer and  $n$  a positive integer,

$$a^{\frac{m}{n}} = \sqrt[n]{a^m} = \left(\sqrt[n]{a}\right)^m$$

Simple interest:

$I = Prt$ , where  $P$  is the principal,  $r$  is the rate per year and  $t$  is the time in years

Compound interest:

$A = P(1 + r)^t$  compounded annually

$A = P\left(1 + \frac{r}{n}\right)^{nt}$  compounded  $n$  times a year

Differentiation:

If  $f(x) = y$  then  $f'(x) = \frac{dy}{dx}$

Powers:

If  $f(x) = x^n$  then  $f'(x) = nx^{n-1}$

or

If  $y = x^n$  then  $\frac{dy}{dx} = nx^{n-1}$

Product rule:

If  $y = f(x) g(x)$

or

If  $y = uv$

then  $y' = f'(x) g(x) + f(x) g'(x)$

then  $\frac{dy}{dx} = \frac{du}{dx} v + u \frac{dv}{dx}$

Integration:

$$\int x^n dx = \frac{x^{n+1}}{n+1} + c, n \neq -1$$

Antiderivative:

Given  $\frac{dy}{dx} = x^n$  then  $y = \frac{x^{n+1}}{n+1} + c, n \neq -1$

See next page

Space and measurement

In any triangle  $ABC$ :

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$\cos A = \frac{b^2 + c^2 - a^2}{2bc}$$

$$A = \frac{1}{2} ab \sin C, \text{ where } A \text{ is the area}$$

Circle:

$C = 2\pi r = \pi D$ , where  $C$  is the circumference,  $r$  is the radius

and  $D$  is the diameter

$A = \pi r^2$ , where  $A$  is the area

Triangle:

$A = \frac{1}{2} bh$ , where  $b$  is the base and  $h$  is the perpendicular height

Parallelogram:

$$A = bh$$

Trapezium:

$A = \frac{1}{2}(a + b)h$ , where  $a$  and  $b$  are the lengths of the parallel sides and  $h$  is the perpendicular height

Prism:

$V = Ah$ , where  $V$  is the volume,  $A$  is the area of the base and  $h$  is the perpendicular height

Pyramid:

$$V = \frac{1}{3} Ah$$

Cylinder:

$S = 2\pi rh + 2\pi r^2$ , where  $S$  is the total surface area  
 $V = \pi r^2 h$

Cone:

$S = \pi rs + \pi r^2$ , where  $s$  is the slant height

$$V = \frac{1}{3} \pi r^2 h$$

Sphere:

$$S = 4\pi r^2$$

$$V = \frac{4}{3} \pi r^3$$

Chance and data

Probability:

For any event  $A$  and its complement  $\bar{A}$

$$P(A) + P(\bar{A}) = 1$$

In a normal distribution approximately:

68% of values lie within one (1) standard deviation of the mean

95% of values lie within two (2) standard deviations of the mean

99.7% of values lie within three (3) standard deviations of the mean.

*Note: Any additional formulas identified by the examination panel as necessary will be included in the body of the particular question.*