

Copyright for test papers and marking guides remains with *West Australian Test Papers*.
Test papers may only be reproduced within the purchasing school according to the advertised Conditions of Sale.
Test papers should be withdrawn after use and stored securely in the school until Thursday 2nd July 2020.



SEMESTER ONE

MATHEMATICS SPECIALIST UNIT 1


2020

SOLUTIONS

Calculator-free Solutions

1. (a) (i) $-\mathbf{b}$ ✓
- (ii) $\frac{1}{2} \mathbf{a} + \mathbf{b}$ ✓
- (b) $\overrightarrow{RX} = \overrightarrow{RP} + \frac{2}{7} \overrightarrow{PQ}$ ✓
- $= -\mathbf{b} + \frac{2}{7} (\mathbf{b} - \mathbf{a})$ ✓
- $= -\frac{2}{7} \mathbf{a} - \frac{5}{7} \mathbf{b}$ ✓ [5]
2. (a) (i) 5 ✓
- (ii) 21 ✓
- (b) (i) $x = 3$ ✓
- (ii) $x = 4$ ✓
- (iii) $x = 7$ ✓
- (iv) $x = 2$ ✓
- (c) (i) $\begin{pmatrix} 7 \\ 5 \end{pmatrix} = 21$ ✓
- (ii) $\begin{pmatrix} 2 \\ 1 \end{pmatrix} \times \begin{pmatrix} 5 \\ 4 \end{pmatrix} + \begin{pmatrix} 2 \\ 2 \end{pmatrix} \times \begin{pmatrix} 5 \\ 3 \end{pmatrix}$ ✓
- $= 20$ ✓
- (d) (i) ${}^5P_5 = 120$ ✓
- (ii) $4 \times 3 \times 2 \times 1 \times 2! = 48$ ✓✓ [12]

3. (a) (i)

(ii) $OA \times OB = OC^2$ But $OC = OX$

$$\therefore OA \times OB = OX^2$$

This is the converse of the secant/tangent theorem

 $\therefore OX$ is a tangent

$$OA \times OB = OC^2$$

But $OC = OY$

$$\therefore OA \times OB = OY^2$$

This is the converse of the secant/tangent theorem

 $\therefore OY$ is a tangent(b) ${}^{16}C_{10}$ or ${}^{16}C_6$ (c) (i) ${}^{16}C_{10} \times {}^{16}C_4 \times {}^2C_2$ (ii) $3! = 6$ ways

[10]

4. (a) (i) $-i - 13j$

(ii) 5

(iii) $-9i - 12j + 2i + 6j$

$$= -7i - 6j$$

(iv) $\mathbf{b} - \mathbf{a} = -6\mathbf{i} + 2\mathbf{j}$

$$|\mathbf{b} - \mathbf{a}| = \sqrt{40} = 2\sqrt{10}$$

$$\therefore \hat{\mathbf{d}} = -\frac{3}{\sqrt{10}}\mathbf{i} + \frac{1}{\sqrt{10}}\mathbf{j} = \frac{\sqrt{10}}{10}(-3\mathbf{i} + \mathbf{j})$$

(b) $|\mathbf{a}| = |3\mathbf{i} - 6\mathbf{j}| = \sqrt{45} = 3\sqrt{5}$


$$|\mathbf{c}| = |\mathbf{i} + 3\mathbf{j}| = \sqrt{10}$$

$$\begin{pmatrix} 1 \\ 3 \end{pmatrix} \cdot \begin{pmatrix} 3 \\ -6 \end{pmatrix} = (3\sqrt{5})(\sqrt{10})\cos\theta$$

$$\therefore \cos\theta = \frac{3 - 18}{15\sqrt{2}} = -\frac{15}{15\sqrt{2}} = -\frac{1}{\sqrt{2}}$$

$$\therefore |\mathbf{c}| \cos\theta \hat{\mathbf{a}} = \sqrt{10} \times -\frac{1}{\sqrt{2}} \times \frac{1}{3\sqrt{5}} \begin{pmatrix} 3 \\ -6 \end{pmatrix} = \begin{pmatrix} -1 \\ 2 \end{pmatrix}$$


(c)



5. (a) $\angle ABC = 90^\circ$ ✓ [12]
 Thale's Theorem ✓
 (b) $\angle CAD = 68^\circ$ ✓
 Revolution equal to 360 degrees ✓
 (c) $\angle OCB = 45^\circ$ ✓
 Equal angles at base of isosceles triangle ✓ [6]
6. Assume $\exists x, y \in \mathbb{Z} : 8x - 24y = 5$ ✓
 $\therefore 8(x - 3y) = 5$ ✓
 $\therefore x - 3y = \frac{5}{8}$ ✓
 Since $x, y \in \mathbb{Z}$, then $(x - 3y) \in \mathbb{Z}$, but $\frac{5}{8} \notin \mathbb{Z}$. ✓
 This is a contradiction as $(x - 3y)$ cannot both belong and not belong to the integer set. ✓
 \therefore False, there are no integers for which this statement is true. ✓ [5]

Calculator-Assumed Solutions

7.



$$A = \begin{pmatrix} -30\sin(60) \\ 30\cos(60) \end{pmatrix} \quad B = \begin{pmatrix} -22\sin(45) \\ -22\cos(45) \end{pmatrix} \quad C = \begin{pmatrix} 42\sin(\theta) \\ 42\cos(\theta) \end{pmatrix}$$

$$30\cos(60) - 22\cos(45) + 42\cos(\theta) = 0$$

$$15 - 11\sqrt{2} + 42\cos(\theta) = 0$$

$$\cos(\theta) = \frac{11\sqrt{2} - 15}{42}$$

$$\theta = 89.24^\circ \therefore \text{bearing} = 089.24^\circ T$$

$$-30\sin(60) - 22\sin(45) + 42\sin(89.24^\circ) = v$$

$$v = 0.46 \text{ km/h}$$

✓✓

✓

✓

✓


✓

[6]

8. (a) (i) **LM** parallel to x -axis so $b = 5$ ✓
 $(a - 1)^2 + (5 - 2)^2 = 5^2$ ✓
 $(a - 1)^2 = 16$
 $a - 1 = \pm 4$
 $a = 5$ or $a = -3$, but $a > 3 \therefore a = 5$ ✓
 $M = (5, 5)$
 $\mathbf{KM} \cdot \mathbf{NM} = \begin{pmatrix} 4 \\ 3 \end{pmatrix} \cdot \begin{pmatrix} 5 - c \\ 5 \end{pmatrix} = 0$
(ii) ✓
 $20 - 4c + 15 = 0$
 $35 = 4c$
 $c = \frac{35}{4}$ ✓
(b) (i) $x = 5$ ✓
 $y = 2$ ✓
(ii) If $y = 2$ & $x \geq 6$, then **$BC \parallel AD$** ✓
If **$AB \parallel DC$** ✓
 $k \begin{pmatrix} 1 \\ 2 \end{pmatrix} = \begin{pmatrix} 6 - x \\ 4 - y \end{pmatrix}$ ✓
 $2(6 - x) = (4 - y)$
 $2x - y = 8$ ✓
(iii) $\mathbf{OA} = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$ $\mathbf{OB} = \begin{pmatrix} 2 \\ 4 \end{pmatrix} = 2 \begin{pmatrix} 1 \\ 2 \end{pmatrix} = 2\mathbf{OA}$ ✓
 $\therefore \mathbf{OA} \parallel \mathbf{OB} \therefore A$ and B are collinear ✓

[13]

9. (a)



✓✓

$$\frac{\sin(\theta)}{190} = \frac{\sin 110}{40}$$

(b)

$$\theta = 10.95^\circ$$

✓

$$\therefore \text{Bearing} = 110^\circ \text{T}$$

✓

$$(c) \quad 940^2 = 190^2 + |r|^2 - 2 \times r \times 190 \cos 110^\circ$$

✓

$$\therefore |r| = 857.9 \text{ km/h}$$


✓

$$\therefore \text{Time taken} = \frac{4350}{857.9} = 5 \text{ h } 4 \text{ mins}$$

✓

[7]

10. (a)



 \therefore 50 like tea only

✓

(b) None like neither

✓

(c)



$$n(J \cup T \cup C) = n(T) + n(C) + n(J) - n(T \cap C) - n(T \cap J) - n(J \cap C) + n(T \cap C \cap J)$$

$$\therefore 100 = 70 + 50 + 60 - 20 - 40 - 30 + x$$

✓

$$\therefore = 90 + x$$

✓

$$\therefore x = 10$$

✓

$$(d) \quad (i) \quad \frac{70}{100}$$

✓

$$(ii) \quad \frac{20}{100}$$

$$(e) \quad 41$$


✓

✓

[8]

11. (a) $\overrightarrow{OE} = \frac{1}{2} \overrightarrow{OB}$ ✓
- $\overrightarrow{OE} = \frac{1}{2} \left(\mathbf{c} + \frac{1}{3} \mathbf{a} \right)$
- $\overrightarrow{OE} = \frac{1}{2} \mathbf{c} + \frac{1}{6} \mathbf{a}$ ✓
- $\overrightarrow{OF} = \overrightarrow{OC} + \frac{1}{2} \overrightarrow{CA}$ ✓
- $\overrightarrow{OF} = \mathbf{c} + \frac{1}{2} (\mathbf{a} - \mathbf{c})$
- $\overrightarrow{OF} = \frac{1}{2} \mathbf{a} + \frac{1}{2} \mathbf{c}$ ✓
- (b) $\overrightarrow{CE} = \overrightarrow{OE} - \overrightarrow{OC} = \frac{1}{2} \mathbf{c} + \frac{1}{6} \mathbf{a} - \mathbf{c} = \frac{1}{6} \mathbf{a} - \frac{1}{2} \mathbf{c}$ ✓
- $\overrightarrow{BF} = \overrightarrow{OF} - \overrightarrow{OB} = \frac{1}{2} \mathbf{a} + \frac{1}{2} \mathbf{c} - \mathbf{c} - \frac{1}{3} \mathbf{a} = \frac{1}{6} \mathbf{a} - \frac{1}{2} \mathbf{c}$ ✓
- $\therefore \overrightarrow{CE} = \overrightarrow{BF}$
- $\overrightarrow{EF} = \overrightarrow{OF} - \overrightarrow{OE} = \frac{1}{2} \mathbf{a} + \frac{1}{2} \mathbf{c} - \frac{1}{2} \mathbf{c} - \frac{1}{6} \mathbf{a} = \frac{1}{3} \mathbf{a}$ ✓
- $\overrightarrow{CB} = \overrightarrow{OB} - \overrightarrow{OC} = \mathbf{c} + \frac{1}{3} \mathbf{a} - \mathbf{c} = \frac{1}{3} \mathbf{a}$ ✓
- $\therefore \overrightarrow{EF} = \overrightarrow{CB}$ [8]
12. (a) $3 > 2$ but $-3 < -2$ is false ✓
- (b) "If the triangle is not isosceles, then the triangle does not have two equal sides." ✓
Yes it is always true since the original implication is always true by definition of isosceles triangles. ✓
- (c) "If n is divisible by 4, then n is divisible by 8". ✓
The converse is not always true, as shown by the counter-example where $n = 12$ ✓
- (d) FOR ALL natural numbers p , EXISTS a real number q , such that q is one less than p . ✓✓ [7]

13. (a)

Let $\angle BAD = x^\circ$ \therefore reflex $\angle BOD = 2x^\circ$

Angle at the centre theorem ✓

Similarly let $\angle BCD = y^\circ$ $\therefore \angle BOD = 2y^\circ$

Angle at the centre theorem ✓

Since $2x + 2y = 360$ then $x + y = 180^\circ$

✓

Hence $\angle BAD + \angle BCD = 180^\circ$

✓

QED

(b) $\angle EAG + \angle BCD = 180^\circ$ Cyclic quadrilateral $\therefore \angle EAG \equiv \angle ECF$

✓

In $\triangle EAG$ and $\triangle ECF$ $\angle AEG \equiv \angle CEF$ Bisector

✓

 $\angle EAG \equiv \angle ECF$ Proven

✓

 $\therefore \angle CFE \equiv \angle BFG$ Vertically Opposite $\therefore \angle AGF \equiv \angle BFG$ QED

✓

(c) (i) Converse

✓

(ii) Yes

✓

[10]

14. (a) (i) $\mathbf{b} \cdot \mathbf{c} = 0$

$$\therefore -8x - 8 = 0$$

✓

$$\therefore x = -1$$

✓

(ii) $\mathbf{a} \cdot \mathbf{c} = |\mathbf{a}| \times |\mathbf{c}| \times \cos(60)$

✓

$$\therefore \begin{pmatrix} 5 \\ -3 \end{pmatrix} \cdot \begin{pmatrix} x \\ -8 \end{pmatrix} = \frac{1}{2} \times \sqrt{34} \times \sqrt{x^2 + 64}$$

✓

$$\therefore 5x + 24 = \frac{1}{2} \times \sqrt{34} \times \sqrt{x^2 + 64}$$

✓

$$\therefore x = -0.13$$

✓

(iii) $\mathbf{a} - \mathbf{b} = 13\mathbf{i} - 4\mathbf{j}$

✓

$$\therefore |\mathbf{a} - \mathbf{b}| = \sqrt{185}$$

$$\therefore |\mathbf{c}| = \sqrt{x^2 + 64}$$

✓


$$\therefore \sqrt{x^2 + 64} = 2\sqrt{185}$$

✓

$$\therefore x = \pm 26$$

✓

(b) (i)



(ii) $\cos(0) = 1$

So, the dot product of a vector with itself is its magnitude squared multiplied by 1.

(iii) $\mathbf{a} \cdot \mathbf{a} = (\mathbf{b} - \mathbf{c}) \cdot (\mathbf{b} - \mathbf{c})$

$$\therefore |\mathbf{a}|^2 = (\mathbf{b} - \mathbf{c}) \cdot (\mathbf{b} - \mathbf{c})$$

$$\therefore |\mathbf{a}|^2 = \mathbf{b} \cdot \mathbf{b} - 2(\mathbf{b} \cdot \mathbf{c}) + \mathbf{c} \cdot \mathbf{c}$$

$$\therefore |\mathbf{a}|^2 = |\mathbf{b}|^2 + |\mathbf{c}|^2 - 2 \times |\mathbf{b}| \times |\mathbf{c}| \times \cos\theta$$

$$\therefore \frac{|\mathbf{a}|^2 - |\mathbf{b}|^2 - |\mathbf{c}|^2}{2 \times |\mathbf{b}| \times |\mathbf{c}|} = \cos\theta$$

[14]

15. (a) $\mathbf{F}_1 = -\frac{3\sqrt{2}}{2} \mathbf{i} + \frac{3\sqrt{2}}{2} \mathbf{j}$

$$\mathbf{F}_2 = \frac{3\sqrt{3}}{2} \mathbf{i} + \frac{3}{2} \mathbf{j}$$

$$\mathbf{F}_3 = -2\mathbf{i} - 2\sqrt{3}\mathbf{j}$$

$$\therefore \mathbf{F}_1 + \mathbf{F}_2 + \mathbf{F}_3 = \begin{pmatrix} -1.523 \\ 0.157 \end{pmatrix}$$

$$\tan^{-1}\left(\frac{0.157}{1.523}\right) = 5.89^\circ$$

 \therefore $\therefore 276^\circ\text{T}$

(b) $\mathbf{a} + \mathbf{b} = -5\mathbf{i} + \mathbf{j}$

$$\mathbf{c} = 5\mathbf{i} - \mathbf{j}$$

$$|\mathbf{c}| = \sqrt{26}$$

Direction = 101°T

[10]

16. let $\angle BEQ = \alpha \therefore \angle QBE = \alpha$ and let $\angle QEC = \beta \therefore \angle QCE = \beta$

$$\therefore \angle BEC = \alpha + \beta, \angle BQE = 180 - 2\alpha \text{ and } \angle EQC = 180 - 2\beta$$

$$\therefore 180 - 2\alpha + 180 - 2\beta = 180$$

$$\therefore 180 = 2\alpha + 2\beta$$

$$\therefore 90 = \alpha + \beta = \angle BEC \therefore \perp$$

[4]

$$\begin{aligned}
 17. \quad & \frac{n!}{(n-3)!} \div \frac{n!}{(n-5)!5!} = \frac{1}{5} \quad \checkmark \\
 & \frac{(n-5)!5!}{(n-3)!} = \frac{1}{5} \\
 \therefore & \frac{120}{(n-3)(n-4)} = \frac{1}{5} \\
 \therefore & 600 = n^2 - 7n + 12 \quad \checkmark \\
 \therefore & n^2 - 7n - 588 = 0 \quad \checkmark \\
 \therefore & (n-28)(n+21) = 0 \rightarrow n = 28 \quad \checkmark \quad [5]
 \end{aligned}$$

18. (a) Independent term is the middle term. \checkmark

$$\begin{aligned}
 T_6 &= \binom{10}{5} (x)^5 \left(\frac{2}{x}\right)^5 \quad \checkmark \\
 \therefore & 8064 \quad \checkmark
 \end{aligned}$$

(b)

$$(a + \sqrt{3})^n = a^n + \binom{n}{1} a^{n-1}(\sqrt{3}) + \binom{n}{2} a^{n-2}(\sqrt{3})^2 + \binom{n}{3} a^{n-2}(\sqrt{3})^3 + \binom{n}{4} a^{n-3}(\sqrt{3})^4 + \dots$$

$$(a + \sqrt{3})^n = a^n + n\sqrt{3}a^{n-1} + \binom{n}{2} 3a^{n-2} + \binom{n}{3} 3\sqrt{3}a^{n-2} + \binom{n}{4} 9a^{n-3} + \dots \quad \checkmark$$

$$(a - \sqrt{3})^n = a^n + \binom{n}{1} a^{n-1}(-\sqrt{3}) + \binom{n}{2} a^{n-2}(-\sqrt{3})^2 + \binom{n}{3} a^{n-2}(-\sqrt{3})^3 + \binom{n}{4} a^{n-3}(-\sqrt{3})^4 + \dots$$

$$(a - \sqrt{3})^n = a^n - n\sqrt{3}a^{n-1} + \binom{n}{2} 9a^{n-2} - \binom{n}{3} 3\sqrt{3}a^{n-2} + \binom{n}{4} 9a^{n-3} + \dots \quad \checkmark$$

Hence,

$$(a + \sqrt{3})^n + (a - \sqrt{3})^n = 2a^n + 6\binom{n}{2}a^{n-2} + 18\binom{n}{4}a^{n-4} + \dots \quad \checkmark \checkmark$$

$$\therefore \forall_n n \text{ is an integer} \quad \checkmark \quad [8]$$