



**Shenton College
Semester 1 Examination, 2016**

Question/Answer Booklet

CHEMISTRY

ATCHE – Year 12 – ATAR

Student Name _____

Your marks	Marks available
	25
	60
	70
	155
%	

Time allowed for this paper

Reading time before commencing work: ten minutes
Working time for paper: two and a half hours

Materials required/recommended for this paper

To be provided by the supervisor

This Question/Answer Booklet
Multiple-choice Answer Sheet
Chemistry Data Sheet

To be provided by the candidate

Standard items: pens, pencils, eraser, correction fluid, ruler, highlighters

Special items: non-programmable calculators satisfying the conditions set out by the Curriculum Council for this course

Important note to candidates

No other items may be taken into the examination room. It is your responsibility to ensure that you do not have any unauthorised notes or other items of a non-personal nature in the examination room. If you have any unauthorised material with you, hand it to the supervisor before reading any further.

Structure of this paper


Section	Number of questions available	Number of questions to be answered	Suggested working time (minutes)	Marks available	Percentage of exam
Section One: Multiple-choice	25	25	50	50	25
Section Two: Short answer	11	11	60	70	35
Section Three: Extended answer	6	6	70	80	40
					100

Instructions to candidates

1. The rules for the conduct of Western Australian external examinations are detailed in the *Year 12 Information Handbook 2012*. Sitting this examination implies that you agree to abide by these rules.
2. Answer the questions according to the following instructions.

Section One: Answer **all** questions on the separate Multiple-choice Answer Sheet provided. For each question shade the box to indicate your answer. Use **only a blue or black pen** to shade the boxes. If you make a mistake, place a cross through that square, do not erase or use correction fluid, and shade your new answer. Marks will not be deducted for incorrect answers. No marks will be given if more than one answer is completed for any question.

Sections Two and Three: Write answers in this Question/Answer Booklet **with a blue or black, biro or pen**.

3. When calculating numerical answers, show your working,  reasoning **and answers** clearly unless instructed otherwise. Final answers should be to three significant numbers.
4. You must be careful to confine your responses to the specific questions asked and to follow any instructions that are specific to a particular question.
5. Spare pages are included at the end of this booklet. They can be used for planning your responses and/or as additional space if required to continue an answer.
 - Planning: If you use the spare pages for planning, indicate this clearly at the top of the page.
 - Continuing an answer: If you need to use the space to continue an answer, indicate in the original answer space where the answer is continued, i.e. give the page number. Fill in the number of the question(s) that you are continuing to answer at the top of the page.


Section One: Multiple-choice

25% (50 Marks)

This section has **25** questions. Answer **all** questions on the separate Multiple-choice Answer Sheet provided. For each question shade the box to indicate your answer. Use only a blue or black pen to shade the boxes. If you make a mistake, place a cross through that square, do not erase or use correction fluid, and shade your new answer. Marks will not be deducted for incorrect answers. No marks will be given if more than one answer is completed for any question.

Suggested working time for this section is 50 minutes.

Use the potential energy diagram shown below to answer question 1.





1. Which of the following gives the correct values for the enthalpy change (ΔH) and the activation energy (E_a) for the reverse reaction?

	ΔH (kJ mol ⁻¹)	E_a (kJ mol ⁻¹)
(a)	-300	+700
(b)	+300	+400
(c)	-300	+400
(d)	+300	+700

2. A catalyst was added to the reaction mixture. Comparing the catalysed reaction to the uncatalysed reaction, which of the following will remain the same?

- (a) the enthalpy change of the reaction.
- (b) the activation energy for the forward reaction.
- (c) the energy of the transition state
- (d) the activation energy of the reverse reaction.

Question 3 refers to the following graph, which represents the concentrations of SO₂ and SO₃ in the reaction shown below.




3. At the 20 minute mark, what changes could have been made to the system to produce the effects shown on the graph?
- The system temperature is increased or the concentration of NO is increased.
 - The system temperature is increased or the concentration of NO₂ is increased.
 - The system temperature is decreased or the concentration of NO is decreased.
 - The system temperature is decreased or the concentration of NO₂ is decreased.
4. If solid calcium carbonate is heated in a sealed container, the following equilibrium is established at 500°C and 600kPa pressure.



Which of the following statements about this equilibrium is correct?

- Adding more CO₂ to the system will reduce the amount of CaO present.
- Reducing the temperature of the system will increase the amount of CaO present.
- Increasing the pressure of the system to 1000 kPa by adding inert nitrogen gas will decrease the amount of CaCO₃ present.
- Adding more CaCO₃ to the system will cause an increase in CaO and CO₂ present.


5. The next question refers to the chemical process shown below.
This process has reached equilibrium in a closed system.



A small amount of $\text{HCl}(\text{g})$ is added to the system and dissolved. Which of the following lists correctly shows the changes in **concentration** for $\text{H}_2\text{O}(\text{l})$, $\text{CrO}_4^{2-}(\text{aq})$ and $\text{Cr}_2\text{O}_7^{2-}(\text{aq})$ as equilibrium is re-established?


	$[\text{H}_2\text{O}(\text{l})]$	$[\text{CrO}_4^{2-}(\text{aq})]$	$[\text{Cr}_2\text{O}_7^{2-}(\text{aq})]$
(a)	increased	decreased	decreased
(b)	decreased	increased	increased
(c)	unchanged	decreased	increase
(d)	unchanged	increased	decrease

6. The following equation represents a closed equilibrium system.



The value of K for this system at 2300°C is 2.5×10^{12} . This value tells us that;

- (a) The reaction is extremely exothermic
 (b) The reaction is extremely fast
 (c) The reaction essentially goes to completion
 (d) The reaction essentially does not occur
7. The reaction of iron(III) oxide with carbon monoxide gas is shown below:



Which one of the following changes to the system will decrease the rate of the forward reaction?

- (a) decreasing the volume of the reaction vessel
 (b) decreasing the pressure of $\text{CO}(\text{g})$ in the vessel
 (c) decreasing the $\text{Fe}_2\text{O}_3(\text{s})$ particle size
 (d) increasing the concentration of $\text{CO}_2(\text{g})$ in the system
8. When pure water is kept in the fridge, the pH rises above 7. This is because;
- (a) The water has become slightly basic
 (b) There is a higher concentration of OH^- ions than H_3O^+ ions
 (c) The self-ionisation of water occurs to a lesser extent

(d) The value of K_w is higher

9. What is the conjugate base of HC_2O_4^- ?

- (a) $\text{C}_2\text{O}_4^{2-}$
- (b) $\text{H}_2\text{C}_2\text{O}_4$
- (c) $\text{H}_3\text{C}_2\text{O}_4^+$
- (d) $\text{H}_2\text{C}_2\text{O}_4^-$

10. Which of the following pairs of words best describe the acidic properties of a 0.005 mol L^{-1} solution of H_2CO_3 ?

- (a) Dilute, strong
- (b) Dilute, weak
- (c) Concentrated, strong
- (d) Concentrated, weak

Use the information in the table below to answer questions 4 and 5.

Indicator	Colour (low pH – high pH)	pH range
Methyl yellow	Red – yellow	2.4 – 4.0
Bromocresol purple	Yellow – purple	5.2 – 6.8
Phenol red	Yellow – red	6.8 – 8.4
Cresol red	Yellow – red	7.2 – 8.8

11. Which indicator in the table above would be most suitable to identify the end point in a titration of hydrochloric acid solution against sodium carbonate solution?

- (a) Methyl yellow
- (b) Bromocresol purple
- (c) Phenol red
- (d) Cresol red

12. In an acid-base titration with an end point of pH 8.2, a chemist uses bromocresol purple as the indicator. The acid is added from the burette to the base in a conical flask and the base has an initial pH of 10.5.

What effect will this procedure have on the calculation of the unknown concentration for the base?

- (a) The concentration calculated will be higher than its true concentration.
- (b) The concentration calculated will be lower than its true concentration.
- (c) The concentration calculated will be accurate.
- (d) A calculation cannot be done as no colour change will be seen during the titration.

13. Sodium oxide (Na_2O) is a strong base, while sodium carbonate (Na_2CO_3) is a weak base. If an equal number of moles of each solid was placed into separate beakers containing 100 mL water, which of the following would be correct?

- (a) The beaker containing sodium oxide would have a lower pH
- (b) Both beakers would contain the same concentration of sodium ions
- (c) Both beakers would contain the same concentration of hydroxide ions
- (d) The sodium carbonate would not have dissolved as it is insoluble

14. When freshly cleaned aluminium metal is placed in dilute hydrochloric acid, a redox reaction occurs. Which of the following statements regarding the reaction is **NOT** correct?

- (a) The pH of the reaction mixture would fall as the reaction proceeded
- (b) The aluminium metal would be oxidised to form aluminium ions
- (c) The hydrogen ions would be reduced to form hydrogen gas
- (d) The gas formed would be positive for the 'pop test'

15. Which of the following would be classified as basic substances?

- (i) NaOH
- (ii) NH_3
- (iii) Na_2CO_3
- (iv) CH_3COOH
- (v) H_2CO_3

- (a) (i), (ii) and (iii)
- (b) (i), (iii) and (iv)
- (c) (i) and (iii)
- (d) (ii) and (v)

16. In which of the following equations is water acting as a Bronsted-Lowry base?

- (a) $\text{HCO}_3^- (\text{aq}) + \text{H}_2\text{O} (\text{l}) \rightarrow \text{CO}_3^{2-} (\text{aq}) + \text{OH}^- (\text{aq})$
- (b) $\text{SO}_4^{2-} (\text{aq}) + \text{H}_2\text{O} (\text{l}) \rightarrow \text{OH}^- (\text{aq}) + \text{HSO}_4^- (\text{aq})$
- (c) $\text{ClO}^- (\text{aq}) + \text{H}_2\text{O} (\text{l}) \rightarrow \text{HClO} (\text{aq}) + \text{OH}^- (\text{aq})$
- (d) $\text{H}_2\text{PO}_4^- (\text{aq}) + \text{H}_2\text{O} (\text{l}) \rightarrow \text{H}_3\text{O}^+ (\text{aq}) + \text{HPO}_4^- (\text{aq})$

17. Which solution would have the highest pH?

- (a) $1 \text{ mol L}^{-1} \text{HNO}_3$
- (b) $1 \text{ mol L}^{-1} \text{H}_2\text{SO}_4$
- (c) $1 \text{ mol L}^{-1} \text{H}_2\text{CO}_3$
- (d) $1 \text{ mol L}^{-1} \text{HCl}$

18. Which of the following pairs of half-cells would produce the greatest voltage under standard conditions?

- (a) Pb / Pb²⁺ and Ag⁺ / Ag
- (b) Zn / Zn²⁺ and Ni²⁺ / Ni
- (c) Ni / Ni²⁺ and Pb²⁺ / Pb
- (d) Zn / Zn²⁺ and Ag⁺ / Ag

19. A beaker held a dilute solution of copper(II) sulfate. A piece of metal was added to the beaker. Slowly the blue colour of the solution faded and a salmon pink metal began to precipitate. Which of the following metals would **not** cause these observations?

- (a) Magnesium
- (b) Iron
- (c) Nickel
- (d) Silver

20. If you had an equal mass of each of the metallic ions below, which would require the greatest number of moles of electrons to be reduced to the elemental metal?

- (a) Mn²⁺
- (b) Ni²⁺
- (c) Cd²⁺
- (d) Co²⁺

21. In which of the substances below is the underlined element in an oxidation state of +3?

- (i) HClO₂
- (ii) C₂H₂O₄
- (iii) H₂SO₃
- (iv) N₂O₃
- (v) HPO₂

- (a) (i), (ii) and (iv)
- (b) (i), (iv) and (v)
- (c) (ii), (iii) and (iv)
- (d) (i), (ii), (iv) and (v)

22. In which of the following is chlorine in the highest oxidation state?

- (a) HClO
- (b) Cl₂O₇
- (c) HCl
- (d) ClO₂

23. In which of the following reactions is oxygen reduced?

- (a) F_2O to O_2
- (b) H_2O to H_2O_2
- (c) SO_2 to SO_3
- (d) H_2O_2 to O_2

24. Which of the following is the strongest oxidant (oxidising agent) under standard conditions?

- (a) Cl_2 (g)
- (b) KMnO_4 (aq) / H^+ (aq)
- (c) Br_2 (l)
- (d) $\text{Na}_2\text{Cr}_2\text{O}_7$ (aq) / H^+ (aq)

25. Which of the following statements regarding redox reactions are true?

- (i) A reductant (reducing agent) will gain electrons
 - (ii) The oxidation number of the reductant (reducing agent) will increase
 - (iii) Oxidation numbers correspond to the number of electrons being transferred during a reaction
 - (iv) Oxidation cannot happen without reduction
 - (v) Redox reactions always involve the transfer of electrons
- (a) (i), (iii) and (v)
 - (b) (ii), (iv) and (v)
 - (c) (i), (ii) and (iv)
 - (d) (ii), (iii) and (iv)

End of Section One

Section Two: Short answer

% (57 Marks)

This section has **twelve (12)** questions. Answer **all** questions. Write your answers in the space provided.

Spare pages are included at the end of this booklet. They can be used for planning your responses and/or as additional space if required to continue an answer.

- Planning: If you use the spare pages for planning, indicate this clearly at the top of the page.
- Continuing an answer: If you need to use the space to continue an answer, indicate in the original answer space where the answer is continued, i.e. give the page number. Fill in the number of the question(s) that you are continuing to answer at the top of the page.

Suggested working time for this section is 60 minutes.

Question 26


(1 mark)

Equation	$\text{NH}_3(\text{g}) + \text{H}_2\text{O}(\text{l}) \rightleftharpoons \text{NH}_4^+(\text{aq}) + \text{OH}^-(\text{aq})$
Equilibrium constant expression	

Question 27

(4 marks)

An equilibrium is set up in a test tube by suspending some finely divided copper sulphide in a dilute solution of hydrochloric acid. The equation for the equilibrium is:




Complete the following table, giving your answers as “increases”, “decreases” or “no change”.

Change made to the equilibrium system	Immediate effect on rate of forward reaction	Effect on equilibrium yield of $\text{HS}^- (\text{aq})$
HCl acid is passed into the solution		
CuSO_4 solution is added		
More finely divided CuS is added		

Question 28


(7 marks)

Ammonia is an industrially important gas produced by the Haber process, as illustrated by the equation below:



The reaction is catalysed by iron (III) oxide, Fe_2O_3 .

The following graph shows the partial pressures of the three species involved in the reaction:



- (a) At the start, why does the partial pressure of the H_2 gas decrease more rapidly than that of N_2 gas? (1 mark)


- (b) What has occurred at the 30 minute mark to cause the changes shown in the graph? (1 mark)

- (c) By the 45 minute mark, what difference will the change imposed at the 30 minute mark have made to the rate of:

(i) the forward reaction? _____

(ii) the reverse reaction? _____ (2 marks)

At 50 minutes, the contents of the reaction vessel are rapidly compressed by reducing the volume. The changes in the partial pressures of the species are shown on the following graph, starting at 40 minutes.




- (d) Complete the graph above up to 70 minutes by showing how the partial pressures of each species change as a new equilibrium is achieved. (3 marks)

Question 29

(3 marks)

Methanol gas [CH_3OH] can be manufactured by combining hydrogen gas and carbon monoxide gas in an exothermic equilibrium reaction.
The equation for this reaction that reflects all the information given above.



- (a) List one condition that would favour a fast reaction rate and one condition that would favour a high yield of methanol gas in an industrial setting. (2 marks)

Conditions for a fast reaction rate	Conditions for a high yield

- (b) What temperature conditions would suit this reaction in an industrial setting? (1 mark)

Question 30**(2 marks)**

Write the equation for the reaction that occurs in each of the following procedures. If no reaction occurs, write 'no reaction'. For full marks show only net ionic equations.

(a) Magnesium oxide solid is mixed with hydrochloric acid solution.

(1 mark)

Equation: _____

(b) Barium nitrate solution is mixed with sulfuric acid solution.

(1 mark)

Equation: _____

Question 31**(3 marks)**

Water ionises according to the equation $\text{H}_2\text{O} (\text{l}) \rightleftharpoons \text{H}^+ (\text{aq}) + \text{OH}^- (\text{aq})$

At 25°C, $K_w = 1.0 \times 10^{-14} \text{ mol}^2 \text{ L}^{-2}$. At 50°C, the K_w value changes to approximately $5.5 \times 10^{-14} \text{ mol}^2 \text{ L}^{-2}$.

Use the information above, and Le Châtelier's principle, to predict whether the self-ionisation of water is an endothermic or exothermic process. Explain.

Question 32**(7 marks)**

(a) 30.0 mL of a sodium carbonate solution is made up to 300 mL with distilled water. The resultant solution has a sodium carbonate concentration of 0.108 mol L^{-1} . What mass of sodium carbonate was present in the original solution.

(2 marks)

- (b) 10.0 mL of a 2.00 mol L⁻¹ H₂SO₄ solution is added to 10.0 mL of 1.50 mol L⁻¹ NaOH. What is pH of the resultant solution? (5 marks)


Question 34

(8 marks)

There are many different theories used to describe acids and bases, such as Arrhenius, Bronsted-Lowry and Lewis.

- (a) Define an acid and a base according to the Bronsted-Lowry theory. (2 marks)

Consider the following acid-base reaction.



- (b) Label the conjugate acid-base pairs in the reaction above. (2 marks)
- (c) The HPO₄²⁻ ion is a weak base according to the Bronsted-Lowry theory. Write an equation showing how this ion behaves as a base in aqueous solution. (2 marks)

- (d) H_2PO_4^- has a K_a value of 6.2×10^{-8} . Explain why this value makes H_2PO_4^- a weak acid (2 marks)

Question 35

(5 marks)

A variety of substances are listed below. Use these substances to answer the following questions. Not all substances must be used, but each substance can only be used **once**.



- (a) Which two substances could be mixed together to form a buffer? (1 mark)

- (b) Using an equation, explain what happens when OH^- is added to the buffer mixture. (2 marks)

- (c) Which substance could be classified as a 'basic salt'? Write a hydrolysis equation to support your answer. (2 marks)

Question 36

(7 marks)

A sample of copper (II) iodide powder was dissolved in water. This solution was then split into three (3) test tubes and a separate reaction was performed in each.


- (a) Chlorine gas was bubbled through the first test tube. Write the oxidation and reduction half-equations for the reaction that would have taken place. (2 marks)

oxidation: _____

reduction: _____

- (b) Zinc metal was added to the second test tube. Describe the predicted observations for this reaction. (2 marks)

Lead(II) nitrate was added to the third test tube and the molecular equation for the reaction that took place is shown below.




- (c) Is this a redox reaction? Explain your answer. (2 marks)

Question 37

(2 marks)


Write a balanced half equation for the following:



Question 38

(10 marks)

The following diagram shows how a car battery can be recharged while the car is being driven. Both electrodes in a car battery are lead-based and the electrolyte in the cell is sulfuric acid. The reactions occurring at each electrode are also indicated in the diagram.



(a) On the diagram above label the cathode and anode, and label and show the direction of electron flow and the direction of cation flow. (3 marks)

(b) Write the overall equation for the recharging process. (1 mark)

(c) Identify the oxidant (oxidising agent) and reductant (reducing agent) in the recharging equation. Use oxidation numbers to support your answer. (3 marks)

(d) What is the function of the sulfuric acid electrolyte? (1 mark)

(e) What would happen to the pH of the electrolyte as the battery recharges? Explain your answer. (2 marks)

(f) Using your data sheet, calculate the overall voltage of this cell, under standard conditions (1 mark)

End of Section Two

Section Three: Extended answer

% (66 Marks)

This section contains **five (5)** questions. You must answer **all** questions. Write your answers in the space provided.

Spare pages are included at the end of this booklet. They can be used for planning your responses and/or as additional space if required to continue an answer.

- Planning: If you use the spare pages for planning, indicate this clearly at the top of the page.
- Continuing an answer: If you need to use the space to continue an answer, indicate in the original answer space where the answer is continued, i.e. give the page number. Fill in the number of the question(s) that you are continuing to answer at the top of the page.


Suggested working time for this section is 70 minutes.

Question 39


(7 marks)

The Pilbara iron ore industry uses vast amounts of ammonium nitrate explosive to break up the rock and ore. Much of the ammonium nitrate is produced in Kwinana, Western Australia, using the following process:


Step 1: Natural gas (from the North West Shelf) is reacted with steam.




Step 2: Hydrogen produced in the above process is reacted with nitrogen from the air using the Haber Process.




Step 3: Ammonia is reacted with oxygen in air.




Step 4: Nitrogen monoxide is reacted with oxygen in air.




Step 5: The nitrogen dioxide produced in the reaction above is reacted with water and oxygen to form nitric acid.



Step 6: Finally, nitric acid is reacted with ammonia to form ammonium nitrate.



The equation for Step 3 of the process is reproduced below.




It is an exothermic reaction ($\Delta H = -1130 \text{ kJ}$) and is carried out at 900°C and atmospheric pressure in the presence of a catalyst.

Use your understanding of reaction rates and equilibrium principles to explain why these conditions are employed for this reaction.

[illegible]

Question 40**(11 marks)**

A reaction that reaches equilibrium is studied in the laboratory. In this system, two colourless gases, nitrogen trichloride and oxygen, react to produce brown nitrogen dioxide gas and light green chlorine gas. The equation for the system is



- (a) State *three observable properties* that would stay constant when equilibrium is reached. (3 marks)

- (b) What is meant by the statement “dynamic equilibrium”. (2 marks)

- (c) Write the equilibrium constant expression for the system. (1mark)

- (d) State the effect of an increase in the pressure on the yield of chlorine? Explain (2 marks)


- (e) Complete the graph below to show what happens to the concentration of chlorine after the pressure was increased. (2 marks)



Question 41

(16 marks)

An electrochemical cell consists of a tin electrode in a solution of 1.0 mol L^{-1} tin(II) nitrate, to create a Sn/Sn^{2+} half cell, and a similarly constructed half cell composed of a chromium electrode in a solution of 1.0 mol L^{-1} chromium(III) nitrate. The two electrodes are joined by a piece of copper wire. A salt bridge, as shown in the diagram below, joins the two solutions.



(a) On the diagram, label (3 marks)

- the anode
- the direction of electron flow
- the direction of cation flow in the salt bridge.

(c) Would sodium carbonate be suitable as a salt for the salt bridge? Explain. (2 marks)

[illegible]

1.0 mol L ⁻¹ Sn(NO ₃) ₂	1.0 mol L ⁻¹ Cr(NO ₃) ₃
Sn	Salt Bridge
Cr	

STAGE 3 23 CHEMISTRY

(d) Why does the rate of production of electrical current from an electrochemical cell decrease as it operates? (1 mark)

(e) During the operation of an electrochemical cell, why is it important that the anode and cathode do not come into contact with each other? (1 mark)

(20 marks)

The acidity of wine is due mainly to the presence of the **soluble salt** potassium tartrate ($\text{KC}_3\text{H}_4\text{O}_4\text{COOH}$). It contains the tartrate ion which is a **weak monoprotic acid**. A student conducted an investigation to find the concentration of potassium tartrate in some red wine. To do this three 50 mL samples of wine were titrated with $0.0105 \text{ mol L}^{-1} \text{ NaOH}$. The data recorded by the student is shown here.

Volume of NaOH	First titration	Second titration	Third titration
Final reading (mL)	9.80 mL	10.30 mL	10.20 mL
Initial reading (mL)	0.05 mL	0.2 mL	0.1 mL
Volume used (mL)			

(a) Write the formula for the **tartrate ion** present in potassium tartrate ($\text{KC}_3\text{H}_4\text{O}_4\text{COOH}$).
(2 marks)

(b) With reference to the information given write an ionisation or hydrolysis equation for the tartrate ion? (2 marks)

(c) Use the information given to write a NET IONIC equation for the reaction occurring between the tartrate ion and sodium hydroxide solution? (2 marks)

- (d) Name the apparatus used to measure the wine samples and the sodium hydroxide samples. State what is used for the final rinse of the apparatus? (4 marks)

Wine sample measured with _____

Apparatus rinsed with _____

Sodium hydroxide sample measured with _____

Apparatus rinsed with _____

- (e) Use the data in the table to find the average volume of NaOH used. (2 marks)

- (f) Calculate the percentage of potassium tartrate in the wine. (4 marks)

- (f) A solution of potassium tartrate ($\text{KC}_3\text{H}_4\text{O}_4\text{COOH}$) can act as a buffer. What does this mean? Explain how this could occur. (4 marks)

Question 43**(6 marks)**

The **dry cell** and the **lead acid cell** are two commercial galvanic cells. The following questions refer to the use and operation of these two cells.

- (a) Some of the components of the **dry cell** and **lead acid cell** are listed here. State to which cell they belong and in a few words state their role in the operation of the cell.

Zinc Cell _____ role

_____ (1)

PbO₂ Cell _____ role

_____ (1)

Porous paper Cell _____ role

_____ (1)

Graphite Cell _____ role

_____ (1)

- (b) Write the **oxidation half reaction** for each cell.

dry cell


_____ (1)


lead acid cell


_____ (1)

Question 44**(6 marks)**

Write separate half equations and an overall redox equation for the following reactions.







END OF QUESTIONS