



MATHEMATICS METHODS Year 12

Section One:

Calculator-free

Student name \_\_\_\_\_

Teacher name \_\_\_\_\_

Time and marks available for this section

Reading time before commencing work: 2 minutes

Working time for this section: 15 minutes

15 marks

Materials required/recommended for this section

*To be provided by the supervisor*

This Question/Answer Booklet

Formula Sheet

*To be provided by the candidate*

Standard items: pens (blue/black preferred), pencils (including coloured), sharpener, correction fluid/tape, eraser, ruler, highlighters

Special items: nil

Important note to candidates

No other items may be taken into the examination room. It is **your** responsibility to ensure that you do not have any unauthorised notes or other items of a non-personal nature in the examination room. If you have any unauthorised material with you, hand it to the supervisor **before** reading any further.

**Instructions to candidates**

1. Write your answers in this Question/Answer Booklet.
2. Answer all questions.
3. **Show all your working clearly.** Your working should be in sufficient detail to allow your answers to be checked readily and for marks to be awarded for reasoning. Incorrect answers given without supporting reasoning cannot be allocated any marks. For any question or part question worth more than two marks, valid working or justification is required to receive full marks. If you repeat an answer to any question, ensure that you cancel the answer you do not wish to have marked.
4. It is recommended that **you do not use pencil**, except in diagrams.

**Additional working space**

Question number: \_\_\_\_\_

(4 marks)

(1 mark)

(1 mark)

(2 marks)

See next page

Evaluate each of the following integrals:

(a)  $\int (1 + 5x)^3 dx$


(b)  $\int x e^{x^2-1} dx$

(c)  $\int \cos 2x \sin^3 2x dx$

Question 8

(6 marks)

The diagram below shows an area bounded by the  $x$ -axis, the function  $y = \sqrt{ax}$  and the function  $ax - (a - 4)y = 4a$ .



(a) Write an expression involving integrals to calculate this shaded area. (3 marks)

(b) Given that the shaded area is 31.5 squared units, determine the value of  $a$ . (3 marks)

End of questions

**Question 2**

**(3 marks)**

It is given that  $f(x) = x \cos x$  and  $f'(x) = \cos x - x \sin x$ .

Use the above facts to find  $\int x \sin x \, dx$ .

**Question 7 continued**

(c)  $\int_0^2 2f(-x)dx$  (2 marks)


(d)  $\int_{-1}^1 f(x-1)dx$  (2 marks)

(e)  $\int_0^1 (\sin \pi x - f(x)) \, dx$  (2 marks)

Question 7

(10 marks)

The graph of the function  $y = f(x)$  is shown below over the domain  $-2 \leq x \leq 2$ .



The area of the regions P, Q, R and S enclosed by the curve and the x-axis are 5, 1, 2, and 3 squared units respectively.

Determine

- (a) the area enclosed by the curve and the x-axis for  $-1 \leq x \leq 1$ . (2 marks)

(b)  $\int_{-2}^2 f(x) dx$

(2 marks)

See next page

See next page

value of the constant  $k$ .

The area of the region bounded by the curve with equation  $y = kx^{\frac{1}{2}}$ , where  $k$  is a positive constant, the x-axis and the line with equation  $x = 9$  is  $27 \text{ units}^2$ . Find the

Question 3


(4 marks)

**Question 4**

**(4 marks)**

The graph of  $y = e^{x-1}$  is shown in the diagram below.

Calculate the exact area between the graphs of  $y = e^{x-1}$ ,  $y = 2 - x$  and the two axes.



End of questions

**Question 6 continued**

(c) What was the initial speed of the particle? (2 marks)

(d) Find the value(s) of  $t$  when the particle comes to rest, and the distance(s) from the origin at that time(s). (3 marks)

(e) Calculate the acceleration of the particle when the velocity is  $6 \text{ m/s}$ . (3 marks)

See next page

Question 6

(11 marks)


A particle moves in rectilinear motion, such that its displacement ( $x$ ) from the origin  $O$ , at any time  $t$  (seconds), is given as  $x = t(2t^2 - 5t + b) + 1$  metres.

It is known that at  $t = 3$ , the particle is 4 m to the right of the origin.

- (a) Determine the value of  $b$ . (1 mark)

- (b) Determine the velocity of the particle when  $t = 3$ .

(2 marks)



MATHEMATICS METHODS Year 12  
Section Two:  
Calculator-assumed

Student name \_\_\_\_\_

Teacher name \_\_\_\_\_

**Time and marks available for this section**  
Reading time before commencing work: 3 minutes  
Working time for this section: 30 minutes  
Marks available: 30 marks

**Materials required/recommended for this section**  
*To be provided by the supervisor*  
This Question/Answer Booklet  
Formula Sheet (retained from Section One)

***To be provided by the candidate***  
Standard items: pens (blue/black preferred), pencils (including coloured), sharpener, correction fluid/tape, eraser, ruler, highlighters

Special items: drawing instruments, templates, and up to three calculators approved for use in the WACE examinations

**Important note to candidates**


No other items may be taken into the examination room. It is **your** responsibility to ensure that you do not have any unauthorised notes or other items of a non-personal nature in the examination room. If you have any unauthorised material with you, hand it to the supervisor **before** reading any further.

**Instructions to candidates**

1. Write your answers in this Question/Answer Booklet.
2. Answer all questions.
3. **Show all your working clearly.** Your working should be in sufficient detail to allow your answers to be checked readily and for marks to be awarded for reasoning. Incorrect answers given without supporting reasoning cannot be allocated any marks. For any question or part question worth more than two marks, valid working or justification is required to receive full marks. If you repeat an answer to any question, ensure that you cancel the answer you do not wish to have marked.
4. It is recommended that **you do not use pencil**, except in diagrams.

**Question 5****(3 marks)**

The velocity function of a particle is given by  $v(t) = 3 \cos 5\pi t$  cm/sec.



Find the exact distance travelled by the particle from time  $t = 0$  to  $0.8$  sec.