

Copyright for test papers and marking guides remains with *West Australian Test Papers*.  
Test papers may only be reproduced within the purchasing school according to the advertised Conditions of Sale.  
Test papers should be withdrawn after use and stored securely in the school until Friday 30<sup>th</sup> November 2018.



**SEMESTER TWO**

**MATHEMATICS  
SPECIALIST  
UNITS 1 & 2**

**2018**


**SOLUTIONS**

**Calculator-free Solutions**

1. (a) (i)  ${}^8P_5$  or  ${}^8C_5 \times 5!$  ✓
- (ii)  ${}^4C_3 \times {}^4C_2 \times 5!$  ✓✓
- (b) (i)  $6^4 \times 5^2 \times 6!$  ✓
- (ii) '9' from first set only:  ${}^5C_3 \times {}^4C_2 \times 6!$  ✓
- '9' from second set only:  ${}^5C_4 \times {}^4C_1 \times 6!$  ✓
- no '9' chosen:  ${}^5C_4 \times {}^4C_2 \times 6!$  ✓
- total  ${}^5C_3 \times {}^4C_2 + {}^5C_4 \times {}^4C_1 + {}^5C_4 \times {}^4C_2 \times 6!$  ✓ [8]
2. (a) (i)  $i^{n+2} = i^n \times i^2 = -i \times -1 = i$  ✓
- (ii)  $i^{2n+1} = (i^n)^2 \times i = (-i)^2 \times i = -i$  ✓✓
- (b)  $\frac{1-i}{i+\frac{2}{i}} \times \frac{i}{i} = \frac{i-i^2}{i^2+2} = \frac{1+i}{-1+2} = 1+i$  ✓✓
- (c)  $(1+i)^4 - (1-i)^4$
- $i[(1+i)^2 + (1-i)^2] \times [(1+i)^2 - (1-i)^2]$
- $i[1+2i-1+1-2i-1] \times [1+2i-1-1+2i+1]$
- $i0 \times 4i = 0$  ✓✓ [7]
3. (a)  $5^3 = 5 + 1 \times 10 + 1 \times 10 \times 6 + 1 \times 4 \times 15 = 5 + 60 + 60 = 125$  ✓
- (b)  ${}^5C_2, {}^4C_1$  ✓✓
- (c)  $n^3 = n + {}^{n-1}C_0 \times {}^{n+1}C_1 \times {}^nC_2 + {}^nC_0 \times {}^{n-1}C_1 \times {}^{n+1}C_2$  ✓✓ [5]
4. (a) (i)  $\begin{bmatrix} 0 & -1 & 1 \\ 1 & 1 & -1 \\ 1 & 1 & 0 \end{bmatrix} \times \begin{bmatrix} 1 & 1 & 0 \\ -1 & -1 & 1 \\ 0 & -1 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$  ✓✓
- A and B are inverses of each other. ✓
- (ii)  $\begin{bmatrix} 0 & -1 & 1 \\ 1 & 1 & -1 \\ 1 & 1 & 0 \end{bmatrix} \times \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} -6 \\ 8 \\ 3 \end{bmatrix}$  ✓
- $\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 & -1 & 1 \\ 1 & 1 & -1 \\ 1 & 1 & 0 \end{bmatrix}^{-1} \times \begin{bmatrix} -6 \\ 8 \\ 3 \end{bmatrix} = \begin{bmatrix} 1 & 1 & 0 \\ -1 & -1 & 1 \\ 0 & -1 & 1 \end{bmatrix} \times \begin{bmatrix} -6 \\ 8 \\ 3 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \\ -5 \end{bmatrix}$  ✓✓

$$\therefore x=2, y=1, z=-5$$

4. (b) (i)  $T_1$  performs a rotation of  $180^\circ$



$$\therefore y = -(x-1)^2$$

✓✓

- (ii) Reflection of the line  $y=x$

✓

$$T_2 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$$

✓

- (iii) New Area  $|T_3| \times 10 \frac{2}{3} = 6 \times \frac{32}{3} = 64 \text{ unit s}^2$

✓✓

[12]

5. (a) (i)  $y = -3 \cos \left[ 4 \left( x - \frac{\pi}{12} \right) \right] = -3 \cos \left[ 4x - \frac{\pi}{3} \right]$


$$\therefore A = -3\omega = 4\theta = \frac{-\pi}{3}$$

✓✓✓

- (ii)  $y = -3 \sin \left( 4x + \frac{\pi}{6} \right)$

✓✓

(b)



✓ Scale factor  
( $y=2$  at  $x=\frac{5\pi}{24}$ )

✓ Period of  $\frac{\pi}{2}$

✓ Vertical Asymptote  
 $x = \frac{\pi}{6}, \frac{\pi}{2}$



6. (a) (i)  $\sqrt{2} \times \sqrt{8} = \sqrt{16} = 4$  ✓
- (ii) "If  $ab$  is irrational, then both  $a$  and  $b$  must be irrational" ✓
- (b)  $A \Rightarrow B$ : If the triangle has two equal sides, then it is isosceles,  
and therefore it has two congruent sides.  
 $\therefore A \Rightarrow B$  is valid and True. ✓
- $B \Rightarrow A$ : If the triangle has two congruent sides, then it is isosceles,  
and therefore it has two equal sides.  
 $\therefore B \Rightarrow A$  is valid and True. ✓
- $\therefore A \Leftrightarrow B$
- (c)  $\forall x \in P, \exists y \in P : xy \in Q$  ✓✓ [6]
7. Assume  $n$  is even and that  $n^3$  is odd. ✓
- Let  $m \in \mathbb{N} : n = 2m$  is even. ✓
- $n^3 = (2m)^3 = 8m^3 = 2(4m^3)$  ✓
- Since  $n^3$  cannot be both even and odd simultaneously, then this  
is a contradiction. And therefore  $n$  must be odd. ✓ [4]

**Calculator-assumed Solutions**

8. (a)  $z = \frac{4 \pm \sqrt{16-24}}{4} = 1 \pm \frac{\sqrt{2}}{2}i$  ✓

$$P(z) = \left(z - 1 - \frac{\sqrt{2}}{2}i\right) \left(z - 1 + \frac{\sqrt{2}}{2}i\right) \quad \checkmark$$

(b) Since  $a, b, c \in R$  then  $\bar{z} = 2 - i$  is also a solution ✓

$$\therefore R(z) = (z+1)(z-2-i)(z-2+i) \quad \checkmark$$

$$iz^3 - 3z^2 + z + 5$$

$$\therefore a = -3b = 1c = 5 \quad \checkmark$$

[5]

9. LHS  $i \frac{\tan \theta + \tan \frac{\pi}{4}}{1 - \tan \theta \times \tan \frac{\pi}{4}}$  ✓

$$i \frac{\tan \theta + 1}{1 - \tan \theta} \quad \checkmark$$

$$i \frac{\frac{\sin \theta}{\cos \theta} + 1}{1 - \frac{\sin \theta}{\cos \theta}} \quad \checkmark$$

$$i \frac{\frac{\sin \theta}{\cos \theta} + 1}{1 - \frac{\sin \theta}{\cos \theta}} \times \frac{\cos \theta}{\cos \theta} \quad \checkmark$$

$$i \frac{\sin \theta + \cos \theta}{\cos \theta - \sin \theta} \quad \checkmark$$

$$i \frac{\sin \theta + \cos \theta}{\cos \theta - \sin \theta} \times \frac{\cos \theta + \sin \theta}{\cos \theta + \sin \theta} \quad \checkmark$$

$$i \frac{\sin^2 \theta + \cos^2 \theta + 2 \sin \theta \cos \theta}{\cos^2 \theta - \sin^2 \theta} \quad \checkmark$$

$$i \frac{1 + \sin 2\theta}{\cos 2\theta} = i \text{RHS} \quad \checkmark$$

[8]

10. (a) (i)  $\begin{pmatrix} 3 \\ 1 \end{pmatrix} = k \begin{pmatrix} \alpha \\ -2 \end{pmatrix} \rightarrow k = \frac{-1}{2} \rightarrow \alpha = -6$  ✓✓

(ii)  $a + c = \begin{pmatrix} 3 + \alpha \\ -1 \end{pmatrix} \rightarrow \begin{pmatrix} 3 + \alpha \\ -1 \end{pmatrix} \cdot \begin{pmatrix} 1 \\ -3 \end{pmatrix} = 0 \rightarrow \alpha = -6$  ✓✓

(b)  $\begin{pmatrix} 1 \\ 5 \end{pmatrix} \cdot \begin{pmatrix} 6 \\ 4 \end{pmatrix} = \begin{vmatrix} 1 \\ 5 \end{vmatrix} \times \begin{vmatrix} 6 \\ 4 \end{vmatrix} \times \cos \theta \rightarrow \cos \theta = \frac{1}{\sqrt{2}}$  ✓

$$|m| = \left| \frac{1}{5} \right| = \sqrt{26} \quad \checkmark$$

$$\hat{n} = \frac{1}{2\sqrt{13}} \begin{pmatrix} 6 \\ 4 \end{pmatrix} \quad \checkmark$$

$${}_m \text{proj}_n = |m| \cos \theta \hat{n} = \sqrt{26} \times \frac{1}{\sqrt{2}} \times \frac{1}{2\sqrt{13}} \begin{pmatrix} 6 \\ 4 \end{pmatrix} = \begin{pmatrix} 3 \\ 2 \end{pmatrix} = 3i + 2j \quad \checkmark \quad [8]$$



11. (a)  $S_3 = \frac{1}{2!} + \frac{2}{3!} = \frac{1}{2} + \frac{2}{6} = \frac{5}{6}$  ✓

$$S_4 = \frac{1}{2!} + \frac{2}{3!} + \frac{3}{4!} = \frac{1}{2} + \frac{2}{6} + \frac{3}{24} = \frac{23}{24} \quad \checkmark$$

(b)  $\frac{3!-1}{3!} = \frac{6-1}{6} = \frac{5}{6}$  ✓

$$\frac{4!-1}{4!} = \frac{24-1}{24} = \frac{23}{24} \quad \checkmark$$

(c)  $S_n = \frac{1}{2!} + \frac{2}{3!} + \frac{3}{4!} + \dots + \frac{n-1}{n!} = \frac{n!-1}{n!}$  ✓

(d) For  $n=2$ :  $S_2 = \frac{1}{2}$  and  $\frac{2!-1}{2!} = \frac{1}{2} \therefore$  True for  $n=2$  ✓

Assume true for  $n=k$ :

$$S_k = \frac{1}{2!} + \frac{2}{3!} + \frac{3}{4!} + \dots + \frac{k-1}{k!} = \frac{k! - 1}{k!} \quad \checkmark$$

For  $n=k+1$ :

$$S_{k+1} = \frac{1}{2!} + \frac{2}{3!} + \frac{3}{4!} + \dots + \frac{k-1}{k!} + \frac{(k+1)-1}{(k+1)!}$$

$$\textcolor{red}{+} S_k + \frac{k}{(k+1)!} = \frac{k!-1}{k!} + \frac{k}{(k+1)!} \quad \checkmark$$

$$i \frac{k!-1}{k!} \times \frac{k+1}{k+1} + \frac{k}{(k+1)!}$$

$$i \frac{k!(k+1) - (k+1)}{(k+1)!} + \frac{k}{(k+1)!}$$


$$i \frac{(k+1)! - 1}{(k+1)!} \text{ as required}$$

Therefore, True for  $n = k + 1$ , and since true for  $n = 2$ ,

by induction the conjecture is true for all whole numbers.

[10]

12. (a)



$$(b) \quad w = \begin{pmatrix} -80 \cos 45^\circ \\ 80 \sin 45^\circ \end{pmatrix} = \begin{pmatrix} -40\sqrt{2} \\ 40\sqrt{2} \end{pmatrix} \quad \checkmark$$

$$r = \begin{pmatrix} r \cos 42^\circ \\ r \sin 42^\circ \end{pmatrix} \quad \checkmark$$

$$p = \begin{pmatrix} 870 \cos(90^\circ - \theta) \\ 870 \sin(90^\circ - \theta) \end{pmatrix} = \begin{pmatrix} 870 \sin \theta \\ 870 \cos \theta \end{pmatrix} \quad \checkmark$$

$$(c) \quad p + \begin{pmatrix} -40\sqrt{2} \\ 40\sqrt{2} \end{pmatrix} = \begin{pmatrix} r \cos 42^\circ \\ r \sin 42^\circ \end{pmatrix}$$

$$\therefore |p| = \left| \begin{pmatrix} 40\sqrt{2} + r \cos 42^\circ \\ -40\sqrt{2} + r \sin 42^\circ \end{pmatrix} \right| = 870 \quad \checkmark$$

$$\text{CAS: } r = 862.14 \text{ kmh}^{-1} \quad \checkmark$$

$$\theta = 53.27^\circ \quad \checkmark$$

$$(d) \quad t = \frac{d}{v} = \frac{878}{862.14} = 1.018 \text{ hrs} = 1 \text{ hour } \wedge 1 \text{ minute} \quad \checkmark \checkmark$$

[11]

$$13. (a) \quad P(Q - I) = Q + I$$

$$\therefore P = (Q + I)(Q - I)^{-1} \quad \checkmark$$

$$\therefore \begin{bmatrix} 2 & 3 \\ -2 & 2 \end{bmatrix} \times \begin{bmatrix} 0 & 3 \\ -2 & 0 \end{bmatrix}^{-1} = \begin{bmatrix} 2 & 3 \\ -2 & 2 \end{bmatrix} \times \frac{1}{6} \begin{bmatrix} 0 & -3 \\ 2 & 0 \end{bmatrix} \quad \checkmark$$

$$\therefore P = \begin{bmatrix} 1 & -1 \\ \frac{2}{3} & 1 \end{bmatrix} \quad \checkmark$$

$$(b) \quad A^4 = A^2 \times A^2 = (3A - 2I) \times (3A - 2I)$$

$$= 9A^2 - 12A + 4I \quad \checkmark$$

$$\hookrightarrow 9(3A - 2I) - 12A + 4I$$

✓

$$\hookrightarrow 27A - 18I - 12A + 4I$$

$$\therefore A^4 = 15A - 14I$$

✓

[6]

14. (a)  $f(x) = R \cos(2x + \theta)$

$$\hookrightarrow R \cos 2x \cos \theta - R \sin 2x \sin \theta = \cos 2x - \sqrt{3} \sin 2x$$

✓

$$\therefore R \sin \theta = \sqrt{3} R \cos \theta = 1$$

✓

$$\Rightarrow R = 2, \theta = 60^\circ$$

✓

$$f(x) = 2 \cos(2x + 60^\circ)$$

✓

(b)  $f(x) = 2 \cos(2x + 60^\circ) = -1$

$$\cos(2x + 60^\circ) = \frac{-1}{2}$$

$$\therefore 2x + 60^\circ = 120^\circ, 240^\circ, 480^\circ, 600^\circ$$

✓

$$x = 30^\circ, 90^\circ, 210^\circ, 270^\circ$$

✓

[6]

15. (a) (i)  $P = \begin{bmatrix} -2 & 3 \\ 2 & -3 \end{bmatrix} \times \begin{bmatrix} 2 & 1 \\ 0 & 2 \end{bmatrix}^{-1} = \begin{bmatrix} -1 & 2 \\ 1 & -2 \end{bmatrix}$

✓

$$\therefore k = -2$$

✓

(ii)  $|P| = k(k+1) - 2 = k^2 + k - 2 \neq 0$

✓

$$\therefore k \neq -2, 1$$

✓

(b) (i)  $M^{2n+1} = (M^2)^n \times M = I^n \times M = M$

✓

(ii)  $M^{-2n} = (M^2)^{-n} = I^{-n} = I$

✓


(c)  $A^n = \begin{bmatrix} 1 & 0 \\ 2^n - 1 & 2^n \end{bmatrix}$

✓✓

[8]

16. (a)

Customers [x10]



tude  
d = 7hrs  
all waves



16. (b)  $C(t) = -80 \cos\left[\frac{2\pi}{7}(t-8.5)\right] + 100$

$$A = -80, \omega = \frac{2\pi}{7}, \phi = \frac{17\pi}{7}, v = 100$$

✓✓✓✓

OR  $A = 80, \omega = \frac{2\pi}{7}, \phi = \frac{24\pi}{7}, v = 100$

(c)  $-80 \cos\left[\frac{2\pi}{7}t - \frac{17\pi}{7}\right] + 100 = 140$

✓

CAS:  $t_1 = 10.83, t_2 = 13.17, t_3 = 17.83, t_4 = 20.18$

✓✓

Between 10:50am and 1:11pm

✓

and 5:50pm and 8:11pm

✓

[12]

17. (a) (i)  $5 - 4i$

✓✓

(ii)  $6i$


✓✓

(iii)  $3 + 5i$


✓✓

(iv)  $3 - 5i$

✓✓



17. (b) (i)



✓✓✓

(ii) Rotation ✓ of  $45^\circ$  anticlockwise ✓(iii) Powers that are multiples of 8 give  $z^{8n}=1$ 

✓

[14]

18. (a)  $|T| = \cos \theta \cos \phi + \sin \theta \sin \phi = 0$ 

✓

$$\therefore \cos(\theta - \phi) = 0$$

✓

$$\therefore \theta - \phi = \pm 90^\circ$$

✓

$$(b) (i) T = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$$

Rotation of  $90^\circ$  anti-clockwise

✓✓

$$(ii) T = \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix}$$

Reflection over the line  $y = -x$ 

✓✓

(c) i.e.  $T^2 = I$ 

$$\begin{bmatrix} \cos \theta & -\sin \phi \\ \sin \theta & \cos \phi \end{bmatrix} \times \begin{bmatrix} \cos \theta & -\sin \phi \\ \sin \theta & \cos \phi \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} \cos^2 \theta - \sin \theta \sin \phi & -\sin \phi (\cos \theta + \cos \phi) \\ \sin \theta (\cos \theta + \cos \phi) & \cos^2 \phi - \sin \theta \sin \phi \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

✓

$$\sin \theta (\cos \theta + \cos \phi) = -\sin \phi (\cos \theta + \cos \phi) = 0$$

$$\therefore \sin \theta = \sin \phi = 0 \text{ and } \cos^2 \theta = \cos^2 \phi = 1$$

✓✓

OR  $\cos \theta + \cos \phi = 0$  and  $\sin \theta = \sin \phi$ 

✓✓

[12]