

# Physics

## TRIAL EXAMINATION

### Written Paper Stage 2A

### Question/Answer Booklet

Student Number: In figures

--	--	--	--	--	--	--	--

In words


Student name: \_\_\_\_\_ Teacher name: \_\_\_\_\_

#### ***Time allowed for this paper***

Reading time before commencing work: Ten minutes  
Working time for paper: Three hours

#### ***Material required/recommended for this paper***

##### **To be provided by the supervisor**

This Question/answer booklet; Formulae and constants sheet

##### **To be provided by the candidate**

Standard items: pens, pencils, eraser or correction fluid, ruler, highlighter

Special items: scientific non-programmable calculator

#### ***Important note to candidates***

No other items may be taken into the examination room. It is **your** responsibility to ensure that you do not have any unauthorised notes or other items of a non-personal nature in the examination room. If you have any unauthorised material with you, hand it to the supervisor **before** reading any further.

**All calculations are to be set out in detail.** Marks may be awarded for correct equations and clear setting out, even if you cannot complete the calculation. Express **numerical answers** to two (2) or three (3) significant figures and include units where appropriate.

#### Student Marks

Section		Percentage of paper	Maximum mark	Student Mark
Section One 15 - 20 questions	Short Answer 35-45%	42%	45	
Section Two 5 – 7 questions	Problem Solving 45-55%	47%	51	
Section Three 1 -2 questions	Comprehension 5-15%	11%	12	
Overall mark	Units and Significant figures		2	
Student total Mark				
Student Percentage				

**Structure of this paper**

Section of exam	Suggested working time	Number of questions	Number of questions to be attempted	Marks available
Section One	70 minutes	14	all	45
Section Two	80 minutes	6	all	51
Section Three	30 minutes	1	all	12
Overall				2
[Total marks]				110

**Note:** the 'overall' section represents marks allocated to appropriate use of units and significant digits in final answers to numerical problems.

**Instructions to candidates**

1. The rules for the conduct of WACE examinations are detailed in the *Student Information Handbook*. Sitting this examination implies that you agree to abide by these rules.
2. Answer **all** questions in the spaces provided in this Question/Answer Booklet.
3. A blue or black ballpoint or ink pen should be used except for graphs where a pencil should be used.

**This paper is a combination of material from  
The Curriculum Council of Western Australia  
and  
Karen Harper; TDC Physics,  
Department of Education and Training**

**All material is copyright.**

*Part of this paper is published by the Curriculum Council of Western Australia  
27 Walters Drive  
OSBORNE PARK WA 6017*

**SECTION ONE: SHORT ANSWER**

This section has **FOURTEEN (14)** questions. Attempt **ALL** questions. Answer in the spaces provided.

Allow approximately 50 minutes to complete this section *[46 marks]*.

---

**Question 1**

Atom Q has an Atomic Number of 11 and has 12 neutrons. Atom R is a singly positively charged ion which also has an Atomic Number of 11 but has 11 neutrons. Use this information to fill in the table below: *[3 marks]*

atom	Number of electrons	Mass Number	Number of protons
Q			
R			

**Question 2**

Jemima measured the activity of a 10.00 g of radioisotope X. The original activity of the sample was 21.0 kBq and 12 days later the activity was 1.3125 kBq.

**(a)** What is the half-life of the radioisotope? *[3 marks]*

**(b)** Jemima also had a 100 g sample of X. Would this second sample have a greater, equal or smaller half-life than the original 10 g sample of X?

Circle the correct answer:

GREATER


EQUAL

SMALLER

*[1 mark]*

**Question 3**

Phosphorus-32 decays to sulfur-32 by the emission of a beta particle:



The masses of the particles involved are:

particle	Mass ( $\mu$ )
${}_{15}^{32}\text{P}$	31.97272
${}_{16}^{32}\text{S}$	31.97091
${}_{-1}^0\beta$	0.0005486

(a) Calculate the mass difference created by this decay.

[1 mark]

(b) This mass difference shows that energy is released during this reaction. Explain.

[1 mark]

---



---



---

(c) Calculate the energy in joules released created by this decay.

[2 marks]

**Question 4**

A nucleus of copper-65 ( ${}_{29}^{65}\text{Cu}$ ) absorbs a neutron and shortly afterward emits a  $\beta$  particle. Write two balanced nuclear equations to show each of these processes.

Absorbing a neutron: \_\_\_\_\_

[1 mark]

Emitting a  $\beta$  particle: \_\_\_\_\_

[1 mark]

**Question 5**

A beta particle has a mass of  $9.11 \times 10^{-31}$  kg and a speed of  $2.50 \times 10^6$  m s<sup>-1</sup>. How fast would an alpha particle (mass  $6.68 \times 10^{-27}$  kg) have to move in order to have the same momentum as the beta? [2 marks]

**Question 6**

A 64.0 kg athlete stored elastic potential energy in her leg muscles when she was crouched, ready to jump vertically. When she jumped, she found that she could raise her centre of mass through a vertical distance of 1.34 m.

(a) Calculate how much gravitational potential energy she had at the highest point of the jump. [2 marks]

(b) Calculate the speed with which she left the ground as she took off. [2 marks]

**Question 7**

Identify "X" in each of the following equations and place its symbol in the brackets to the right of the equation. Only where appropriate, name the **type of radiation** produced, leave blank otherwise. [3 marks]



(note: *m* denotes in excited state)

**Question 8**

From rest, a sneeze can reach  $161 \text{ kmh}^{-1}$  within 5.00 cm of your nose.

(a) Calculate the acceleration of the sneeze.

[2 marks]

(b) Calculate how long it would take for the sneeze to reach 5.00 cm.

[2 marks]

**Question 9**

Rodney the Rodent applied a force of constant magnitude to push a 0.50 kg block of cheese across a smooth flat tabletop. The cheese accelerated from rest to a speed of  $0.30 \text{ m s}^{-1}$  in a time of 20 seconds. Calculate the magnitude of the force that Rodney applied to the cheese. [2 marks]

**Question 10**

A car's engine supplies  $3.45 \times 10^6$  J of energy every minute to run the car.

(a) What is the power output of the car?


[2 marks]

(b) Calculate the car's maximum speed against a total resistance force of  $1.50 \times 10^3$  N.

[2 marks]

**Question 11**

The graph shows how the force exerted by a cricket bat on a cricket ball varies with time.



(a) The ball's change in momentum can be determined from this graph. Is the change in momentum related to the gradient, or the area under the curve?

Circle

the correct answer:

GRADIENT

AREA UNDER THE CURVE

[1 mark]

(b) Calculate the magnitude of the ball's change in momentum between  $t = 0$  and  $t = 0.02$  s.

[2 marks]

**Question 12**

A student is measuring the decay of a nuclear source. She finds that the source has a count of 15.60 kBq decays in a three hour period. Calculate the activity of the source. (2 marks)

**Question 13**

Heather is helping her Mum around the house by doing the weekly washing in the new washing machine. She reads in the instruction booklet that the spin cycle on a washing machine can spin as fast as  $6.00 \times 10^2$  rpm (revolutions per minute). Using this information, Heather correctly calculates the speed and velocity of a sock on the edge of the tub after one and a quarter turns of the tub? Assume the tub has a diameter of 0.550 m. You must show all working for full marks.

[3 marks]





**Question 14**

Chris and Rohan were playing golf and Chris was using one of the new jumbo sized metal “woods” to hit the golf ball. Find the **change in momentum** of the golf ball and the **force** delivered to the **golf ball** by the metal wood from the following data:

Speed of metal ‘wood’ at impact	= 70 m s <sup>-1</sup>
Mass of the club head	= 150 g
Mass of golf ball	= 50 g
Speed of ball on leaving the club head	= 140 m s <sup>-1</sup>
Time for which contact was made	= 0.02 s

[4 marks]

**SECTION TWO: PROBLEM-SOLVING**

This section has **SIX (6)** questions. Attempt **ALL** questions. Answer in the spaces provided.

Allow approximately 45 minutes to complete this section [51 marks].

**Question 1 [11 marks]**

One way to treat certain cancers is to irradiate them. Cobalt-60 is one radioisotope that can project a beam of gamma radiation to dose cancer patients.

- (a)** Two other kinds of radiation can be given off from radioisotopes, apart from gamma radiation. Name these two other radiations, and describe what each consists of. [4 marks]

*Particle One:* Name: \_\_\_\_\_

What it consists of: \_\_\_\_\_

\_\_\_\_\_

*Particle Two:* Name: \_\_\_\_\_

What it consists of: \_\_\_\_\_

\_\_\_\_\_

- (b)** Considering its use as a gamma source in cancer treatment, would you expect the half-life of the cobalt-60 to be long or short? Explain.

**Circle** the correct answer:                      LONG                      SHORT [1 mark]

Explanation: \_\_\_\_\_

\_\_\_\_\_

[1 mark]

- (c)** How does gamma treatment of cancer work? That is, what effect does a beam of gamma radiation have on cells? [2 marks]

\_\_\_\_\_

\_\_\_\_\_

- (d)** The cobalt-60 gives off gamma radiation all the time, in all directions. This radiation must be stopped when it is not needed. Name a material suitable for blocking gamma emissions:

\_\_\_\_\_

[1 mark]

- (e)** Beta radiation can also be used in medical technology to diagnose certain conditions but alpha radiation is never used. Fully explain, using your understanding of alpha radiation, why alpha radiation is never used to diagnose medical conditions. [2 mark]

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

**Question 2 [7 marks]**

Celia used a Geiger-Muller counter to measure the activity of a radioactive sample. The label said the sample activity was 47.8 kBq, but when Celia set up the G-M tube 50 cm from the source, she found the counter measured an average of 390 counts per second.

- (a) Celia recorded count rates of 388, 372 and 410. Explain why all the counts were different.

[1 mark]

---

---

---

- (b) Suggest two possible explanations for difference between the activity according to the label and Celia's measurement.

[2 marks]

---

---

---

---

- (c) Celia moved the G-M tube further from the source. State how you would expect the reading to change as a result, and explain your reasoning.

☒ Circle the correct answer:      INCREASE      STAY THE SAME      DECREASE

[1 mark]

Explain your choice:

---

---

---

[1 mark]

- (d) Celia put away the radioactive source, but found the G-M tube was still measuring about 30 Bq, even with no radioactive material nearby. Explain what this radiation is and where it comes from.

[2 marks]

---

---

---

---

---

---

**Question 3 [8 marks]**

In 2001 a very rich American tried to fly a balloon non- stop around the world. It was launched at Northam and moved vertically upwards with a constant velocity of  $4.00 \text{ m s}^{-1}$ . Unfortunately after only 5.00 mins an essential piece of equipment became dislodged and fell out of a gap in the basket and fell towards the ground. (Assume no air resistance.)

**(a)** How high was the balloon after 5.0 mins?

*[2 marks]*

**(b)** With what speed will it impact on the ground?


*[3 marks]*

**(c)** Neglecting air resistance, how long does it take for the essential piece of equipment to reach the ground?  
*marks]*

*[3*

**Question 4 [8 marks]**


The mass of a rocket resting on the launch pad is  $3.00 \times 10^3$  kg.



- (a) Calculate the weight of the rocket, in newtons, while it is sitting on the launch pad with its engine turned off. [2 marks]
- (b) Calculate the magnitude of the resultant force required to accelerate the rocket upwards from the launch pad at  $40.0 \text{ m s}^{-2}$ . [2 marks]
- (c) Using your answers to (a) and (b) above, calculate the thrust force provided by the rocket motor at the moment of lift-off. [2 marks]


**Question 4 continues**

- (d) Clearly show, using arrows to represent force vectors, the magnitude and directions of all the forces acting on the rocket at the moment it lifts off the launch pad with an acceleration of  $40.0 \text{ m s}^{-2}$  upwards: *[2 marks]*



**Question 5 [8 marks]**

Sam and Max used a motion detector to measure the speed of a motor scooter. Their results made a straight line graph when they plotted kinetic energy on one axis and the square of the scooter's speed on the other axis:



(b) Show that the gradient of this graph represents  $\left( \frac{1}{2} \times \text{scooter's mass} \right)$ . [3 marks]

(c) Use the gradient to calculate the mass of the scooter. If you could not obtain an answer to part (a), use gradient = 65 kg. [2 mark]

### Question 6 [9 marks]

At the Royal Show a gunman was demonstrating an unusual stunt. He used a 70.0 cm long rifle which was placed on the ground so that its barrel points upwards. A 300 g rubber ball was dropped downwards towards the rifle barrel by an assistant. The gunman fires the gun so that the ball, which is travelling at  $4.00 \text{ ms}^{-1}$  downwards, is hit when it is 10 cm from the top of the barrel. The 20.0 g bullet was travelling at  $460 \text{ ms}^{-1}$  upwards when it hits the ball and remains in the ball causing the ball to change direction and travel upwards. The assistant measured the maximum height above the ground that the ball reached after it had been hit by the bullet. What measurement did the assistant get? (Assume no loss of energy to other forms and no friction.)

- a. Using your understanding of conservation of momentum, calculate the initial velocity with which the ball / bullet combination moves upwards. Don't forget direction. (4 marks)
- b. Knowing the initial velocity of the ball / bullet combination, find the distance upwards the ball will travel after being hit by the bullet. (3 marks)
- c. Now find the maximum height the ball will reach above the ground. (2 marks)



**SECTION THREE: COMPREHENSION**


This section has **ONE (1)** question. Attempt **ALL** sections. Answer in the spaces provided.

Allow approximately 25 minutes to complete this section [12 marks].

A group of testers at a car design centre drove a number of cars directly into a solid barrier and measured the 'collapse length'  $CL$ . The testers used the same type of car in each test. They defined the collapse length as the difference between the overall length of the car before collision ( $\ell_i$ ), and the length after collision ( $\ell_f$ ).

Mathematically, this can be shown as:  $CL = \ell_i - \ell_f$

The diagram shows how the testers measured  $\ell_i$  and  $\ell_f$ .




The following table shows their results.

Impact speed, $v$ [m s <sup>-1</sup> ]	Collapse length, $CL$ [m]	$\frac{v^2}{CL}$ [m s <sup>-2</sup> ]
6.0	0.22	164
10.0	0.35	286
14.0	0.50	392
19.0	0.74	488
22.0	0.93	520

The test group found that the force  $F$  acting on a car of mass  $m$  during a collision at an impact speed of  $u$  is given by

$$F = \frac{mv^2}{2CL}.$$

- (a) Use the table of values to plot a graph of  $\frac{v^2}{CL}$  (y-axis) vs  $v$  (x-axis) on the graph paper. Draw a smooth line of best fit, by eye. [3 marks]



- (b) Use the graph to determine the value of  $\frac{v^2}{CL}$  for a car impacting the barrier at  $16 \text{ m s}^{-1}$ . [1 marks]
- 

- (c) Calculate the force acting on a  $1500 \text{ kg}$  car impacting the barrier at a speed of  $16 \text{ m s}^{-1}$ . If you could not obtain an answer to part (c), use  $\frac{v^2}{CL} = 550 \text{ m s}^{-2}$ . [2 marks]

- (d) If the car in the previous questions was  $3.20 \text{ m}$  in length, what would be the resultant length after

the crash?

[1 mark]

- (e) One of the testers called the force acting on the car during the collision an 'average force'. Explain why this force is an average force.

[2 marks]

---

---

---

---

---

---

---

- (f) Cars today are designed so that the bonnet does crumple. Explain, using your understanding of physics principles, why it is safer on the passengers in the car if the bonnet crumples. [3 marks]

---

---

---

---

---

---

---

---

---

---

**END OF 2A PAPER**

Part of this paper is published by the Curriculum Council of Western Australia  
27 Walters Drive  
OSBORNE PARK WA 6017