

# SOLUTIONS

## 2018

### Semester Two

#### MATHEMATICS

#### METHODS

#### UNITS 3 & 4



Test papers should be withdrawn after use and stored securely in the school until Wednesday 10th October 2018.  
Test papers may only be reproduced within the purchasing school according to the advertised Conditions of Sale.

Copyright for test papers and marking guides remains with West Australian Test Papers.

WATP

19. (a)  $SA = (3x)(x) + (3x)(h)(2) + (x)(h)(2)$
- $= 3x^2 + 8xh$
- $V = 3x^2h = 18 \rightarrow h = \frac{6}{x}$
- $SA = 3x^2 + 8x\left(\frac{6}{x}\right)^2 = 3x^2 + \frac{48}{x}$
- $\frac{d(SA)}{dx} = 6x - \frac{48}{x^2}$
- $\therefore \text{Max occurs when } 6x - \frac{48}{x^2} = 0 \rightarrow x = 2$
- $\therefore h = 1.5 \text{ and } SA = 36 \text{ m}^2$
- (b)  $\frac{d(SA)}{dx} \approx \frac{\delta SA}{\delta x}$
- $\delta SA = \left( \frac{d(SA)}{dx} \right) (\delta x) = \left( 6x - \frac{48}{x^2} \right) (0.1)$
- When  $x = 2$ ,  $\frac{dS}{dt} = 0.35 \text{ cm}^3/\text{sec}$
- (c)  $\frac{dS}{dt} = \frac{ds}{dx} \times \frac{dx}{dh} \times \frac{dh}{dt}$
- $When x = 1, \frac{dS}{dt} = 0.35 \text{ cm}^3/\text{sec}$
- (d)  $\frac{dS}{dt} = \left( 6x - \frac{48}{x^2} \right) \left( -\frac{1}{12x^3} \right) (0.1)$
- When  $x = 1.1, \frac{dS}{dt} = 0.0794$
- (e) (i)  $X \sim \text{Bin}(4, 0.1587)$
- $P(X > k) = 0.9 \rightarrow k = 2.11$
- $\therefore P(X = 1) = 0.378$
- (ii)  $0.1587 \times 0.5^3 \times (4C_1)$
- (iii)  $0.1587 \times 0.5 \times (4C_1) = 0.0378$
20. (a)  $X \sim N(1.6, 0.4^2)$
- $P(X > 2) = 0.1587$
- (b)  $P(X < k) = 0.9 \rightarrow k = 2.11$
- (c)  $P(X = 1) = 0.378$
- (d)  $X \sim \text{Bin}(4, 0.1587)$
- $P(X > k) = 0.9 \rightarrow k = 2.11$
- $\therefore P(X = 1) = 0.378$
- (e)  $0.1587 \times 0.5^3 \times (4C_1) = 0.0378$

**Calculator-free Solutions**

1. (a)  $\log_x 9 + \log_x x^2$  ✓  
 $= \frac{\log_3 9}{\log_3 x} + 2\log_x x$  ✓  
 $= \frac{2}{p} + 2$  or  $\frac{2+2p}{p}$  ✓  
(b)  $3^{2\log_3 3} = 3^2$  ✓  
 $= 9$  ✓  
(c)  $3^{2x^2}(2x) = 2x(3^{2x^2})$  ✓✓ [7]
2. (a)  $\frac{dV}{dt} = 4\pi(12t - t^2)^2(12 - 2t)$  ✓  
 $\frac{dV}{dt} = 0$  when  $t(12 - t) = 0$  or  $(12 - 2t) = 0$  ✓  
 $\therefore t = 0$  or  $6$  or  $12$  ✓  
Using the sign table yields Maximum occurs at  $t = 6$  ✓  
(b) (i)  $V = \frac{4}{3}\pi r^3 = \frac{4}{3}\pi(12t - t^2)^3$  ✓  
 $\therefore r = 12t - t^2$  ✓  
(ii) When  $t = 6$  then  $r = 36$  ✓  
 $r = 36$  is the maximum value of  $r$ , so max Volume. ✓ [8]
3. (a)  $p = 0.5[(2) + (1.5) + (1.2)] = 0.5(4.7)$  ✓  
 $= 2.35$  ✓  
 $q = 0.5[(3) + (2) + (1.5)] = 0.5(6.5)$  ✓  
 $= 3.25$  ✓  
 $\int_1^{2.5} \frac{3}{x} dx = [3 \ln x]_1^{2.5}$  ✓  
(b)  $= 3 \ln 2.5 - 3 \ln 1 = 3 \ln 2.5$  ✓ [6]
4. (a) 0.07 ✓  
(b) 0.25 ✓  
(c)  $P(1) = (0.1)^2 = 0.01$   
 $P(2) = (0.15)^2 = 0.0225$   
 $\therefore P(1) + P(2) = 0.0325$  ✓  
(d) Unbiased would be uniform ✓ [5]

16.  $X \sim \text{Bin}(36, 0.8)$  ✓✓  
(a)  $P(X \leq 30) = 0.7536$  ✓  
(b) (i)  $E(\hat{p}) = 0.8$  ✓  
(ii)  $\text{VAR}(\hat{p}) = \sqrt{\frac{(0.8)(0.2)}{36}} = \frac{1}{15}$  or  $0.067$  ✓✓  
(c)  $(0.9)^{10} = 0.349$  ✓✓ [7]
17. (a) Take proportions as follows:  
 $20 - 29 : \frac{727}{3100} \times 100 = 23$   
 $30 - 39 : \frac{1050}{3100} \times 100 = 34$   
 $40 - 49 : \frac{800}{3100} \times 100 = 26$   
 $50 - 59 : \frac{523}{3100} \times 100 = 17$  ✓✓✓  
(b) (i) Normal distribution ✓  
(ii)  $\mu = 0.3387$  ✓  
Standard deviation =  $\sqrt{\frac{(0.3387)(0.6613)}{100}} = 0.0473$  ✓  
(c)  $Y \sim N(0.3387, 0.0473^2)$  ✓  
 $P(X \geq 40) = P(Y \geq 0.4) = 0.0975$  ✓✓ [9]
18. (a)  $P(7) = \frac{1}{6}$  ✓  
(b)  $P(<7) = \frac{15}{36}$  ✓  
(c)  $P(>7) = \frac{15}{36}$  ✓  
(d) \$12 ✓  
(e)
- | Game   | 7s             | Unders          | Overs           |
|--------|----------------|-----------------|-----------------|
| P(X)   | $\frac{6}{36}$ | $\frac{15}{36}$ | $\frac{15}{36}$ |
| Return | 20             | 1               | 1               |
- $\therefore E(X) = \frac{6}{36} \times 20 + \frac{15}{36} \times 1 + \frac{15}{36} \times 1 = \frac{150}{36} = 4.16$  ✓✓  
 $\therefore$  Club makes \$5 - \$4.16 = 84 cents per roll ✓ [7]

5. (a)  $\cos 5x - 5x \sin 5x$

(b)  $\int \frac{dx}{x} (x \cos 5x) = \int (\cos 5x - 5x \sin 5x) dx$

6.  $4e^{2x} - 9e^x - 9 = 0$   
Let  $y = e^x$   
 $(4y^2 - 9y - 9) = 0$   
 $4y^2 - 9y - 9 = 0$   
 $y = -\frac{3}{4}$  or  $y = 3$   
 $e^x = -\frac{3}{4}$  → no solution  
 $e^x = 3 \rightarrow x = \ln 3$   
and  $e^x = 3 \rightarrow x = \ln 3$

7. (a)

(b)  $A = \int_{\pi}^{\frac{3\pi}{2}} (\sin x - \cos x) dx$

8.  $E_1 = 1.645 \sqrt{\frac{n_1}{p(1-p)}}$

$E_2 = 1.645 \sqrt{\frac{9n_1}{p(1-p)}}$


$E_1 = \frac{\sqrt{9n_1}}{\sqrt{n_1}}$

15. (a) (i)  $v(t) = \int (3t + 5) dt = \frac{3t^2}{2} + 5t + 20$   
 $v(3) = 48.5 \text{ m/sec}$

(ii)  $x(t) = \int \left( \frac{3t^2}{2} + 5t + 20 \right) dt = \frac{t^3}{2} + \frac{5t^2}{2} + 20t - 10$   
 $x(3) = 86 \text{ m}$

(b)  $v = \frac{3t^2}{2} + 5t + 20 = 0$  when stopped  
However  $\frac{3t^2}{2} + 5t + 20 \neq 0$  so never stops

(c) Distance travelled =  $x(3) - x(0)$   
 $86 - (-10) = 96 \text{ m}$



- $= \frac{1}{3}$  ✓ [3]
9. (a)  $\ln |\sin x| + c$  ✓✓
- (b)  $\left[ \ln |\sin x| \right]_{\frac{\pi}{4}}^{\frac{\pi}{2}}$  ✓
- $$= \ln(1) - \ln\left(\sin \frac{\pi}{4}\right) = 0 - \ln\left(\frac{1}{\sqrt{2}}\right) = \ln \sqrt{2}$$
- ✓
- $$= \frac{1}{2} \ln 2$$
- ✓ [5]
- Calculator-assumed Solutions**
10. (a)  $p + q = 0.4$  ✓
- $$0.2 + 4p + 2.7 + 16q + 2.5 = 8.2$$
- ✓
- $$\therefore \text{From CAS } p = 0.3 \text{ and } q = 0.1$$
- ✓
- $$(b) E(X) = 0.2 + 0.6 + 0.9 + 0.4 + 0.5 = 2.6$$
- ✓
- $$(c) E(Y) = 1 - 5.2 = -4.2$$
- ✓ [5]
11. (a)  $\hat{p} = \frac{86}{200} = 0.43$  ✓
- (b)  $CI = 0.43 \pm 1.645 \sqrt{\frac{(0.43)(0.57)}{200}}$  ✓✓
$$= 0.43 \pm 0.0576$$
 ✓
$$\therefore 0.372 \leq p \leq 0.488$$
 ✓
- (c) The mean suggested by the CI is  $\frac{0.62 + 0.82}{2} = 0.72$  ✓
This does not lie within the 90% CI as calculated in (b) ✓
 $\therefore$  Evidence doesn't support the claim. ✓
(d) (i) Binomial requires independent events and a constant probability of success. Neither criteria apply here. ✓
- (ii)  $P(X=1) = \frac{7C_1 \times 3C_3}{10C_4}$  ✓✓
$$= 0.1$$
 ✓ [12]

12. (a)
- 
- $$f(x) = \begin{cases} \frac{1}{4000} & 6000 < x < 1000 \\ 0 & \text{otherwise} \end{cases}$$
- (b) (i)  $P(8000 \leq X \leq 9000) = 0.25$  ✓✓
(ii)  $P(X = 8500) = 0$  ✓
- (c)  $E(X) = \int_{6000}^{10000} \left( x \times \frac{1}{4000} \right) dx = 8000$  ✓✓
- (d) (i)  $\text{VAR}(X) = \int_{6000}^{10000} \left( (x - 8000)^2 \times \frac{1}{4000} \right) dx = 1333333.33$  ✓✓
 $\therefore \text{st dev} = 1154.7 = \$1155$  ✓
(ii) Standard deviation = \$1155 Not affected by change of origin ✓ [10]
13. (a) (i)  $\ln V = \frac{7.22 - 6.91}{4} t + 6.91$  ✓
 $\therefore \ln V = 0.0775t + 6.91$  ✓
(ii)  $V = e^{0.0775t + 6.91}$  ✓
 $\therefore V = 1002.25e^{0.0775t}$  ✓
- (b)  $2000 = 1002.25e^{0.0775t}$  ✓
 $\therefore t = 8.91 \text{ years}$  ✓
- (c)  $V = V_0 e^{kt}$ 
 $\therefore 2V_0 = V_0 e^{\frac{rt}{100}}$  ✓
 $e^{\frac{rt}{100}} = 2$  ✓
 $\frac{rt}{100} = \ln 2 \rightarrow t = \frac{100 \ln 2}{r}$  ✓
 $\therefore t = \frac{69}{r}$  ✓ [9]