

# **SOLUTIONS**

**2018**

**MATHEMATICS  
METHODS  
UNITS 1 & 2**

**SEMESTER TWO**



Copyright for test papers and marking guides remains with West Australian Test Papers.  
Test papers may only be reproduced within the purchasing school according to the advertised Conditions of Sale.  
Test papers should be withdrawn after use and stored securely in the school until 30 November 2018.

**Calculator-free Solutions**

1. (a)  $\frac{2^{3x}}{2^y} = 2^4 = 16$  ✓✓  
     (b)  $u^{\frac{7}{2}} - \left(\frac{5}{2} + \frac{1}{2}\right)$  ✓  
          $= u^{\frac{1}{2}} = \sqrt{u}$  ✓ [4]
2.  $ab = 15$  and  $a + b = 8$   
 $a = 3, b = 5$  or  $a = 5, b = 3$  ✓  
 $abx^2 + (2b + 7a)x + 14 = 15x^2 + cx + 14$   
 $2b + 7a = 2(5) + 7(3)$  or  $2b + 7a = 2(3) + 7(5)$  ✓✓  
 $c = 31$  or  $41$  [4]
3. (a)  $m = \frac{4}{3}$  ✓  
 $-5 = \frac{4}{3}(3) + c$   
 $\therefore c = -9$   $\therefore y = \frac{4}{3}x - 9$  ✓  
 $-4x + 3y = -27$  ✓  
 $-8x + 6y = -14$   
(b)  $9x - 6y = 12$   
 $x = -2$   $y = -5$  ✓  
 $D(-2, -5)$  ✓  
(c)  $4(k-2) - 3(2k-3) = 7$   
 $-2k + 1 = 7$   
 $k = -3$  ✓ [7]
4.  $\frac{1.2 \times 10^{-4}}{3 \times 10^{-7}} = 4 \times 10^2 = 400$  ✓✓ [2]
5. (a)  $2(3xy) + 2(3x^2) + 2xy = 32$  ✓  
 $3xy + 3x^2 + xy = 16$  ✓  
 $3x^2 + 4xy = 16$   
(b)  $V = 3x^2y$  and  $y = \frac{16 - 3x^2}{4x}$  ✓  
 $V = 3x^2\left(\frac{16}{4x} - \frac{3x^2}{4x}\right)$  ✓  
 $V = 12x - \frac{9x^3}{4}$   
(c)  $V' = 12 - \frac{27x^2}{4}$   
 $12 - \frac{27x^2}{4} = 0$  for stationary point ✓  
 $x^2 = \frac{16}{9}$  ✓  
 $x = \frac{4}{3}$  (discard  $-\frac{4}{3}$ ) ✓



(d)

$V(x)$	$\uparrow$	$\frac{4}{3}$	$\downarrow$
$V'(x)$	$+$	0	$-$

 $V$  has a maximum value


✓✓

[9]

6. (a)  $T_{n+1} = T_n + 10 \quad T_1 = -3$

✓✓

25. (a)



✓✓✓

- (b) (i) 0.5323  
(ii) 0.2726  
(iii) 0.51  
(iv) 0.8026

[7]

(b)  $T_2 = -8$

✓

[4]

$T_3 = -32$

✓

7. (a)  $y = 10x^2 - 2x^3 - 16x + c$

✓

$3 = 10(2)^2 - 2(2)^3 - 16(2) + c$

✓

$c = 11$

$y = 10x^2 - 2x^3 - 16x + 11$

✓

26. The particle's initial displacement is 5 m to the right of the origin.

✓

$v = 3t^2 - 12t \therefore \text{Initial velocity} = 0$

✓

[2]

(b) Gradient of  $x$ -axis = 0

✓

$\frac{dy}{dx} = 20(2) - 6(2)^2 - 16 = 0$

✓

 $\therefore m = 0$  Tangent is horizontal and parallel to the  $x$ -axis.

✓

[5]

8. (a) Vertical translation 12 units down

✓

Horizontal dilation by factor  $\frac{1}{2}$ .

✓

(b)  $7^x = 7^{2x} - 12$

✓

Let  $7^x$  be  $k$ .  $k^2 - k - 12 = 0$

✓

$(k - 4)(k + 3) = 0$

✓

$7^x = 4 \quad 7^x \neq -3$

✓

One solution, therefore intersects at only one point.

[5]

9. (a)  $y = \frac{x^3}{2} + 2x - 7$

✓

$\frac{dy}{dx} = \frac{3}{2}x^2 + 2$

✓

$m = 8$

✓


$1 = 8(2) + c$

✓

$y = 8x - 15$


✓

(b)



✓✓

[6]



10. (a) Period =  $4\pi$   
Amplitude = 2

✓

[5]

[6]

### Calculator-assumed Solutions

✓✓✓

11. (a) The number of phones she is given to repair for the week.  
(b) She fixes 23 per day, for 4 days  $23 \times 4 = 92$  phones

✓✓✓

[6]

12. (a)  $x^2 + y^2 = 4$   
(b)  $\tan \theta = \sqrt{3} \therefore \theta = \frac{\pi}{6}$   
(c) radius = 2 units  
(d)  $T_1 = 155 \quad T_2 = 128 \quad T_3 = 99 \quad T_4 = 68$   
5 hours and 34 minutes

✓✓✓

[6]

13. (a)  $R = R_0(0.95)^t$   
 $R = R_0(1.085)^t$   
 $W_0(1.085)^t = 10 \times W_0(0.95)^t$   
 $t = 17.329$  years  
 $t = 18$  years the wallabies will be more than triple.

✓✓✓

[6]

14. (a)  $W = W_0(1.085)^t$   
 $R = R_0(0.95)^t$   
 $10W_0 = R_0$   
 $10W_0 = 1.085W_0$   
 $W_0 = 655$

✓✓✓

[6]

15. (a)  $r = -\frac{3}{2}$   
 $-2\left(1 - \left(-\frac{3}{2}\right)^n\right) = 30$   
 $1 - \left(-\frac{3}{2}\right)^n = 15$   
 $\left(-\frac{3}{2}\right)^n = -10$   
 $n = 9.003$   
10 terms

✓✓✓

[6]

16. (a)  $\text{Area of shaded part} = \text{triangle} - \text{sector}$   
 $= \frac{3}{2} \text{ units}^2$   
 $\text{Area of sector} = \frac{1}{2} r^2 \theta = \frac{1}{2} (2)^2 \left(\frac{3}{\pi}\right)$   
 $\text{Therefore area of } \Delta AOB = \frac{1}{2}(4)\sqrt{3} = 2\sqrt{3} \text{ units}^2$   
 $\cos 3 = 4$

✓✓✓

[6]

17. (a)  $\therefore \text{Sum of next 15 terms} = S_{30} - S_{15} = 1050$   
 $S_{30} = 15(2(4) + 29(3)) = 1425$   
 $a = 4 \quad d = 3$   
 $T_5 = a + 4d \quad T_7 = a + 6d$   
 $\therefore 2a + 10d = 38 \quad (\text{eq 1})$

✓✓✓

[6]

18. (a)  $r = \frac{3}{2}$   
 $-2\left(1 - \left(-\frac{3}{2}\right)^n\right) = 30$   
 $1 - \left(-\frac{3}{2}\right)^n = 15$   
 $\left(-\frac{3}{2}\right)^n = -10$   
 $n = 9.003$   
10 terms

✓✓✓

[6]

19. (a)  $\text{Area of shaded part} = \text{triangle} - \text{sector}$   
 $= 2\sqrt{3} - \frac{3}{2\pi} = 1.37 \text{ units}^2$   
 $\text{units}^2$  (3 sig fig)

✓✓✓

[6]

20. (a)  $\therefore \text{Sum of next 15 terms} = S_{30} - S_{15} = 1050$   
 $S_{30} = 15(2(4) + 29(3)) = 1425$   
 $a = 4 \quad d = 3$   
 $T_5 = a + 4d \quad T_7 = a + 6d$   
 $\therefore 2a + 10d = 38 \quad (\text{eq 1})$

✓✓✓

[6]

21. (a)  $y = 16 - 6 - w$   
 $(ii) y = 36 + w^2 - 12w \cos y$   
 $(iii) (10 - w)^2 = 36 + w^2 - 12w \cos y$   
 $64 - 20w = -12w \cos y$   
 $\therefore \cos y = \frac{5w - 16}{3w}$

✓✓✓

[6]

22.  $y = 3x^2 - 12x + k$   
 $b^2 - 4ac = 0 \text{ for one solution}$   
 $k = 144 - 4(3)k = 0$

✓✓✓

[6]

23. (a)  $W = W_0(1.085)^t$   
 $R = R_0(0.95)^t$   
 $10W_0 = R_0$   
 $10W_0 = 1.085W_0$   
 $W_0 = 985$

✓✓✓

[6]

24. (a)  $r = -\frac{3}{2}$   
 $1 - \left(-\frac{3}{2}\right)^n = 30$   
 $\left(-\frac{3}{2}\right)^n = -10$   
 $n = 9.003$   
10 terms

✓✓✓

[6]

13. (a) 5.1 seconds  
 $v(t) = -9.8t + 25$
- (b)  $v(0) = 25$
- (c) Maximum height is 31.89 m when  $t = 2.55$  sec  
 $\frac{31.89 \times 2}{2.55 \times 2} = 12.5$  m/s

14. (a)  $f(x) = (2+x)^2 = 4 + 4x + x^2$   
 $\lim_{h \rightarrow 0} \frac{(2+x+h)^2 - (2+x)^2}{h} = 2x + 4$

(b) (i) -7.5  
(ii) 18

(c)  $p'(x) = 3x^2 - 3a$   
 $0 = 3(\sqrt{2})^2 - 3a$   
 $a = 2$   
 $-\sqrt{2} = (\sqrt{2})^3 - 3(2)(\sqrt{2}) + b$   
 $b = 3\sqrt{2}$

15. (a)  $3x^2 - \frac{4}{x^2} - 11 = 0$   
 $x = -2$  or  $x = 2$

(b)

$y'$	+	0	-	0	+
$y$	↑	-	↓	-	↑


$x = -2$  Maximum

$x = 2$  Minimum

(c)  $y = x^3 - 11x + \frac{4}{x} + c$

$c = 13$

$y = x^3 - 11x + \frac{4}{x} + 13$



✓

[5]

✓✓

[7]

✓

✓

✓

✓

✓

✓

✓

✓

✓

✓✓

✓✓

✓

✓

✓

✓

✓✓✓

[10]

16. (a) (i)  $T_{100} = 23 + (99)(9)$   
 $= 914$

(ii)  $357980 = \frac{n}{2}(23 + 2534)$   
 $n = 280$

(b) (i)  $T_{n+1} = \frac{1}{4}T_n$   $T_1 = 12$   
 $S_\infty = \frac{12}{1 - \frac{1}{4}} = 16$

(ii) ✓✓

17. (a) 92°C (initial temp of tea)  
22°C (room temp)

(b) After 4.12 mins and before 7.45 mins  
 $4.12 \leq t \leq 7.45$

(c) Horizontal asymptote  $y = 22$   
The tea will cool at a decreasing rate as it approaches room temperature which is 22°.

18. (a)  $y = \frac{2}{x-3} + 2$   
(b)  $y = -3\sqrt{x+4}$   
(c)  $y = \left(\frac{1}{2}\right)^x - 4$

19. (a) 0.2  
(b) 0.5  
(c) 0.3  
(d)  $\Pr(X \cap Y) = \Pr(X) \times \Pr(Y)$  if independent  
 $\Pr(X \cup Y) = \Pr(X) + \Pr(Y) - \Pr(X \cap Y)$   
 $\Pr(X) + \Pr(Y) - \Pr(X \cup Y) = \Pr(X) \times \Pr(Y)$   
Let  $\Pr(Y) = k$   
 $0.5 + k - 0.8 = 0.5k$   
 $k = 0.6 = \Pr(Y)$

20. (a) 1, 3, 7, 15, ...  
 $T_{n+1} = T_n + 2^n$   $T_1 = 1$
- (b) 20 sets

✓

✓

✓

✓

✓✓

[8]

[6]

[6]

[6]

[3]