

--	--	--

In words

Number of additional
answer booklets used
(if applicable):

Number of additional
answer booklets used
(if applicable):

1

This Question/Answer booklet

To be provided by the candidate

Special items: nil

No other items may be taken into the examination room. It is **your** responsibility to ensure that you do not have any unauthorised material. If you have any unauthorised material with you, hand it to the supervisor **before** reading any further.

3-SST-100NS

Structure of this paper

Section	Number of questions available	Number of questions to be answered	Working time (minutes)	Marks available	Percentage of examination
Section One: Calculator-free	8	8	50	52	35
Section Two: Calculator-assumed	13	13	100	98	65
Total					100

CALCULATOR-FREE

11

METHODS UNIT 3

Supplementary page

Question number: _____

- Question 8**
- (a) Determine an expression for $\frac{d}{dt} 6t \cos\left(\frac{\pi}{6}\right)$. (2 marks)

- The volume of water in a tank, v litres, is changing at a rate given by $v'(t) = m \sin\left(\frac{\pi}{6}\right)$, where t is the time in hours. The rate of change is shown in the graph below.
-
- (b) Using the result from part (a) or otherwise, determine the change in volume of water in the tank between $t = 0$ and $t = 12$ hours. (5 marks)

- Section One: Calculator-free** 35% (52 Marks)
- This section has **eight** questions. Answer **all** questions. Write your answers in the spaces provided.
- Working time: 50 minutes.
-
- Question 1** (5 marks)
- Determine the area bounded by the line $y = -2x$ and the parabola $y = x^2 - 6x$.

Question 2**(5 marks)**

A curve, defined for $x > 0$, passes through the point $P(2, 3)$ and its gradient is given by

$$\frac{dy}{dx} = 6x^2 - \frac{4}{x^2} - 23$$

- (a) Verify that P is a stationary point, determine the value of the second derivative at P and hence describe the nature of the stationary point. (3 marks)

- (b) Determine the equation of the curve. (2 marks)

Question 7**(8 marks)**

Initially, particle P is stationary and at the origin. Particle P moves in a straight line so that at time t seconds its acceleration $a \text{ cms}^{-2}$ is given by $a = 8 - 3\sqrt{t}$ where $t \geq 0$.

- (a) Determine the speed of P after 1 second. (3 marks)

- (b) Determine the speed of P when it returns to the origin. (5 marks)

Question 6

(5 marks)

The graph of $y=f(x)$ has a stationary point at $(-1,2)$ and $f'(x)=ax^2+4x+6$, where a is a constant.

Determine the interval over which $f'(x)>0$ and $f''(x)<0$.

Question 3

(7 marks)

A bag contains 40 counters, 15 marked with 0 and the remainder marked with 1. The random variable X is the number on a randomly selected counter from the bag.

(a) Explain why X is a Bernoulli random variable and determine the mean and variance of X .
(3 marks)

Each of the 32 students in a class randomly select a counter from the bag, note the number on the counter and then replace it back in the bag. The random variable Y is the number of students in the class who select a counter marked with 0.

(b) Define the distribution of Y and determine the mean and variance of Y .
(3 marks)

(c) Explain why it is important that the students replace their counters for the distribution of Y in part (b) to be valid.
(1 mark)

Question 4**(8 marks)**

Determine

(a) $f'(x)$ when $f(x) = \sqrt{4x-3}$.

(2 marks)

(b) $\frac{d}{d\theta}(\theta^3 e^{4\theta})$ when $\theta = 2$.

(3 marks)

(c) $f\left(\frac{\pi}{4}\right)$ when $f(t) = \frac{1+\cos t}{\sin t}$.

(3 marks)

Question 5**(7 marks)**Functions f and g are such that

$$\begin{aligned} f(4) &= 2, f'(x) = 18(3x-10)^{-2} \\ g(-4) &= 2, g'(x) = 18(3x+10)^{-2} \end{aligned}$$

(a) Determine $f(6)$.

(3 marks)

(b) Use the increments formula to determine an approximation for $g(-3.98)$.

(3 marks)

(c) Briefly discuss whether using the information given about f and the increments formula would yield a reasonable approximation for $f(6)$. (1 mark)