

MATHEMATICS METHODS

Calculator-free

ATAR course examination 2016

Marking Key

Marking keys are an explicit statement about what the examining panel expect of candidates when they respond to particular examination items. They help ensure a consistent interpretation of the criteria that guide the awarding of marks.

Question 1

(5 marks)

- (a) Given that $\log_8 x = 2$ and $\log_2 y = 5$, evaluate $x - y$. (2 marks)

Solution
$8^2 - 2^5 = 32$
Specific behaviours
<ul style="list-style-type: none"> ✓ determines x and y ✓ recognises the inverse relationship between logarithms and exponentials

- (b) Express y in terms of x given that $\log_2(x + y) + 2 = \log_2(x - 2y)$. (3 marks)

Solution
$\log_2(x + y) + 2 = \log_2(x - 2y)$ $\log_2(x + y) + \log_2 4 = \log_2(x - 2y)$ $\log_2(4(x + y)) = \log_2(x - 2y)$ $4(x + y) = (x - 2y)$ $4x + 4y = x - 2y$ $6y = -3x$ $y = \frac{-1}{2}x$
Specific behaviours
<ul style="list-style-type: none"> ✓ expresses all terms as logarithms ✓ uses log laws to combine terms ✓ expresses y in terms of x

Question 2

(5 marks)

(a)

Determine $\frac{dx}{d}\left(2xe^{2x}\right)$.

(2 marks)

Solution
$\frac{d}{d}\left(2xe^{2x}\right)=2x\left(2e^{2x}\right)+e^{2x}(2)$ $=2\left(2x+1\right)e^{2x}$
Specific behaviours
✓ uses product rule ✓ differentiates exponential term

(b)

Use your answer in part (a) to determine $\int 4xe^{2x}dx$.

(3 marks)

Solution
$\frac{d}{d}\left(2xe^{2x}\right)=\left(4xe^{2x}\right)+e^{2x}(2)$ $\int \frac{dx}{d}\left(2xe^{2x}\right)dx=\int 4xe^{2x}dx+\int 2e^{2x}dx$ $2xe^{2x}=\int 4xe^{2x}dx+e^{2x}$ $\int 4xe^{2x}dx=\left(2x-1\right)e^{2x}+c$
Specific behaviours
✓ uses linearity of anti-differentiation ✓ uses fundamental theorem ✓ obtains an expression for required integral with a constant

Question 3

(7 marks)

Consider the function $f(x) = \frac{(x-1)^2}{e^x}$.

- (a) Show that the first derivative is $f'(x) = \frac{-x^2 + 4x - 3}{e^x}$. (2 marks)

Solution
$f'(x) = \frac{e^x 2(x-1) - e^x (x-1)^2}{e^{2x}}$ $= \frac{e^x (x-1)(2-x+1)}{e^{2x}}$ $= \frac{-(x-1)(x-3)}{e^x}$ $= \frac{-x^2 + 4x - 3}{e^x}$
Specific behaviours
<ul style="list-style-type: none"> ✓ uses quotient rule ✓ simplifies expression

- (b) Use your result from part (a) to explain why there are stationary points at $x = 1$ and $x = 3$. (2 marks)

Solution
$f'(x) = \frac{-(x-1)(x-3)}{e^x}$ $f'(1) = 0 = f'(3)$
Specific behaviours
<ul style="list-style-type: none"> ✓ identifies stationary points as $f'(x) = 0$ ✓ shows that this is true for $x = 1, 3$

This document – apart from any third party copyright material contained in it – may be freely copied, or communicated on an intranet, for non-commercial purposes in educational institutions, provided that it is not changed and that the School Curriculum and Standards Authority is acknowledged as the copyright owner, and that the Authority's moral rights are not infringed.

Copying or communication for any other purpose can be done only within the terms of the *Copyright Act 1968* or with prior written permission of the School Curriculum and Standards Authority. Copying or communication of any third party copyright material can be done only within the terms of the *Copyright Act 1968* or with permission of the copyright owners.

Any content in this document that has been derived from the Australian Curriculum may be used under the terms of the Creative Commons Attribution-NonCommercial 3.0 Australia licence.

Published by the School Curriculum and Standards Authority of Western Australia
303 Sevenoaks Street
CANNINGTON WA 6107

It can be shown that the second derivative is $f''(x) = \frac{e^x}{x^2 - 6x + 7}$.

(c) Use the second derivative to describe the type of stationary points at $x = 1$ and $x = 3$. (3 marks)

Solution	
$f''(x) = \frac{e^x}{x^2 - 6x + 7}$	
$f''(1) = \frac{e}{2}$	
$f''(3) = \frac{e}{-2}$	
when $x = 1$ $f'' > 0$ hence local minimum	
when $x = 3$ $f'' < 0$ hence local maximum	
Specific behaviours	
✓ evaluates second derivatives for $x = 1$ and $x = 3$	
✓ uses sign to determine nature	
✓ states nature for each stationary point	

See next page

(b) Using calculus, determine the value of θ that maximises the area A of the inscribed triangle. State this area in terms of r exactly. Justify your answer.
(Hint: you may need the identity $\sin^2 x = 1 - \cos^2 x$ in your working.) (5 marks)

Solution	
$A = r^2 \sin \theta (1 + \cos \theta)$	
$\frac{dA}{d\theta} = r^2 [\sin \theta (1 + \cos \theta) + \cos \theta (\sin \theta)]$	
$\frac{dA}{d\theta} = r^2 [\sin \theta + \cos \theta \sin \theta + \cos^2 \theta]$	
$\frac{dA}{d\theta} = r^2 [\cos \theta + \cos^2 \theta + \sin^2 \theta]$	
$\frac{dA}{d\theta} = r^2 [\cos \theta + 1]$	
$\frac{dA}{d\theta} = r^2 [2 \cos \theta + 1] = 0$	
$\cos \theta = -\frac{1}{2}, \theta = \frac{2\pi}{3}, \cos \theta \neq -1, 0 < \theta < \pi$	
$A = r^2 \sin \theta (1 + \cos \theta) = r^2 \left(\frac{\sqrt{3}}{2} \right) \left(1 + \frac{1}{2} \right) = \frac{3\sqrt{3}}{4} r^2$	
Specific behaviours	
✓ differentiates area with respect to θ using calculus	
✓ equated derivative to zero to solve for optimal value	
✓ rearranges derivative to allow solving for θ exactly	
✓ solves for $0 < \theta < \pi$ allowing for one solution only	
✓ states exact area for this optimal value	

End of questions

Question 4

(8 marks)

The displacement x micrometres at time t seconds of a magnetic particle on a long straight superconductor is given by the rule $x = 5 \sin 3t$.

- (a) Determine the velocity of the particle when $t = \frac{\pi}{2}$. (3 marks)

Solution
$x = 5 \sin 3t$ $v = \frac{dx}{dt} = 15 \cos 3t$ $v\left(\frac{\pi}{2}\right) = 15 \cos \frac{3\pi}{2} = 0$ Velocity = 0 micrometres/second
Specific behaviours
✓ differentiates to determine velocity ✓ uses chain rule ✓ evaluates velocity at $t = \frac{\pi}{2}$

- (b) Determine the rate of change of the velocity when $t = \frac{\pi}{2}$. (3 marks)

Solution
$\frac{dv}{dt} = \frac{d}{dt}(15 \cos 3t)$ $= -45 \sin 3t$ $\frac{dv}{dt} = 45$, $t = \frac{\pi}{2}$ Rate of change of velocity = 45 micrometres/second squared
Specific behaviours
✓ recognises $\frac{dv}{dt}$ as rate of change ✓ differentiates velocity ✓ evaluates rate at $t = \frac{\pi}{2}$


See next page

Question 8

(7 marks)

An isosceles triangle $\triangle PQR$ is inscribed inside a circle of fixed radius r and centre O . Let θ be defined as in the diagram below.

- (a) Show that the area A of the triangle $\triangle PQR$ is given by $A = r^2 \sin \theta (1 + \cos \theta)$. (2 marks)

Solution

Specific behaviours
✓ determines an expression of height in terms of r, θ ✓ determines an expression for base in terms of r, θ

Let v = velocity of the particle at t seconds.

(c) Determine $\int_{\frac{\pi}{2}}^0 \frac{dv}{dt} dt$. (2 marks)

Solution
$\int_{\frac{\pi}{2}}^0 \frac{dv}{dt} dt = v \left(\frac{\pi}{2} \right) - v(0)$ $= 0 - 15$ $= -15$ <p>Integral = -15 micrometres/second</p>
Specific behaviours
✓ uses fundamental theorem ✓ subtracts velocities at the two limits

Question 7 (7 marks)

Consider the graph $y = f(x)$. Both arcs have a radius of four units.

Using the graph of $y = f(x)$, $x \geq 0$, evaluate exactly the following integrals.

(a) $\int_{12}^0 f(x) dx$ (3 marks)

Solution
$36 + \frac{\pi 4^2}{4} + 4 \times 2 + \frac{1}{2} 2^2 = 46 + 4\pi$
Specific behaviours
✓ determines areas of two rectangles ✓ determines area of triangle and sector ✓ adds areas together

(b) $\int_{18}^0 f(x) dx$ (2 marks)

Solution
$46 + 4\pi - \left[\frac{1}{2} 2^2 + \left(4 \times 6 - \frac{\pi 4^2}{4} \right) \right] = 20 + 8\pi$
Specific behaviours
✓ determines area under axis ✓ uses signed areas to find net result

(c) Determine the value of the constant α such that $\int_{\alpha}^0 f(x) dx = 0$. There is no need to simplify your answer. (2 marks)

Solution
$6(\alpha - 18) + 26 - 4\pi = 46 + 4\pi$ $6(\alpha - 18) = (20 + 8\pi)$ $\alpha = \frac{6}{(20 + 8\pi)} + 18$
Specific behaviours
✓ determines a value so that signed areas balance ✓ derives an expression for α

Question 5


(6 marks)

Consider the graph of $y = f(x)$ below.

Let $A(x)$ be defined by the integral $A(x) = \int_{-1}^x f(t) dt$ for $-1 \leq x \leq 6$.

It is known that $A(2) = 15$, $A(5) = 0$ and $A(6) = 8$.

Sketch on the axes below the function $A(x)$ for $-1 \leq x \leq 6$ labelling clearly key features such as x intercepts, turning points and inflection points if any.



Question 6

(4 marks)

The graphs $y = 6 - 2e^{x-4}$ and $y = -\frac{1}{4}x + 5$ intersect at $x = 4$ for $x \geq 0$.

Determine the exact area between $y = 6 - 2e^{x-4}$, $y = -\frac{1}{4}x + 5$ and the y axis for $x \geq 0$.

