



**PERTH MODERN SCHOOL**  
Exceptional schooling. Exceptional students.  
**Independent Public School**

## Course      Methods test 4      Year 12

Student name: \_\_\_\_\_ Teacher name: \_\_\_\_\_

**Task type:**                      **Response**

**Time allowed for this task:**   40   mins

**Number of questions:**          6  

**Materials required:** Calculator with CAS capability (to be provided by the student)

Standard items:                Pens (blue/black preferred), pencils (including coloured),  
sharpener, correction fluid/tape, eraser, ruler, highlighters

Special items:                 Drawing instruments, templates, notes on one unfolded sheet  
of  
A4 paper, and up to three calculators approved for use in the  
WACE examinations

**Marks available:**         45  marks

**Task weighting:**         10 %


**Formula sheet provided:** Yes

**Note: All part questions worth more than 2 marks require working to obtain full marks**

---


## Q1 (3 marks)

Mary who goes to Perth Modern scored 67% in her exam which had a mean of 65% and a standard deviation of 18%. John who goes to Rossmoyne scored 62% in an exam with a mean of 58% and a standard deviation of 12%. Using standard scores, explain who had the better result.


Solution	
	
John has the higher z score hence better result.	
Specific behaviours	
P shows calculation for z scores P states both z scores P states that John had the better result	

## Q2 (3 marks)

Given that  $X \sim N(62, 196)$  and that  $P(k \leq x \leq 73) = 0.41$ , determine the value of  $k$  to two decimal places.



### Solution



Accept below as working but not recommended in WACE!!!

$\text{InvNormCDF}(L, 0.37398, 14, 62) = 57.50$

$k = 57.50$

### Specific behaviours

P determines prob less than 73

P subtracts to find area below k

P determines value of k to **2 dp**


(Note- max 2 marks if working not shown)

Q3 (10 marks)


The lengths of telephone calls,  $X$  in minutes, in Perth are Normally distributed with  $X \sim N(15, 3^2)$ . Calls that are longer than 17 mins are charged at double rate and are known as SUPER calls.

a) Determine the proportion of calls that are between 11 and 17 mins.


(2 marks)

Solution	
	
Specific behaviours	
<p>P uses correct parameters</p> <p>P states prob</p>	


- b) Given that a call is less than 13 mins, determine the probability that the call is greater than 9 mins. (3 marks)

Solution	
$\frac{P(9 \leq X \leq 13)}{P(X \leq 13)}$ 	
Specific behaviours	
<p>P uses conditional prob formula</p> <p>P shows calculation for numerator and denominator</p> <p>P states prob</p>	

- c) If the length of the call is in the lower 20% of calls, the call is not charged at all. Determine the longest length of call that is not charged to the nearest second.  
(2 marks)

Solution

Longest call 12 mins 29 secs (accept 12 mins 28 secs)
Specific behaviours
P uses correct parameters P states to nearest second (do not accept decimal mins)

- d) In Sydney the calls are also Normally distributed with a mean of 19 mins. If 25% of all calls are greater than 24 mins, determine the standard deviation to two decimal places. (3 marks)

Solution

Specific behaviours
P solves for z score P sets up equation for standard deviation P solves for standard deviation

Q4 (15 marks)

In a particular city it is estimated that 30% of the population have blue eyes. A sample of 250 people is chosen and the number of blue eyed people are counted. The statistician will assume a Binomial distribution.


- a) State two reasons why the statistician believes that a Binomial distribution is appropriate.

(2 marks)

Solution
The probability is independent of other people.
The probability is fixed.
Specific behaviours
P states one reason P states two reasons

- b) Determine the probability that at least 75 people of the sample have blue eyes.

(2 marks)

Solution

Specific behaviours
P uses Binomial with correct parameters P states prob (Do not accept Normal-zero)

For each sample of 250 people the number of blue eyed people are counted and a sample proportion is calculated.


- c) State the **approximate** distribution of these sample proportions including all features.

(2 marks)

Solution
Normal Distribution Mean = 0.3 Standard deviation = 0.0290

<b>Specific behaviours</b>
P states Normal P states mean and approx. st dev

- d) Determine the approximate probability that a sample proportion will be greater than 0.35.  
(2 marks)

<b>Solution</b>

<b>Specific behaviours</b>
P uses Normal with correct parameters P states prob

- e) One sample of 250 people had 65 with blue eyes. Determine a 90% confidence interval for the population proportion to 3 decimal places.  
(3 marks)

<b>Solution</b>
$\frac{65}{250} \pm 1.645 \sqrt{\frac{\frac{65}{250}(1 - \frac{65}{250})}{250}}$ $0.214 \leq p \leq 0.306$
<b>Specific behaviours</b>
P states z score P states interval P rounds to 3 dp

- f) Another three samples of 250 people were taken and the number of blue eyed people were counted. See below. One of these samples is not from the city above. Explain which sample it is and justify your answer using confidence intervals

(4 marks)

Sample	Number of blue eyed people
A	55
B	72
C	41

Solution
<p><math>A: 0.177 - 0.263</math>  <math>B: 0.241 - 0.335</math>  <math>C: 0.126 - 0.203</math>  <math>e: 0.214 - 0.306</math></p> <p>Most probably Sample C as the confidence interval does not overlap with interval first calculated.  OR  We simply do not know whether any of the above intervals contain the true value of <math>p</math> as not all confidence intervals are expected to contain the true population <math>p</math>. Therefore no inference can be made.</p>
Specific behaviours
<p>P obtains at least one confidence interval of samples A,B&amp;C  P obtains confidence intervals for all samples  P gives an argument that mentions confidence intervals  P identifies sample C OR no inference possible</p>



Q5 (2, 2 & 3 = 7 marks)


Consider a 90% confidence interval that has a margin of error of  $a$  units. Assume that the sample proportion does not change.

Answer in terms of  $a$ .


- a) Determine the new margin of error if the sample size is nine times the original.

Solution
$M \propto \frac{1}{\sqrt{n}} = \frac{1}{3}a$
Specific behaviours
P uses inverse proportion with square root P states in terms of $a$

- b) Determine the new margin of error if the sample size is unchanged but the confidence is decreased to 75%. (2 decimal places).

Solution

Margin = 0.70a
Specific behaviours
P determines z scores for both confidences P states in terms of $a$ (no need to round)

- c) Determine the new width of the confidence interval if a confidence of 97% is used and the sample size being one third of the original size. (2 decimal places)


Solution
 <p> <math display="block">\text{invNormCDf}("C", 0.9, 1, 0) = -1.644853627</math> <math display="block">\text{invNormCDf}("C", 0.97, 1, 0) = -2.170090378</math> <math display="block">\frac{-2.170090378}{-1.644853627} \sqrt{3} = 2.285131473</math> </p> <p>New width = double of 2.285a=4.57a</p>
Specific behaviours
<p>P determines all z scores</p> <p>P divides z scores and multiplies by root 3</p> <p>P doubles and states in terms of a</p>

Q6 (7 marks)


A traffic consultant has been hired to determine the proportion of people who drive to work.

- a) A confidence interval of 82% will be used with an error within 0.015. Determine the minimum sample size that should be selected. (3 marks)

Solution
----------

	
n= 1998	Accept 1996-1999 due to various rounding of z score.
<b>Specific behaviours</b>	
P sets up equation for n using p=0.5 P solves for decimal value P round up	

- b) If the maximum sample size is 720 people, determine the maximum error in estimating a 95% confidence interval. (2 marks)

<b>Solution</b>	
	
<b>Specific behaviours</b>	
P determines z score P determines error (no need to round)	

- c) The consultant will stand outside a train station and will interview every fourth person who enters. Describe two sources of bias with this sampling scheme. (explain). (2 marks)

<b>Solution</b>
Location would favour those who do not drive to work  Every fourth is not random selection
<b>Specific behaviours</b>
P identifies one source P two sources

**Extra working space**