



CHEMISTRY Stage 3

Hale School Semester Two Examination Sample

Write your name below:

Soluhais

For Examiners only	
Part 1	
Part 2	
Part 3	
Total	

Time allowed for this paper

Reading time before commencing work: Ten minutes

Working time for paper: Three hours

Materials required/recommended for this paper

To be provided by the supervisor

This Question/Answer Booklet

Multiple-choice Answer Sheet

Chemistry Data Sheet

To be provided by the candidate

Standard items: pens, pencils, eraser, correction fluid, ruler, highlighters

Special items: non-programmable calculators satisfying the conditions set out by the Curriculum Council for this course

Important note to candidates

No other items may be taken into the examination room. It is your responsibility to ensure that you do not have any unauthorised notes or other items of a non-personal nature in the examination room. If you have any unauthorised material with you, hand it to the supervisor before reading any further.

Structure of this paper

Section	Number of questions available	Number of questions to be answered	Suggested working time (minutes)	Marks available	Percentage of exam
Section One: Multiple-choice	25	25	50	25	25
Section Two: Short answer	12	12	60	70	35
Section Three: Extended answer	5	5	70	80	40
					100

Instructions to candidates

- The rules for the conduct of Curriculum Council examinations are detailed in the *Student Information Handbook*. Sitting this examination implies that you agree to abide by these rules.
- Answer the questions according to the following instructions:

Section One

Answer **all** questions, using a 2B, B or HB pencil, on the separate Multiple Choice Answer Sheet provided. Do not use a ball point or ink pen.

Section Two

Answer in the spaces provided in this Question/Answer Booklet.

Section Three

Write your answers in the Section 3 Question/Answer Booklet which is provided.

- A blue or black ball point or ink pen should be used. Marks for answers in pencil (besides section 1) can't be contested.
- For full marks, chemical equations should refer only to those species consumed in the reaction and the new species produced. These species may be **ions** [for example $\text{Ag}_{(\text{aq})}^+$], **molecules** [for example $\text{NH}_{3(\text{g})}$, $\text{NH}_{3(\text{aq})}$, $\text{CH}_3\text{COOH}_{(\text{l})}$, $\text{CH}_3\text{COOH}_{(\text{aq})}$] or **solids** [for example $\text{BaSO}_{4(\text{s})}$, $\text{Cu}_{(\text{s})}$, $\text{Na}_2\text{SO}_{4(\text{s})}$]

See next page

Section One: Multiple-choice**25% (25 Marks)**

This section has **25** questions. Answer **all** questions on the separate Multiple-choice Answer Sheet provided. For each question shade the box to indicate your answer. Use only a blue or black pen to shade the boxes. If you make a mistake, place a cross through that square, do not erase or use correction fluid, and shade your new answer. Marks will not be deducted for incorrect answers. No marks will be given if more than one answer is completed for any question.

Suggested working time for this section is 50 minutes.

1. Which of the following best describes the molecular shape and molecular polarity of a chloroform molecule whose formula is CHBr_3 ?

- (a) pyramidal, non polar
- (b) tetrahedral, non polar
- (c) pyramidal, polar
- ~~(d) tetrahedral, polar~~

2. An element X has the following five successive ionisation energies (in kJ mol^{-1})

680	1600	8000	11600	14500
-----	------	------	-------	-------

What would be the formula of the compound formed when “X” reacts with oxygen ?

- (a) X_2O
- ~~(b) XO~~
- (c) X_2O_3
- (d) XO_2

3. When 1.0 mol L^{-1} solutions of the following are mixed, which combinations will result in the formation of precipitates?

- i) $\text{Ba}(\text{NO}_3)_2$ and HCl
- ii) $\text{Ca}(\text{NO}_3)_2$ and Na_2CO_3
- iii) $\text{Cu}(\text{NO}_3)_2$ and KOH
- iv) $\text{Pb}(\text{NO}_3)_2$ and H_2SO_4

- (a) i), ii) and iii) only
- (b) ii) and iii) only
- (c) i), ii), iii) and iv)
- ~~(d) ii), iii) and iv) only~~


See next page

4. The conjugate base of the acid HCrO_4^- is:
- (a) H_2CrO_4
 - (b) H_2CrO_4^-
 - ~~(c) CrO_4^{2-}~~
 - (d) CrO_4^-
5. Which of the following physical properties **decrease** with increasing atomic number for both the alkali metals and the halogens?
- I. Atomic radius
 - II. Ionization energy
 - III. Melting point
- (a) I only
 - ~~(b) II only~~
 - (c) III only
 - (d) I and III only
6. Which of the following equations represents a redox equation? (States not included)
- (a) $\text{NaOH} + \text{HNO}_3 \rightarrow \text{NaNO}_3 + \text{H}_2\text{O}$
 - ~~(b) $2\text{AgNO}_3 + \text{Cu} \rightarrow 2\text{Ag} + \text{Cu}(\text{NO}_3)_2$~~
 - (c) $\text{H}_2\text{SO}_4 + 2\text{KOH} \rightarrow \text{K}_2\text{SO}_4 + 2\text{H}_2\text{O}$
 - (d) $\text{CaCl}_2 + \text{Ba}(\text{OH})_2 \rightarrow \text{Ca}(\text{OH})_2 + \text{BaCl}_2$
7. Which one of the following solids contains covalent bonds only?
- ~~(a) SiO_2~~
 - (b) MgO
 - (c) NH_4Br
 - (d) Ne
8. If the pH of a solution changes from 2 to 4, then the hydronium ion concentration
- (a) is doubled.
 - (b) is halved.
 - (c) increases by a factor of 100.
 - ~~(d) decreases by a factor of 100.~~

9. A crystal of iodine, I_2 , produces a purple vapour when gently heated. Which pair of statements correctly describes this process?

	Type of bond broken	Formula of purple species
(a)	covalent	I
(b)	covalent	I_2
(c)	dispersion forces	I_2
(d)	dipole-dipole	I_2

10. Household bleach contains sodium hypochlorite, NaClO, as the active ingredient. The concentration of NaClO in the bleach can be determined by reacting a known amount with aqueous hydrogen peroxide, H_2O_2 , according to the equation:




When 25.0 mL of bleach is treated with an excess of aqueous H_2O_2 , 0.0350 mol of oxygen gas is given off.

What is the concentration of NaClO in the bleach?

- ~~(a)~~ 1.40 mol L⁻¹
 (b) 0.700 mol L⁻¹
 (c) 0.875 mol L⁻¹
 (d) 8.75×10^{-4} mol L⁻¹

11. In the contact process reaction:



If the equilibrium system **temperature** is increased, what effect will this have on the equilibrium constant, K, and the yield?

	Equilibrium constant, K	Yield increase
(a)	decrease	products
(b)	decrease	reactants
(c)	increase	products
(d)	increase	reactants


12. Deposits of ammonium compounds, including ammonium sulfate, have been discovered in areas of high atmospheric pollution. A chemical reaction believed to occur is:




What does **not** occur in this reaction?

- (a) acid/base neutralisation
 - (b) coordinate (dative) bond formation
 - ~~(c)~~ oxidation/reduction
 - (d) ionic bond formation
13. Galvanic cells are used as portable sources of electrical energy. One common cell is the rechargeable nickel-cadmium cell.

Nickel–Cadmium Cell



The net equation representing the discharge of the nickel-cadmium cell is:




The reaction at the **anode** during the discharge of the cell is:

- ~~(a)~~ $\text{Cd}(\text{s}) + 2\text{OH}^-(\text{aq}) \rightarrow \text{Cd}(\text{OH})_2(\text{s}) + 2\text{e}^-$
- (b) $\text{Cd}(\text{s}) + 2\text{OH}^-(\text{aq}) + 2\text{e}^- \rightarrow \text{Cd}(\text{OH})_2(\text{s})$
- (c) $\text{NiO}_2(\text{s}) + 2\text{H}_2\text{O}(\text{l}) + 2\text{e}^- \rightarrow \text{Ni}(\text{OH})_2(\text{s}) + 2\text{OH}^-(\text{aq})$
- (d) $\text{NiO}_2(\text{s}) + 2\text{H}_2\text{O}(\text{l}) \rightarrow \text{Ni}(\text{OH})_2(\text{s}) + 2\text{OH}^-(\text{aq}) + 2\text{e}^-$

See next page

14. Which one of the following has the same electronic arrangement as Li^+ ?
- (a) Na^+
~~(b) Be^{2+}~~
 (c) F^-
 (d) Ne
15. The largest mass of silver chloride is precipitated when an excess of silver nitrate solution is added to:
- (a) 25.0 mL of a 0.800 mol L^{-1} solution of hydrochloric acid.
~~(b) 30.0 mL of a 0.300 mol L^{-1} solution of iron(III) chloride.~~
 (c) 50.0 mL of a 0.200 mol L^{-1} solution of magnesium chloride.
 (d) 50.0 mL of a 0.500 mol L^{-1} solution of sodium chloride.
16. The IUPAC name for the structure below is:



- ~~(a) 2,2,5-trimethylheptane~~
 (b) 3,6,6-trimethylheptane
 (c) 2-ethyl-5,5-dimethylhexane
 (d) 5-ethyl-2,2-dimethylhexane

17. Which one of the following species does **not** have eight valence electrons surrounding the central atom?


(a) CHCl_3 molecule

~~(b)~~ NO_2 molecule

(c) NH_4^+ ion

(d) OF_2 molecule

18. Select, from the list below, the compound that can be polymerised to give:







~~(a)~~ 2-methylbut-1-ene

(b) 2-methylbut-2-ene

(c) pent-2-ene

(d) pent-1-ene

19. Which of the following rows identifies the structural diagram and the corresponding IUPAC name of the compound with the chemical formula, C_8H_{16} ?

	Structural Diagram	IUPAC Name
(a)		ethylbenzene
(b)		ethylcyclohexane
(c)		cyclopentylpropane
(d)		propylcyclopentene


20. When the compounds HF, H_2O , NH_3 , and CH_4 are listed in order of increasing boiling point, which order is correct?

- (a) $CH_4 < NH_3 < H_2O < HF$
 (b) $NH_3 < CH_4 < H_2O < HF$
~~(c)~~ $CH_4 < NH_3 < HF < H_2O$
 (d) $HF < CH_4 < H_2O < NH_3$

21. The reductant that can convert $1.0\text{ M Fe}^{3+}(\text{aq})$ to $\text{Fe}^{2+}(\text{aq})$ but not $1.0\text{ M Sn}^{2+}(\text{aq})$ to $\text{Sn}(\text{aq})$, at STP is:


- ~~(a)~~ $\text{Cu}(\text{s})$
 (b) $\text{Au}(\text{s})$
 (c) $\text{Ni}(\text{s})$
 (d) $\text{HOOC}(\text{COOH})$

22. A cell was incorrectly connected, as shown below. Which statement is **incorrect**?



- (a) The anode is the zinc electrode.
- (b) There would be no electron current flow from one half cell to the other.
- ~~(c)~~ If electrodes are interchanged the cell emf (potential difference) would be -1.1 V (at 25°C).
- (d) The concentration of Cu^{2+} ions will decrease.
23. Which of the following statements is **correct**?
- (a) Covalent network solids include diamond, graphite and sulfur.
- (b) Metal solids and ionic solids exhibit non-directional interparticle bonding.
- (c) Ionic solids conduct electricity very well in the aqueous and solid states.
- ~~(d)~~ Heated covalent molecular solids tend to decompose before melting.
24. Which of the following statements about the third row of the Periodic Table is correct?
- ~~(a)~~ Elements on the right side of the row form acidic oxides, whilst those on the left side form basic oxides.
- (b) Elements on the left side of the row have a greater range of oxidation states than elements on the right side.
- (c) Elements on the right side of the row are stronger reducing agents than elements on the left side.
- (d) Electronegativity decreases across a row from left to right of the period.

25. A common painkiller has the structure:



Which of the options below best represents its characteristics?

	Type	Functional groups
(a)	aromatic	carboxyl, hydroxyl
(b)	aliphatic	hydroxyl, alkene
(c)	aromatic	hydroxyl, ester
(d)	aliphatic	carbonyl, hydroxyl

END OF SECTION 1

Section Two: Short answer

35% (70 Marks)

This section has 12 questions. Answer **all** questions. Write your answers in the space provided.

Spare pages are included at the end of this booklet. They can be used for planning your responses and/or as additional space if required to continue an answer.

- Planning: If you use the spare pages for planning, indicate this clearly at the top of the page.
- Continuing an answer: If you need to use the space to continue an answer, indicate in the original answer space where the answer is continued, i.e. give the page number. Fill in the number of the question(s) that you are continuing to answer at the top of the page.

Suggested working time for this section is 60 minutes.

Question 26


(5 marks)

A buffer solution is needed for preserving “Tango” fruit juice. A chemist at the fruit juice company prepared a benzoic acid/sodium benzoate buffer with concentrations of 0.105 mol L^{-1} $\text{C}_6\text{H}_5\text{COOH}$ and 0.125 mol L^{-1} $\text{C}_6\text{H}_5\text{COONa}$.

- (a) What is a “buffer solution”, and what is its purpose? (2 mark)

A solution of weak acid and conj. base or visa versa. Buffer keep pH relatively stable when acid or base is added.

- (b) Explain, using Le Chatelier’s principle, how this solution acts as a buffer solution. (use equations in your answer). (3 marks)



acid is added and H^+ reacts with benzoate ion to $\downarrow [\text{H}^+]$

(or reacting with OH^-)

1 mark equation - 2 mark explanation

Question 27

(2 marks)

See next page

Toluene (methyl benzene), C_7H_8 (g) is an important solvent and precursor to many other organic compounds such as trinitrotoluene (TNT). It can be produced according to the following equilibrium:



When 3.00 mol of C_7H_{14} (g) was introduced into a 1.00 L container, 1.20 mol of H_2 (g) was produced at equilibrium.

What were the equilibrium concentrations, in mol L^{-1} , of C_7H_8 and C_7H_{14} ?

$$C_7H_8 = 0.4 \text{ mol.}$$

$$C_7H_{14} = 3 - 0.4 = 2.6 \text{ mol.}$$

Question 28

(4 marks)

Write **observations** for any reactions that occur in the following procedures. In each case describe in full what you would observe, including any:

- colours
- odours
- precipitates (give the colour)
- gases evolved (give the colour or describe as colourless).

If no change is observed, you should state this.

- (a) Potassium sulfide solution is added to lead (II) nitrate solution.

(2 marks)

Observation: Two clear solutions combine to form black ppt in clear solution.

- (b) Sodium metal is added to pentanol.

(2 marks)

Observation: grey solid added to clear solution. Colourless odourless gas produced and solid dissolves to give a clear solution.

Question 29

(3 marks)

- (a) Consider the reaction half equations and then balance the following redox equation:



See next page

3


2

6

4



- (b) Re-write the redox equation, for alkaline (basic) conditions. (1 mark)




Question 30


(4 marks)

Write the equation for the reaction that occurs in each of the following procedures. If no reaction occurs, write 'no reaction'. For full marks, chemical equations should refer only to those species consumed in the reaction and the new species produced. These species may be ions [for example $\text{Ag}^+(\text{aq})$], molecules [for example $\text{NH}_3(\text{g})$, $\text{NH}_3(\text{aq})$, $\text{CH}_3\text{COOH}(\text{l})$] or solids [for example $\text{BaSO}_4(\text{s})$, $\text{Cu}(\text{s})$, $\text{Na}_2\text{CO}_3(\text{s})$].

- (a) Potassium phosphate solution is added to copper (II) sulfate solution. (2 marks)



- (b) Propene gas is shaken with hydrogen bromide gas in the presence of uV light. (2 marks)




Question 31

(7 marks)

Commercial buses, and cars, can operate using a proton exchange membrane fuel cell (PEM) to provide the required energy. They use a solid polymer sandwiched between two sheets of carbon fibre paper as an electrolyte, and porous carbon

See next page



electrodes containing a platinum catalyst. They need only hydrogen, oxygen from the air, and water to operate. They are typically fuelled with pure hydrogen supplied from storage tanks or onboard reformers. The cell operates at a temperature of around 80°C.

A Picture of the Hydrogen Fuel Cell Transit Bus



- (a) What is **one** advantage of a solid polymer membrane electrolyte over the liquid chemical electrolyte, KOH (used in other fuel cells)? (1 mark)

less mass not corrosive.
spillage

- (b) State **one** environmental advantage, and **one** sustainability advantage; of this PEM fuel cell over the use of conventional fuels such as diesel? (2 marks)


no toxic waste (1)
O₂ + H₂ renewable
diesel non-renewable → (1)

- (c) What are **two** disadvantages of the O₂/H₂ fuel cell, as energy sources, over conventional fuels such as diesel? (2 marks)

Fuel needs stored carefully
Needs const. fuel.
High cost of cell.

(any 2)

- (d) Some fuel cells use methanol as a source of hydrogen (as H⁺), which combines with oxygen to produce carbon dioxide and water. Write an overall balanced redox equation given the following skeleton equations; CH₃OH → CO₂ and O₂ → H₂O (2 marks)




See next page

Question 32

(8 marks)

On heating, a mixture of potassium and bromine react to form potassium bromide, according to the equation:



- (a) The melting points of potassium, bromine and potassium iodide are 63.3 °C, -7.2 °C and 734 °C respectively. For each of the substances state the type of **inter-particle** bonding present and the nature of the attractive forces holding each substance together. (6 marks)

Substance	Type of bonding	Nature (strength) of bonding
Potassium	metallic Dispersion	strong.
Bromine		weak
Potassium bromide	ionic	strong

- (b) Briefly explain why the melting point of bromine is much lower than that of sodium bromide. (2 marks)

less energy required to break dispersion forces.






Question 33

(15 marks)

- (a) For each species listed in the table below, draw the structure, representing all valence shell electron pairs either as : or as — **and** state or draw the shape of the molecule or ion. (9 marks)

(for example, water $\text{H}:\ddot{\text{O}}:\text{H}$ or $\text{H}-\ddot{\text{O}}-\text{H}$ or $\text{H}-\bar{\text{O}}-\text{H}$ bent)

See next page

Compound	Electron-dot structure (showing all valence shell electrons)	Shape (sketch or name)
Carbon disulfide CS_2	 	linear
Strontium nitrate $\text{Sr}(\text{NO}_3)_2$	 	3D network 2
Urea H_2NCONH_2		+planar.

- (b) Compare, and explain, the molecular polarity of carbon disulfide and urea. (6 marks)


1 mark

The two bond

area

Explanation


2 marks

Compound	Polar or non-polar	Explanation
CS ₂	NP	dipoles equal and opposite ∴ cancel each other out 
H ₂ NCONH ₂	Polar.	There is an asymmetrical e ⁻ distribution and a highly electronegative O atom

Question 34

(5 marks)

- (a) Draw and label the geometric isomeric forms of 2-pentene (pent-2-ene). (3 marks)




- (b) What chemical test could be used to distinguish between pent-2-ene and pentane? (chemical equation required) (2 marks)

Add Bromine water. Only pent-2-ene

See next page

will decolour the bromine water (1)



Question 35

(10 marks)

Industrially, calcium carbonate can be thermally decomposed (roasted) to form calcium oxide ('quicklime'). This product is used to change pH, in calcium silicate brick manufacture, in aluminium and gold production, and in the building industry (plaster).

The chemical reaction for its production is:



- (a) What is the K_{eq} expression for the reaction?






(1 mark)

$$K = [\text{CO}_2] \text{ at } 25^\circ\text{C}$$

- (b) In practice, conditions can be changed to alter the rate of reaction and the yield of product. Indicate any effects of imposed change on the system in the table below.

(9 marks)

Imposed change	Effect on forward reaction rate	Effect on reverse reaction rate	Effect on the value of K
Increase the partial pressure of the carbon dioxide	\uparrow	\uparrow	no change

The temperature is decreased			
Increase the surface area of the CaCO_3			no change

Question 36

(3 marks)

Older second hand cars can have bubbling of paint on the panels due to iron corrosion ("rusting") under the sealed paint (i.e. an anodic region). This is often noticed close to the wheel hubs or the base of doors or windows. The paint under normal circumstances provides a physical barrier to corrosion.

(a) What is the likely cause of the corrosion?

(2 marks)

'Protective' layer is broken allowing oxidation of the iron underneath

(b) Suggest a practical chemical solution to the problem.


(1 mark)

paint, seal it etc

Question 37


(4 marks)

Vinegar is about 4% by mass acetic acid and is safe to consume in foods. The same concentration sulfuric acid is not safe to consume. Explain why. Include equations.



partial dissociation

See next page



END OF SECTION 2

EXTRA WORKING SPACE

[illegible]

See next page



CHEMISTRY

Stage 3

Hale School

Semester Two Examination Sample

Write your name below:

Section Three: Extended answers

40% (80 Marks)

This section contains **five (5)** questions. You must answer **all** questions. Write your answers in the space provided in this question and answer booklet. Answer should be in blue or black pen.

Spare pages are included at the end of this booklet. They can be used for planning your responses and/or as additional space if required to continue an answer.


- Planning: If you use the spare pages for planning, indicate this clearly at the top of the page.
- Continuing an answer: If you need to use the space to continue an answer, indicate in the original answer space where the answer is continued, i.e. give the page number. Fill in the number of the question(s) that you are continuing to answer at the top of the page.

Suggested working time for this section is 70 minutes.

Question 38**(14 marks)**

A student carried out two acid-base practical investigations, at 25°C:

A titration was performed where 20.0 mL of an unknown solution was pipetted into a conical flask and titrated with another unknown solution from a burette. The pH was monitored with a pH meter, recorded with a data logger and the results displayed on a computer screen throughout the experiment. The changes in pH are shown below:



- (a) What is meant by the term “equivalence point”? (1 mark)

*When acid + base are
in stoichiometric proportions*

- (b) What was the pH of the mixture at the equivalence point? (1 mark)

7

See next page

- (c) (i) A pH meter had to be used in this investigation. Explain, why there was no suitable acid-base indicator that could be used for this titration? (1 mark)

End point is not sharp \therefore difficult to detect accurately.

- (ii) What does this tell us about the acid and base added together? (1 mark)

weak acid / weak base

- (iii) Give an example of the acid and base that could have been used to give these results? (1 mark)

Acid something weak Base something weak.

- (iv) Which species was in the conical flask? (1 mark)

The base


- (e) What was the concentration of $\text{OH}^-(\text{aq})$ (mol L^{-1}) in the mixture after 15.0 mL of titrant had been added? (2 marks)

$\text{pH} = 10 \therefore [\text{H}^+] 1 \times 10^{-10}$

$\therefore [\text{OH}^-] = 1 \times 10^{-4} \text{ mol L}^{-1}$

A commercial brand of antacid, "Easiflux", was tested to see how much acid it could neutralize. A 5.00 mL sample of the mixture was used where the active ingredient was stated to be magnesium hydroxide. The mixture was reacted with 12.90 mL of 72.9 g L^{-1} hydrochloric acid, HCl (stomach acid).

- (f) Write a balanced equation for this reaction. (1 mark)



- (g) How many moles of HCl reacted? (2 marks)

$$m(\text{HCl}) = 0.01290 \text{ L} \times 72.9 \text{ g L}^{-1}$$

$$= 0.94041 \text{ g}$$

$$n(\text{HCl}) = \frac{0.94041}{36.453} = 0.0258 \text{ mol.}$$

- (h) What mass (g) of magnesium hydroxide would react with this amount of HCl? (2 marks)

$$n(\text{Mg}(\text{OH})_2) = \frac{1}{2} n(\text{HCl}) = 0.0129 \text{ mol}$$

$$\therefore \text{mass}(\text{Mg}(\text{OH})_2) = 0.0129 \times \text{MPP} = 0.752 \text{ g.}$$

- (i) The suggested dose for bad indigestion is 800 mg of $\text{Mg}(\text{OH})_2(\text{s})$ in a 5.00 mL dose. How does the sample tested compare with this? (1 mark)

very close 752 mg compared
to 800 mg,


Question 39

(17 marks)


There are many ways of observing and measuring the rates of chemical reactions. A student investigated one reaction involving colour change; an "iodine clock" reaction.

At room temperature, a reaction occurs when potassium iodate solution is mixed with sodium hydrogen sulfate solution that contains a small amount of starch. In a laboratory, 12.00 mL of a 0.0160 M $\text{NaHSO}_3(\text{aq})$ solution containing starch were placed in each of six test tubes. Different volumes of 0.0240 M $\text{KIO}_3(\text{aq})$ and enough distilled water to maintain a constant volume were added to each test tube and the time taken for the dark-blue colour to appear was measured. The data were recorded in the table over the page.


The chemical reaction is:



- (a) Is this a redox reaction? Justify your answer using half equations and oxidation numbers (3 marks)



See next page



- (b) Describe a procedure for diluting the 500mL stock solution of 0.0240 mol L⁻¹ KIO₃ solution to give a 1.00L solution of 0.00800 mol L⁻¹ reaction mixture test solution. (2 marks)

$$c_1 V_1 = c_2 V_2 \quad V_1 (0.024) = (0.008) \times 1$$

$$V_1 = 0.33333 \text{ L.}$$

i. take 333.3 mls of stock and dilute with water to 1000 mls.

- (c) What is the independent variable for the investigation? (1 mark)

$[\text{IO}_3^-]$ solution.

- (d) What variables need to be controlled? (1 mark)


temp, mixing, conc, apparatus etc. (1 only)

- (e) Complete the table below, and graph the data of reaction rate (1/t) versus concentration of potassium iodate (mol L⁻¹). (5 marks)

Concentration of IO ₃ ⁻ (mol L ⁻¹) in reaction mixture.	Reaction time (s).	Reaction rate, 1/t (s ⁻¹).
0.00200	210	0.00476
0.00400	88	0.0114
0.00600	49	0.0204
0.00800	39	0.0256 0.0303

0.0100	33	
0.0120	27	

$[IO_3^-]$ vs rate $\frac{0.0370}{s}$



- (f) What conclusion can be drawn from the graph about the relationship between the rate of the reaction and the concentration of the potassium iodate? (1 mark)

As $[IO_3^-]$ increases so does rxn rate

- (e) Use your graph to predict the time taken for a $0.00500 \text{ mol L}^{-1}$ potassium iodate solution, at standard room temperature, to react. (1 mark)

$\therefore \frac{1}{t} = 0.015, \therefore t = 67 \text{ seconds}$

- (f) The procedure described above was repeated at a number of different temperatures between 10 °C and 70 °C and the reaction times were measured as before. Would you expect the reaction times to increase, decrease, or stay the same, as the temperature was increased? Use Collision Theory to justify your answer.

(3 marks)


increase (1)
 greater av. kinetic energy ∴ more frequent and harder collisions (1)
 ∴ increase frequency of successful collisions (1)

Question 40

(not number)

(15 marks)

Iron is one of the most abundant metals on Earth, is essential to most forms of life and to normal human physiology. Sometimes people take iron supplements. The iron content of a particular brand of iron tablets was determined by titration with a freshly standardised solution of potassium permanganate, KMnO_4 . The equation for the titration reaction is:



- (a) Why are iron tablets sometimes medically prescribed? (1 mark)

anything containing iron

- (b) What is the oxidation number of manganese in the permanganate ion? (1 mark)

+7

- (c) (i) Why must potassium permanganate solutions be standardised? (2 mark)

to ensure accurate concentration prior to using as standard.

- (ii) What reagent is used for this purpose? (1 mark)

oxalic acid

250 mL of Fe^{2+} solution was prepared from ten (10) iron tablets, each of mass 0.328 g.

- (e) Explain why additional dilute sulfuric acid must be added to the titration flask before each titration is carried out. (1 mark)

must be acidified to produce Mn^{2+} . otherwise colour change is difficult.

- (f) How was the end-point detected? (1 mark)

Purple to very pale pink

A standardised 0.0100 M potassium permanganate was used to react with 25.0 mL portions of the iron solution prepared from the ten tablets.

- (g) A number of titrations were performed and the following titre values obtained.

Titre (mL)	21.00	18.79	18.76	18.70	
------------	-------	-------	-------	-------	--

What is the average titre used? (1 mark)

$$(18.79 + 18.76 + 18.70) / 3 = 18.75 \text{ mLs}$$

- (g) What is the **concentration** (mol L^{-1}) of the Fe^{2+} solution? (3 marks)

$$n(\text{MnO}_4^-) = 0.01 \times 0.01875 = 0.0001875$$

$$\therefore n(\text{Fe}^{2+}) \text{ in } 25 \text{ mL} = 5 \times n(\text{MnO}_4^-) = 0.0001875 \times 5 = 0.0009375$$

$$\therefore [\text{Fe}^{2+}] \text{ in } 25 \text{ mLs} = \frac{0.0009375}{0.025} = 0.0375 \text{ M}$$

- (h) What is the **total mass** (mg) of iron in one tablet? (3 marks)

$$n(\text{Fe}) \sim 250 \text{ mLs} = 10 \times 0.0009375 \\ = 0.009375$$

$$\therefore \text{mass Fe} = 0.009375 \times 55.85 \\ = 0.523 \text{ g}$$

$$\text{or } 52.3 \text{ mg}$$

- (i) What is the **percentage, by mass**, of iron in each tablet? (1 mark)


$$\% \text{ Fe} = \frac{0.523}{0.328} \times 100$$

$$= 15.9\%$$

Question 41

(22 marks)

Many drugs are produced by chemically combining several molecules. A newly developed amino acid, in combination with another substance, has the potential to be a new anti-inflammatory drug. The structure of the amino acid ($M_r = 179.214$) is:



- (a) (i) Explain, why it is considered to be an amino acid? (2 marks)


contains amine and carboxylic acid functional groups

- (ii) Would you expect this molecule to be fairly soluble in water? Explain (2 marks)


*Yes (1)
Due to H-bonds at both functional groups (1)*

The amino acid above exists as a zwitterion in aqueous solution. Zwitterion is the general name given these compounds when they form an ion which has a positive charge on one atom and a negative charge on another.


- (b) Draw the structure of this **zwitterion**. (2 marks)



The anti-inflammatory drug is made by combining the amino acid, with the molecule shown below, called compound X. R represents a small side chain.



- (c) (i) Combine both structures to show the structure of the **drug**. (1 mark)



- (ii) What type of chemical linkage forms? (1 mark)

Amide or peptide link.

- (iii) What type of reaction occurs, and what small molecule is released? (2 marks)

Condensation polymerisation / H₂O.

- (d) Spectroscopy analysis showed compound X contained the elements C, H, O, and chlorine (Cl).

Experimentally, the identity of R was found by combusting 0.425 g of compound X in excess air. It was found that 0.844 g of carbon dioxide and 0.171 g of water was produced.

Further testing of a further 0.396 g sample of compound X, produced 25.7 mLs of chlorine gas (Cl_2), under conditions of 40 °C and 100 kPa.

- (i) Determine the **percentage composition**, by mass, of each element in compound X. (7 marks)

$$n(\text{C}) = n(\text{CO}_2) = \frac{0.844}{44.01} = 0.0191775$$

$$n(\text{H}) = \frac{2 \times 0.171}{18.016} = 0.018983$$


$$n(\text{Cl}) = \frac{100 \times 0.0257}{8.315(313.1)} = 9.871 \times 10^{-4}$$

$$\therefore \text{mass}(\text{Cl}) = 9.871 \times 10^{-4} \times 70.9 = 0.0699853$$

C	H	O	Cl
54.19%	4.57%	23.64%	17.67%

(ii) Determine the **empirical formula** of compound X.

(3 marks)



(iii) What is the **molar mass** of compound X?

(2 mark)


$$108.09 + 9.072 + 48 + 35.45 = 200.612$$

Question 42

(12 marks)

Butanoic acid, $\text{CH}_3\text{CH}_2\text{CH}_2\text{COOH}$, is a ubiquitous, oily, colourless substance with a diverse range of origins and uses. It occurs naturally but can be manufactured through the fermentation of sugar and starch, and then the addition of putrefying cheese, with calcium carbonate added to neutralize the acid.

- (a) Draw and name two structural isomers of $\text{C}_4\text{H}_8\text{O}_2$ (other than butanoic acid). (4 marks)




Butanoic acid is a rancid smelling substance that gives parmesan cheese its characteristic odour. It has been used as a nausea inducing repellent by anti-whaling protesters against whalers. Butanoic acid can undergo **esterification** with ethanol, $\text{CH}_3\text{CH}_2\text{OH}$, to form a much more pleasant smelling pineapple flavoured ester.

- (b) What is the common **catalyst** used for this reaction? (1 mark)

sulfuric acid.

- (c) Draw, and name the **ester** formed. (2 marks)



See next page

Name:	ethyl butanoate
-------	-----------------

The perspiration stains in clothes are partly due to the presence of butanoic acid. Soap powders form alkaline solutions, often containing sodium carbonate, which are used to neutralise this acidity.

Fats and oils can be removed from clothing by the action of soaps. Soaps are typically sodium or potassium salts of long chain fatty acids e.g. $\text{CH}_3(\text{CH}_2)_{16}\text{COONa}$.

(d) Discuss, with illustrations, the action of soaps. Use the terms:

- dispersion forces
- surfactant (or emulsifying agent)
- polar and non-polar
- hydrophobic and hydrophilic

(5 marks)

1 mark for each dot point

1 mark for ~~the~~ appropriate diagrams

[illegible]

END OF SECTION 3

[illegible]

