

Chance and data

Probability:

For any event A and its complement \bar{A} , and event B

$P(A) + P(\bar{A}) = 1$

$P(A \cup B) = P(A) + P(B) - P(A \cap B)$

$P(A \cap B) = P(A) P(B|A) = P(B) P(A|B)$

In a binomial distribution:

Mean: $\mu = np$ and standard deviation: $\sigma = \sqrt{np(1-p)}$

A confidence interval for the mean of a population is:

$$\bar{x} - z \frac{\sqrt{n}}{\sigma} \leq \mu \leq \bar{x} + z \frac{\sqrt{n}}{\sigma}$$

where μ is the population mean,

σ is the population standard deviation,

\bar{x} is the sample mean,

n is the sample size and

z is the cut-off value on the standard normal distribution corresponding to the confidence level.

Note: Any additional formulas identified by the examination panel as necessary will be included in the body of the particular question.

Copyright

© School Curriculum and Standards Authority, 2012

This document—apart from any third party copyright material contained in it—may be freely copied, or communicated on an intranet, for non-commercial purposes by educational institutions, provided that it is not changed in any way and that the School Curriculum and Standards Authority is acknowledged as the copyright owner.

Copying or communication for any other purpose can be done only within the terms of the Copyright Act or by permission of the Authority.

Copying or communication of any third party copyright material contained in this document can be done only within the terms of the Copyright Act or by permission of the copyright owners.

This document is valid for teaching and examining until 31 December 2012.

MATHEMATICS

UNITS 3C AND 3D

FORMULA SHEET

2012



School Curriculum and Standards Authority
Government of Western Australia



Number and algebra

Index laws:

For $a, b > 0$ and m, n real,

$$a^m b^n = (a b)^m \quad a^m a^n = a^{m+n} \quad (a^m)^n = a^{mn}$$

$$a^{-m} = \frac{1}{a^m} \quad \frac{a^m}{a^n} = a^{m-n} \quad a^0 = 1$$

For $a > 0$ and m an integer and n a positive integer, $a^{\frac{m}{n}} = \sqrt[n]{a^m} = (\sqrt[n]{a})^m$

If $f(x) = y$ then $f'(x) = \frac{dy}{dx}$

If $f(x) = x^n$ then $f'(x) = nx^{n-1}$

If $f(x) = e^x$ then $f'(x) = e^x$

Product rule:

If $y = f(x) g(x)$ or If $y = uv$
then $y' = f'(x) g(x) + f(x) g'(x)$ then $\frac{dy}{dx} = \frac{du}{dx} v + u \frac{dv}{dx}$

Quotient rule:

If $y = \frac{f(x)}{g(x)}$ or If $y = \frac{u}{v}$
then $y' = \frac{f'(x) g(x) - f(x) g'(x)}{(g(x))^2}$ then $\frac{dy}{dx} = \frac{\frac{du}{dx} v - u \frac{dv}{dx}}{v^2}$

Chain rule:

If $y = f(g(x))$ or If $y = f(u)$ and $u = g(x)$
then $y' = f'(g(x)) g'(x)$ then $\frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx}$

Powers: $\int x^n dx = \frac{x^{n+1}}{n+1} + c, n \neq -1$

Exponentials: $\int e^x dx = e^x + c$

Fundamental Theorem of Calculus:

$\frac{d}{dx} \int_a^x f(t) dt = f(x)$ and $\int_a^b f'(x) dx = f(b) - f(a)$

Incremental formula: $\delta y \approx \frac{dy}{dx} \delta x$

Exponential growth and decay: If $\frac{dy}{dt} = ky$, then $y = Ae^{kt}$

See next page

Space and measurement

Circle: $C = 2\pi r = \pi D$, where C is the circumference,
 r is the radius and D is the diameter
 $A = \pi r^2$, where A is the area

Triangle: $A = \frac{1}{2}bh$, where b is the base and h is the perpendicular height

Parallelogram: $A = bh$

Trapezium: $A = \frac{1}{2}(a + b)h$, where a and b are the lengths of the parallel sides
and h is the perpendicular height

Prism: $V = Ah$, where V is the volume, A is the area of the base
and h is the perpendicular height

Pyramid: $V = \frac{1}{3}Ah$

Cylinder: $S = 2\pi rh + 2\pi r^2$, where S is the total surface area
 $V = \pi r^2 h$

Cone: $S = \pi rs + \pi r^2$, where s is the slant height
 $V = \frac{1}{3}\pi r^2 h$

Sphere: $S = 4\pi r^2$
 $V = \frac{4}{3}\pi r^3$

Volume of solids of revolution:

$V = \int \pi y^2 dx$ rotated about the x -axis

$V = \int \pi x^2 dy$ rotated about the y -axis

See next page