

Copyright for test papers and marking guides remains with *West Australian Test Papers*.  
Test papers may only be reproduced within the purchasing school according to the advertised Conditions of Sale.  
Test papers should be withdrawn after use and stored securely in the school until Thursday 15th October 2020.



# **SEMESTER TWO**

## **MATHEMATICS METHODS UNITS 3 & 4**

**2020**

## **SOLUTIONS**

Calculator-free Solutions

1. (a)

$a$	-1	0	1	2
$P(A = a)$	0.2	0.3	0.4	0.1

✓✓

$$(b) \quad E(A) = (-1)(0.2) + (0)(0.3) + (1)(0.4) + (2)(0.1)$$

$$= -0.2 + 0.4 + 0.2 = 0.4 \text{ or } \$400$$

✓

✓

$$(c) \quad VAR(A) = (-1)^2(0.2) + (1)^2(0.4) + (2)^2(0.1) - (0.4)^2$$

$$= 0.2 + 0.4 + 0.4 - 0.16$$

$$= 0.84$$

✓

✓

✓

[7]

2. (a)

$m$	0	1
$P(M = m)$	$\frac{\binom{1}{0}\binom{4}{1}}{\binom{5}{1}} = \frac{4}{5}$	$\frac{1}{5}$

✓✓

(b) It has two possibilities, independent events and is a DRV

✓

$$E(M) = (0)\left(\frac{4}{5}\right) + (1)\left(\frac{1}{5}\right) = \frac{1}{5}$$

(c)

✓

$$VAR(M) = (0)^2\left(\frac{4}{5}\right) + (1)^2\left(\frac{1}{5}\right) - \left(\frac{1}{5}\right)^2 = \frac{4}{25}$$

$$\therefore SD = \frac{2}{5}$$

✓

(d)  $X \sim B(8, \frac{1}{5})$ 

✓

$$P(X = 3) = \binom{8}{3} \left(\frac{1}{5}\right)^3 \left(\frac{4}{5}\right)^5$$

(e)

✓

[7]

3. (a)  $\frac{d}{dx}[(3x^2 + 5)^4] = 4(3x^2 + 5)^3(6x)$  ✓✓
- (b)  $g'(x) = e^{3x}(3)\cos(2x) + e^{3x}(-\sin(2x))(2)$  ✓
- $g'(\pi) = 3e^{3\pi}\cos(2\pi) + 2e^{3\pi}(-2\sin(2\pi))$  ✓
- $\therefore = 3e^{3\pi}$  ✓
- (c) (i)  $-2\cos(3x) + c$  ✓✓
- (ii)  $\ln(7) - \ln(1) = \ln 7$  ✓
- $y(x) = -\frac{2}{5}e^{-5x} + c$  ✓
- (d) Since  $y(0) = -\frac{2}{5} + c = -3 \rightarrow c = -\frac{13}{5}$
- $\therefore y = -\frac{2}{5}e^{-5x} - \frac{13}{5}$  ✓ [11]
4. (a)  $X \sim N(7, 1.5^2)$
- $\therefore 7 - 3(1.5) < X < 7 + 3(1.5)$  ✓
- $\therefore 2.5 < X < 11.5$  so 9 years ✓
- (b)  $P(X < 5.5) = 0.5 - 0.34 = 0.16$  ✓
- $\therefore 16\%$  ✓
- (c)  $P(X > h) = 0.975 \rightarrow h = 7 - 2(1.5) = 4$  years ✓✓ [6]
5. Width of CI =  $2E$
- Since  $E = \sqrt{\frac{\hat{p}(\hat{p}(1-\hat{p}))}{n}} \rightarrow E \propto \frac{1}{\sqrt{n}}$  ✓
- $n \propto \frac{1}{E^2}$  ✓
- $\therefore$  New  $n = \frac{1}{4}n = \frac{n}{4}$  ✓ [3]

$$(2x)(\ln x^2) + (x^2)\left(\frac{2x}{x^2}\right) = 4x \ln x + 2x$$

6. (a) (i) ✓✓

$$\int (4x \ln x + 2x) dx = x^2 \ln x^2 + c$$

(ii) ✓

$$\therefore \int (4x \ln x) dx + \int 2x dx = x^2 \ln x^2 + c$$

✓

$$\therefore 4 \int (x \ln x) dx + x^2 = x^2 \ln x^2 + c$$

✓

$$\begin{aligned} \int (x \ln x) dx &= \frac{1}{4}(x^2 \ln x^2) - \frac{x^2}{4} + c \\ &= \frac{x^2(2 \ln x - 1)}{4} + c \end{aligned}$$

✓

$$\begin{aligned} \text{(b) Area} &= \int_0^1 |x \ln x| dx \\ &= -\int_0^1 x \ln x dx \\ &= -\left[\frac{1}{4}(2 \ln 1 - 1)\right] - 0 = \frac{1}{4} \end{aligned}$$

✓✓ [10]

$$\frac{dy}{dx} = 3x^2 + 2ax + b$$

7. ✓

$$\frac{d^2y}{dx^2} = 6x + 2a$$

✓

$$\frac{d^2y}{dx^2} = 0 \text{ when } x = 2$$

Since HPOI ✓

$$\therefore 12 + 2a = 0 \rightarrow a = -6$$

✓

$$\frac{dy}{dx} = 0 \text{ when } x = 2$$

And

$$\therefore 12 - 24 + b = 0 \rightarrow b = 12$$

✓

$$(2, 8) \rightarrow 8 = 8 - 24 + 24 + c \rightarrow c = 0$$

✓ [6]


**Calculator-Assumed Solutions**

8. (a)  $\int_0^2 ax^3 dx = 1 \rightarrow \left[ \frac{ax^4}{4} \right]_0^2 = 1$  ✓
- $\therefore 4a = 1 \rightarrow a = \frac{1}{4}$  ✓
- (b)  $P(x < 1) = \int_0^1 \frac{x^3}{4} dx$  ✓
- $= \frac{1}{16}(1)^4 = \frac{1}{16}$  ✓
- (c)  $\int_q^2 \frac{x^3}{4} = \frac{1}{4}$  ✓
- $\therefore \frac{2^4}{16} - \frac{q^4}{16} = \frac{1}{4}$  ✓
- $\therefore q = 1.8612$  ✓ [7]
9. (a) (i)  $f(2) = e^2 = 7.389$  ✓
- (ii)  $f(2) = 1 + \frac{2}{1} + \frac{4}{2} + \frac{8}{6} + \frac{16}{24} = 7$  ✓
- (b)  $f(-1) = 1 + \frac{(-1)}{1} + \frac{1}{2} + \frac{(-1)}{6} + \frac{1}{24} = 0.375$  ✓✓
- (c)  $\frac{d}{dx}(e^x) = \frac{d}{dx} \left( 1 + \frac{x}{1} + \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24} + \dots \right)$  ✓
- $= 1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \dots = e^x$  ✓ [6]
10. (a)  $S = 120 + 20 = 140^\circ$  ✓
- (b)  $S = 120e^{-0.02(120)} + 20 = 30.89^\circ$  ✓✓
- (c)  $25 = 120e^{-0.02t} + 20$  ✓
- $\therefore t = 159 \text{ seconds} \approx 3 \text{ mins}$  ✓✓
- (d)  $\frac{d}{dt} (120e^{-0.02t} + 20) = -2.4e^{-0.02t}$  ✓
- When  $t = 30$ , rate =  $-1.3^\circ/\text{sec}$  ✓ [8]


11. (a)  $X \sim N(85, 20^2)$   
 $P(X < 110) = 0.8944$  ✓  
 $\therefore 89.44\%$  ✓
- (b)  $Y \sim \text{Bin}(15, 0.8944)$
- (i)  $P(Y = 10) = 0.01292$  ✓✓
- (ii)  $(0.8944)^3 \times \text{Bin}(6, 12, 12, 0.8944)$  ✓✓  
 $= 0.7154$  ✓
- (c)  $\$10 \quad \frac{0.0388}{0.1056} = 0.3677$  ✓  
 $\$20 \quad \frac{0.0267}{0.1056} = 0.2532$  ✓  
 $\$30 \quad \frac{0.0401}{0.1056} = 0.3792$  ✓
- (d)  $E(Y) = 10 \times 0.3677 + 20 \times 0.2532 + 30 \times 0.3792$  ✓  
 $= \$20.12$  ✓
- (e)  $\text{VAR}(Y) = 10^2(0.3677) + 20^2(0.2532) + 30^2(0.3792) - (20.12)^2$  ✓✓  
 $= 74.516$  ✓  
 St. Dev.  $= 8.63$  ✓
- (f) New Mean  $= 1.2 \times 20.12 - 1 = \$23.14$  ✓  
 New St. Dev.  $= 1.2 \times 8.63 = \$10.36$  ✓ [17]
12. (a) (i) 17.32 m/min ✓  
 (ii) 20 m/min ✓
- (b)  $x(t) = \int 20\cos\left(\frac{t}{3} + \frac{\pi}{6}\right) dt = 60\sin\left(\frac{t}{3} + \frac{\pi}{6}\right) + c$  ✓  
 $x(0) = 0 \rightarrow c = -30$  ✓
- $x(t) = 60\sin\left(\frac{t}{3} + \frac{\pi}{6}\right) - 30$  ✓  
 $\therefore$  ✓
- (c) (i)  $x(4) = 27.56 \text{ m}$  ✓  
 (ii)  $v(4) = -5.6 \therefore$  returning to start. ✓
- (d)  $a(t) = -\frac{20}{3}\sin\left(\frac{t}{3} + \frac{\pi}{6}\right)$  ✓  
 $-\frac{20}{3}\sin\left(\frac{t}{3} + \frac{\pi}{6}\right) = -5 \rightarrow t = 5.30$  ✓  
 $\therefore$  ✓
- Distance  $= \int_0^{5.30} \left| 20\cos\left(\frac{t}{3} + \frac{\pi}{6}\right) \right| dt = 44.9 \text{ m}$  ✓  
 $\therefore$  ✓ [10]



13. (a)



OR



$$M = z \sqrt{\frac{\hat{p}(1-\hat{p})}{n}} \text{ where } \hat{p} = 0.5$$

(b)

$$0.06 = 1.96 \sqrt{\frac{(0.5)(0.5)}{n}}$$

 $\therefore$ 

$$n = 266.8$$

 $\therefore$ 

Hence 267 trainers should be tested

$$0.08 = 1.96 \sqrt{\frac{(0.8)(0.2)}{n}}$$

(c)

$$n = 96.04$$

 $\therefore$ 

$$\therefore 96 \times 0.8 = 76.8$$

Hence 76 or 77 trainers would be qualified.

[10]

14. (a)

$$\hat{p} \sim N\left(0.04, \left(\sqrt{\frac{0.04(0.96)}{300}}\right)^2\right)$$

$$\text{i.e. } \hat{p} \sim N(0.04, 0.0113^2)$$

$$(b) \quad P(X < 0.045) = 0.6709$$

$$(c) \quad (0.04 \pm 1.96 \times 0.0113)$$

$$\therefore (0.0179, 0.0621)$$

$$(d) \quad \frac{24}{300} = 0.08$$

Since 0.08 is not in the 95% CI  
it would be unlikely to happen.

(e) (i)  $n$  should be larger

Since the formula for standard error has  $n$  in the denominator,  
increasing  $n$  will decrease the standard error.

(ii) From the standard normal distn,  $P(-k < p < k) = 0.98$ 

$$k = 2.326$$

✓ [15]



15. (a)  $\ln 0$  doesn't exist. ✓
- (b)  $f'(x) = 2x - \frac{1}{x}$  ✓
- $f'(x) = 0$  when  $2x = \frac{1}{x} \rightarrow 2x^2 = 1$  ✓
- $x = \pm \frac{1}{\sqrt{2}}$  ✓
- $\therefore$  ✓
- $f''(x) = 2 + \frac{1}{x^2}$  ✓
- $f''\left(-\frac{1}{\sqrt{2}}\right)$  doesn't exist
- and  $f''\left(\frac{1}{\sqrt{2}}\right) > 0$ ,  $f\left(\frac{1}{\sqrt{2}}\right)$  is the minimum
- Since
- $\left(\frac{1}{\sqrt{2}}, \frac{1}{2} + \frac{\ln 2}{2}\right)$  is the minimum point. ✓ [6]
16. (a) (i)  $x = 2$  ✓
- (ii)  $a = 2$  ✓
- (iii) We require another set of co-ordinates, as  $\log_p 1 = 0$  for all  $p$ . ✓
- (b)  $f(x) + 1 = \log_p(x - 2) + 1$  since  $f(x)$  is translated 1 up. ✓
- $\log_p(x - 2) + 1 = 0 \rightarrow x - 2 = \frac{1}{p}$  ✓
- $\therefore$  ✓
- $x = \frac{1}{p} + 2$  ✓
- $\therefore$  ✓
- (c)  $f(27) = \log_p(25) = 2 \rightarrow p^2 = 25$  ✓
- $\therefore p = 5$  ✓ [8]
17. (a)  $\int_1^5 f'(x) dx = [f(x)]_1^5$  ✓
- $= f(5) - f(1) = -1 - (-2) = 1$  ✓
- (b) Area =  $\int_1^5 |f'(x)| dx$  ✓
- $= \int_1^3 f'(x) dx - \int_3^5 f'(x) dx$  ✓

$$= \left[ f(x) \right]_1^3 - \left[ f(x) \right]_5^3 = 2 - (-2) + 2 - (-1) = 7 \quad \checkmark\checkmark \quad [6]$$

18. (a)  $\log w = 0.5 + 0.4(4) = 2.1$  ✓  
 $\therefore w = 125.9 \text{ kg}$  ✓  
 $\log 180 = 0.5 + 0.4h \rightarrow h = 4.39 \text{ m}$  ✓✓  
(b)  $\log 200 = 0.5 + 0.4h \rightarrow h = 4.50 \text{ m}$  ✓  
(c)  $\log 100 = 0.5 + 0.4h \rightarrow h = 3.75 \text{ m}$  ✓  
 $\therefore \frac{4.50}{3.75} = 1.2 \rightarrow 20\% \text{ taller}$  ✓ [7]