


Ref: 14-085



MATHEMATICS

UNITS 3C AND 3D

707 **FORMULA SHEET**

© School Curriculum and Standards Authority, 2014 Copyright

and Standards Authority is acknowledged as the copyright owner. for non-commercial purposes by educational institutions, provided that it is not changed in any way and that the School Curriculum This document—apart from any third party copynight material contained in it—may be freely copied, or communicated on an intranet,

Copying or communication for any other purpose can be done only within the terms of the Copyright Act or by permission of the

Copying or communication of any third party copyright material contained in this document can be done only within the ferms of the

This document is valid for teaching and examining until 31 December 2014.

Mathematics 3C and 3D Formula Sheet updated January 2014

TORMULA SHEET

UNITS 3C AND 3D SOITAMEHTAM

Chance and data

For any event A and its complement A, and event BProbability:

 $(g|V)_d(g)_d = (V|g)_d(V)_d = (g \cup V)_d$

In a binomial distribution:

 $\frac{u \wedge}{Q} z + \underline{x} \geq n \geq \frac{u \wedge}{Q} z - \underline{x}$

 σ is the population standard deviation, where μ is the population mean,

'ueəm əldmes ətt si \underline{x}

 $(A \cap A) - (A) + (A) + (A) = (A \cup A) - (A \cap A)$ $I = (\underline{V})d + (V)d$

Mean: $\sigma = \eta du = 0$ and standard deviation: $\sigma = \eta du = 0$

A confidence interval for the mean of a population is:

bns exis eldmes ent si n

to the confidence level. z is the cut-off value on the standard normal distribution corresponding

included in the body of the particular question. Note: Any additional formulas identified by the examination panel as necessary will be

MATHEMATICS UNITS 3C AND 3D

2 **FORMULA SHEET**

Number and algebra

For a, b > 0 and m, n real, Index laws:

$$a^m b^m = (a b)^m$$

$$a^m a^n = a^{m+n}$$

$$a^m a^n = a^{m+n} (a^m)^n = a^{mn}$$

$$a^{-m} = \frac{1}{a^m}$$

$$a^{-m} = \frac{1}{a^m} \qquad \qquad \frac{a^m}{a^n} = a^{m-n}$$

$$a^0 = 1$$

For a > 0 and m an integer and n a positive integer, $a^{\frac{m}{n}} = \sqrt[n]{a^m} = \left(\sqrt[n]{a}\right)^m$

If f(x) = y then $f'(x) = \frac{dy}{dx}$ Differentiation:

If
$$f(x) = x^n$$
 then $f'(x) = nx^{n-1}$

If
$$f(x) = e^x$$
 then $f'(x) = e^x$

Product rule: If y = f(x) g(x)

If
$$y = uv$$

then
$$y' = f'(x) g(x) + f(x) g'(x)$$

then
$$y' = f'(x) g(x) + f(x) g'(x)$$
 or then $\frac{dy}{dx} = \frac{du}{dx}v + u\frac{dv}{dx}$

Quotient rule:

If
$$y = \frac{f(x)}{g(x)}$$

then
$$y' = \frac{f'(x) g(x) - f(x) g'(x)}{(g(x))^2}$$

$$\begin{aligned} &\text{If } y = \frac{f(x)}{g(x)} \\ &\text{then } y' = \frac{f'(x) \ g(x) - f(x) \ g'(x)}{(g(x))^2} \end{aligned} \qquad \text{or} \\ &\text{then } \frac{dy}{dx} = \frac{\frac{du}{dx} \ v - u \frac{dv}{dx}}{v^2} \end{aligned}$$

Chain rule:

If
$$y = f(g(x))$$

If
$$y = f(u)$$
 and $u = g(u)$

then
$$y' = f'(g(x)) g'(x)$$

or
$$fy = f(u) \text{ and } u = g(x)$$
 then
$$\frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx}$$

Integration:

Powers:

$$\int x^n dx = \frac{x^{n+1}}{n+1} + c, \ n \neq -1$$

Exponentials:

$$\int e^x dx = e^x + c$$

Fundamental Theorem of Calculus:

$$\frac{d}{dx} \left(\int_a^x f(t) dt \right) = f(x) \qquad \text{and} \qquad \int_a^b f'(x) dx = f(b) - f(a)$$

Incremental formula: $\delta y \simeq \frac{dy}{dx} \delta x$

$$\delta y \simeq \frac{dy}{dx} \, \delta x$$

Exponential growth and decay:

If
$$\frac{dy}{dt} = ky$$
, then $y = Ae^{kt}$

See next page

FORMULA SHEET

MATHEMATICS **UNITS 3C AND 3D**

Space and measurement

 $C = 2\pi r = \pi D$, where *C* is the circumference, Circle:

 \emph{r} is the radius and \emph{D} is the diameter

3

 $A = \pi r^2$, where A is the area

 $A = \frac{1}{2}bh$, where b is the base and h is the perpendicular height Triangle:

Parallelogram: A = bh

 $A = \frac{1}{2}(a+b)h$, where a and b are the lengths of the parallel sides Trapezium:

V = Ah, where V is the volume and A is the area of the base Prism:

 $V = \frac{1}{2} Ah$ Pyramid:

Cylinder: $S = 2\pi rh + 2\pi r^2$, where S is the total surface area

 $S = \pi r s + \pi r^2$, where s is the slant height Cone:

 $V = \frac{1}{3}\pi r^2 h$

Sphere: $S = 4\pi r^2$

 $V = \frac{4}{2} \pi r^3$

Volume of solids of revolution:

 $V = \int \pi y^2 dx$ rotated about the x-axis

 $V = \int \pi x^2 dy$ rotated about the *y*-axis