

MATHEMATICS

UNIT 3

METHODS

Semester One

2018

SOLUTIONS

Test papers may only be reproduced within the purchasing school according to the Advertised Conditions of Sale.
Copyleft for test papers and marking guides remains with West Australian Test Papers.

19. (a) $A = A_0 e^{-0.05t}$
- $\therefore A = 0.6e^{-0.05(10)} = 0.99 \text{ Ha}$
- (b) $5 = 0.6e^{-0.05t}$
- $\therefore t = 42.4 \text{ hours}$
- [6]
- Since $f''\left(\frac{\pi}{4}\right) < 0$ then maximum
POI occurs when $f''(x) = 0$
- $\therefore x = \frac{3\pi}{4} \text{ or } \frac{7\pi}{4}$
- $\therefore -\cos x + \sin x = 0 \rightarrow \tan x = -1$
- $\therefore \left(\frac{3\pi}{4}, 0\right) \text{ and } \left(\frac{7\pi}{4}, 0\right)$
- [10]
21. (a) 10 L/sec
- $10 - \frac{20}{t} = 0 \rightarrow t = 200 \text{ secs}$
- (b) $F = \int_0^{200} \left(10 - \frac{t}{20}\right) dt$
- $\therefore F = 1000 \text{ L}$
- [4]
22. (a) $r' = e^{-t}$
- $\therefore r(4) = e^{-4} = 0.018$
- and $r'(5) = e^{-5} = 0.0067$
- $\therefore r''(t) = -e^{-t}$
- Since $-e^{-t} \neq 0$, then the fire is spreading at its greatest rate at the start of the fire.
- Or the graph has greatest slope at $t = 0$.
- greatest rate of spread at its start of the fire.
- [9]

- When $t = 4$ $\frac{dA}{dt} = 2\pi(-e^{-4} + 4)(0.018) = 0.458 \text{ cm}^2/\text{sec}$
- $\therefore \frac{dA}{dt} = \frac{dA}{dr} \times \frac{dr}{dt} = 2\pi r \times e^{-t}$
- $\therefore A = \pi r^2 \rightarrow \frac{dA}{dt} = 2\pi r$
- Or the graph has greatest slope at $t = 0$.
- greatest rate of spread at its start of the fire.
- Since $-e^{-t} \neq 0$, then the fire is spreading at its greatest rate at the start of the fire.
- [9]

Calculator-free Solutions

1.. $f'(x) = 0 \text{ when } \frac{1-x}{e^x} = 0$ ✓
 $\therefore x = 1$ ✓

$f''(x) = \frac{(e^x)(-1) - e^x(1-x)}{e^{2x}}$ ✓
Since $f''(1) < 0$, then maximum ✓

[4]

2. $A = \int_0^k e^{3-x} dx$ ✓
 $\therefore \left[-e^{3-x} \right]_0^k = (-e^{3-k}) - (-e^{3-0})$ ✓

If $A = e^3(1 - e^{-k}) = e^3$ then $e^{-k} = 0$, which is impossible. ✓

If $A = e^3(1 - e^{-k}) = e^3 - 1$ ✓
 $\therefore k = 3$ ✓

[6]

3. (a) Stationary point occurs when $3x^2 - kx = 0$ ✓
 $\therefore M'(6) = 108 - 6k = 0$ ✓
 $\therefore k = 18$ ✓

(b) $M(x) = x^3 - 9x^2 + c$ ✓
Since $M(6) = 1 \rightarrow 216 - 324 + c = 1 \rightarrow c = 109$ ✓

$\therefore M(x) = x^3 - 9x^2 + 109$ ✓
 \therefore Vertical intercept = 109 ✓

[6]

4. (a) $e^1 = 1 + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \frac{1}{4!}$ ✓
 $= 1 + 1 + \frac{1}{2} + \frac{1}{6} + \frac{1}{24} = 2\frac{17}{24}$ ✓

(b) $\frac{d}{dx} \left(1 + \frac{x}{1} + \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24} + \dots \right)$ ✓
 $= 1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24} + \dots$ ✓

$= 1 + \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \dots = e^x$ ✓

[4]

16. (a) $2\cos x \sin x + \cos x = 0$

$\cos x = 0 \text{ or } \sin x = -\frac{1}{2}$
 $\therefore x = \frac{\pi}{2}, \frac{3\pi}{2}, \frac{7\pi}{6}, \frac{11\pi}{6}$

(b) $\left(\frac{\pi}{2}, 0 \right), \left(\frac{3\pi}{2}, 0 \right), \left(\frac{7\pi}{6}, \frac{\sqrt{3}}{2} \right), \left(\frac{11\pi}{6}, -\frac{\sqrt{3}}{2} \right)$

$\int_{\frac{\pi}{2}}^{\frac{7\pi}{6}} (-\cos x - \sin 2x) dx$

(c) $A = \frac{\pi}{2} = 2.25 \text{ units}^2$

17. (a) $\frac{x+1}{x-1} = 0 \rightarrow x = -1$

(b) (i) $\frac{dy}{dx} = \frac{(x-1)-(x+1)}{(x-1)^2} = \frac{-2}{(x-1)^2}$

(ii) True since $\frac{dy}{dx} \neq 0$ for any value of x .

(c) (i) $\frac{d^2y}{dx^2} = \frac{4}{(x-1)^3}$

(ii) True since $\frac{d^2y}{dx^2} \neq 0$ for any value of x .

18. (a) $v(0) = 3$

(b) Stationary when $2t^2 - 5t + 3 = 0$

$\therefore t = 1 \text{ and } 1.5$

(c) (i) $a = 4t - 5$

(ii) v is a minimum when $a = 0 \rightarrow a = 1.25$

(d) $x = \frac{2}{3}t^3 - \frac{5}{2}t^2 + 3t + c$

Since $x(0) = 0 \rightarrow c = 0$

$x(t) = \frac{2}{3}t^3 - \frac{5}{2}t^2 + 3t$

$\therefore x(3) = 4.5$

Distance = $\int_0^3 |v(t)| dt$

= 4.58

(f) Distance travelled \neq Displacement

[13]

[5]

$$\begin{aligned}
 & \text{(a) } y = (\sin x)^{-1} \\
 & \frac{dy}{dx} = -1(\sin x)^{-2}(\cos x) \\
 & \frac{d^2y}{dx^2} = \frac{\sin x}{\cos x} \\
 & \int \frac{\sin x}{\cos x} dx = \int \frac{1 - \cos^2 x}{\cos x} dx \\
 & = \int \frac{\sin^2 x}{\cos x} dx \\
 & = -5 \int \frac{\sin^2 x}{\cos x} dx \\
 & = -5 \int \frac{1 - \cos^2 x}{\cos x} dx \\
 & = -5 \int \frac{\sin^2 x}{\cos x} dx \\
 & = -\frac{\sin x}{\cos x} + C
 \end{aligned}$$

[6]

$$\begin{aligned}
 & \text{(a) } f(0) = -\frac{1}{3} \\
 & \text{Since PDF } \int f(x) = 0.8 + a = 1 \rightarrow a = 0.2 \\
 & \therefore f(x) = 0.8 + 0.2 = 1 \\
 & \therefore \text{PDF} \quad \text{and } 0 \leq f(x) \leq 1 \text{ for all } x \\
 & \therefore \text{PDF} \quad \text{and } 0 \leq f(x) \leq 0 \text{ for all } x \\
 & \text{Since } f(0) = -\frac{1}{3} \\
 & \text{f(x) is not greater than or equal to 0 for all } x \\
 & \therefore \text{Not PDF}
 \end{aligned}$$

[7]

$$\begin{aligned}
 & \text{(a) } E(Y) = 2E(X) + 3 = 8 \\
 & \therefore m = 2.5 \\
 & \therefore V = 5 \\
 & \text{VAR}(Y) = 4\text{VAR}(X) = 20
 \end{aligned}$$

[8]

$$\begin{aligned}
 & \text{(a) } E(X) = 0 + 0.1 + 0.2 + 0.3 + 0.8 = 1.4 \\
 & \text{(b) } P(X < 2 | X \leq 2) = \frac{P(X < 2)}{P(X \leq 2)} = \frac{0.6}{0.7} = \frac{6}{7} \\
 & \text{(c) } \text{Variance} = (0 + 1 \times 0.1 + 4 \times 0.1 + 9 \times 0.1 + 16 \times 0.2) - 1.4^2 = 0.1 + 0.4 + 0.9 + 3.2 - 1.96 = 2.64
 \end{aligned}$$

10. (a) $y = (ex)(e^x)$
 $\frac{dy}{dx} = e(e^x) + (ex)(e^x)$
 $\therefore = e^{x+1}(x+1)$

(b) $y = \frac{\pi \sin x}{\cos x}$
 $\frac{dy}{dx} = \frac{(\cos x)(\pi \cos x) - (\pi \sin x)(-\sin x)}{\cos^2 x}$
 $= \frac{\pi(\cos^2 x + \sin^2 x)}{\cos^2 x}$
 $= \frac{\pi \times 1}{\cos^2 x} = \frac{\pi}{\cos^2 x}$

✓

✓

✓

✓

[5]

Calculator-assumed Solutions

11. (a) $K = \frac{1}{2} r^2 \sin \theta \rightarrow \frac{dK}{d\theta} = 8 \cos \theta$ when $r = 4$
 $\delta K = \frac{dK}{d\theta} \times \delta\theta \rightarrow \delta K = 8 \cos \theta \times (0.05\pi)$
 $\therefore \delta K = 8 \cos \frac{\pi}{4} \times 0.05\pi$ when $\theta = \frac{\pi}{4}$
 $\therefore \delta K = 0.889$

(b) $\Delta K = \frac{1}{2} 16 \left(\sin 0.3\pi - \sin \frac{\pi}{4} \right)$
 $\therefore \Delta K = 0.8153$
 $\text{Error} = \frac{0.8886 - 0.8153}{0.8886} \times 100 = 8.3\%$

✓

✓

✓

✓

✓

✓

[5]

12. $(-3, 0) \rightarrow -27a - 9b - 3c - 9 = 0$
 $\frac{dy}{dx} = 3ax^2 - 2bx + c$
Stationary point at $x = -3 \rightarrow 0 = 27a + 6b + c$
 $\frac{d^2y}{dx^2} = 6ax - 2b$
POI at $-\frac{5}{3} \rightarrow -10a - 2b = 0$
 \therefore Using calc $a = 1, b = -5$ and $c = 3$

✓

✓

✓

✓

✓

✓

[7]

13. (a) $f(x) = -4 \cos \frac{x}{2} + c$
 $f\left(\frac{\pi}{2}\right) = -4 \cos \frac{\pi}{4} + c = 4 - 2\sqrt{2} \rightarrow c = 4$
 $\therefore f(x) = -4 \cos \frac{x}{2} + 4$
 $-4 \cos \frac{x}{2} + 4 = 4 - 2\sqrt{2}$

(b) $\cos \frac{x}{2} = \frac{\sqrt{2}}{2} \rightarrow \frac{x}{2} = \frac{\pi}{4}, \frac{7\pi}{4}, \dots$
 \therefore Next time is $x = \frac{7\pi}{2}$

✓

✓

✓

[6]

14. (a) (i) $0 + 1 + (-5) + 4 = 0$
(ii) $-3 + 1 + (-5) + 4 = -3$
(iii) $1 + 5 + 4 = 10$

✓✓

✓✓

✓✓

(b) (i) $-5 + 4 + \int_{2}^{4} 7 dx = -1 + (28 - 14) = 13$

✓✓

(ii) $2 \int_{3}^{4} f(x) dx = 2(4) = 8$

[10]

15. (a) (i) $X \sim B(30, 0.75)$ where X = the number graduated
 $\therefore P(X = 25) = 0.1047$

✓

✓✓

(ii) $P(X \geq 29 | X \geq 25) = \frac{0.00196}{0.2026}$
 $= 0.00967$

✓✓

(b) $Y \sim B(n, 0.75)$ where Y is the number who graduated out of n
 $\therefore P(Y \geq 10) \geq 0.99 \rightarrow n = 19$ using trial and error ✓✓✓

n	Y
18	0.981
19	0.991
20	0.996

$$\frac{\binom{5}{3} \binom{6}{0} + \binom{5}{2} \binom{6}{1} + \binom{5}{1} \binom{6}{2}}{\binom{11}{3}}$$

✓✓

(c) (i) $\frac{10 + 60 + 75}{165} = 0.87879$

✓

$P(S = n) = \begin{cases} \frac{5}{11}, & n = 1 \\ \frac{6}{11}, & n = 0 \end{cases}$

✓✓✓

[15]