

SOLUTIONS


2016

UNIT 3

MATHEMATICS METHODS

REVISION 1

SEMESTER ONE



SECTION ONE

1. (7 marks)

(a) $y = x^2(2x - 1)$

$$\frac{dy}{dx} = 2x(2x - 1) + 2(x^2)$$
✓✓ -1/error

$$\frac{dy}{dx} = 6x^2 - 2x$$

(2)

(b) $y = \frac{\sin(2x)}{2x}$

$$\frac{dy}{dx} = \frac{2(\cos(2x)) \times 2x - 2(\sin(2x))}{4x^2}$$

$$\frac{dy}{dx} = \frac{4x(\cos(2x)) - 2(\sin(2x))}{4x^2}$$


(3)

(c) $y = (x + e^x)^4$

$$\frac{dy}{dx} = 4(x + e^x)^3 \times (1 + e^x)$$
✓✓

(2)

- (1) In this case the velocity goes from decreasing to increasing so the velocity graph has a minimum turning point.
- (d) At $a(t) = 0$, the velocity graph has a turning point. ✓
- (e) At $a(t) = 0$ on the acceleration graph, there is a point of inflection on the displacement graph. ✓
- (f) The displacement graph is a cubic polynomial. ✓
- (g) At $a(t) = 0$ on the acceleration graph, there is a point of inflection on the displacement graph. ✓



- (e) At $v(t)=0$, then the particle changes direction on the displacement graph. The velocity goes from positive to negative or from negative to positive. ✓ (2)
 On the displacement graph, the first time $v(t)=0$ the displacement had been increasing and the particle turned around and began decreasing.
 The second time $v(t)=0$, the displacement had been decreasing and the particle turned around and began increasing.

3. (6 marks)

(a) (i) $\int \sqrt{2x+1} dx = \frac{2\sqrt{(2x+1)^3}}{3 \times 2} + c = \frac{\sqrt{(2x+1)^3}}{3} + c$ ✓ ✓ (2)
 (ii) $\int 1+x - e^{-x} dx = x + \frac{x^2}{2} + e^{-x} + c$ ✓ ✓ (2)

(b) $\frac{dy}{dx} = 2x + 3x^2 - x^{\frac{1}{2}}$

$$y = x^2 + x^3 - \frac{2x^{\frac{3}{2}}}{3} + c$$

(1,4) belongs to the function

$$4 = 1 + 1 - \frac{2}{3} + c$$

$$c = 2 \frac{2}{3}$$

$$y = x^2 + x^3 - \frac{2x^{\frac{3}{2}}}{3} + \frac{8}{3}$$




(2)

4. (7 marks)

(a) $\int_2^4 (x^2 - 2x + 3) dx = \left[\frac{x^3}{3} - x^2 + 3x \right]_2^4$
 $= \left(\frac{64}{3} - 16 + 12 \right) - \left(\frac{8}{3} - 4 + 6 \right)$
 $= \frac{56}{3} - 4 - 2$
 $= 12 \frac{2}{3}$

20. (5 marks)
 Four Apple MacBooks
 $p = 0.7$
 Three ASUS
 $p = 0.8$ ✓
 $P(x=3) \cap P(y=3)$
 $= 0.4116 \times 0.512$
 $= 0.2107392$
 The probability that 3 Apple MacBooks and 3 ASUS are being used is 0.21 (5)

END OF SECTION TWO



(2) $0.15 + 0.35 = 0.5 \quad \checkmark$

(a) (i) $0.15 \quad 0.25 \quad 0.35 \quad 0.25 \quad 0.25$

(2) $0.15 \quad 0.25 \quad 0.35 \quad 0.25 \quad 0.25$

(b) $\int (\sin(x) - \cos(x)) dx = [-\cos(x) - \sin(x)]_0^{\frac{\pi}{2}} = -(-1 + 0 - (0 + 1)) = -(-1 + 0 - (0 + 1)) = 2$

(a) $P(X = x)$

x	HH	HT or TH	TT
1	1/4	1/2	1/4
2	1/4	1/2	1/4
(1)	(2)	(2)	(2)
(ii) 0.0985 ✓	(iii) 0.8363 ✓	(iv) 0.9937 ✓	(v) (i) 0.9363 ✓



6. (8 marks)

(a) If $F(x) = x^3 - x^2$

(i) $F'(x) = 3x^2 - 2x$ ✓

(1)

(ii) $\int F'(x) dx = [x^3 - x^2]_0^p = p^3 - p^2$ ✓✓

(2)

(b) $F(x) = \int t^3 dt$

$F'(x) = x^3$ ✓✓

(2)

(c) $\frac{d}{dx} \left(\int^{3x} \cos(2y) dy \right) = 3\cos(6x)$
✓✓✓

(3)

7. (8 marks)

Given $f(x) = e^x$, $g(x) = \cos(x)$ and $h(x) = -x$

(a) (i) $y = h(g(x)) = h(\cos(x)) = -\cos(x)$ ✓

(1)

(ii) show $\frac{d}{dx}(h(g(x))) = g\left(\frac{\pi}{2} - x\right)$

$\frac{d}{dx}(h(g(x))) = -(-\sin(x)) = \sin(x)$ ✓

$g\left(\frac{\pi}{2} - x\right) = \cos\left(\frac{\pi}{2} - x\right) = \sin(x)$ ✓

(2)

(b) (i) $y = f(h(x)) = f(-x) = e^{-x}$ ✓

(1)

(ii) $\frac{d}{dx}(f(h(x))) = -e^{-x}$ ✓✓

(2)

(c) $g(f(x)) = g(e^x) = \cos(e^x)$ ✓

(2)

$g(f(0)) = \cos(e^0) = \cos(1)$ ✓

15. (13 marks)

(a) (i) $\int^p \sin(x) dx = 2$ ✓

(1)

(ii) $\int^{\pi/2} \sin(x) dx = 1$ An estimate can be made because the graph

is symmetrical. ✓

(1)

(iii) Area = 8 units² ✓

(1)

(iv) $\int^{4\pi} \sin(x) dx = 0$ ✓

(1)

(b) (i) Estimate from below

Area = $1 \times 0.5 + 1 \times 0.33 + 1 \times 0.25$ ✓✓

= 1.08

Estimate from above

Area = $1 \times 1 + 1 \times 0.5 + 1 \times 0.33$ ✓✓

= 1.83

(4)

(ii) $\int^4 \frac{1}{x} dx = \int^4 x^{-1} dx = \left[\frac{x^0}{0} \right]_1^4$ ✓✓

(3)

(iii) $\int^4 \frac{1}{x} dx = 1.386$ (3dp) ✓✓

(2)

16. (8 marks)

(a) 2000 $t = 0$ $P = 400$

2008 $t = 8$ $P = 550$

$550 = 400(r)^8$

$r = 1.040609622$

The annual rate of increase of the population of numbats was 4.06%. (3)

(b) 2016 $t = 16$ $P = ?$

$P = 400(1.040609622)^{16}$

$P = 756.25$

The expected population in 2016 is 756 numbats. (2)

(c) 2016 $t = 0$ $P = 756$

2020 $t = 4$ $P = 780$

$780 = 756(r)^4$

$r = 1.0078$

The rate of increase has dropped from 4.05% to 0.78%.

It is possible that predators had found a way in. (3)

END OF SECTION ONE

(a)

$$\begin{aligned}
 & \text{SD}(X) \approx 1.7 \\
 & \text{SD}(X) = 1.707825 \\
 & \text{Var}(X) = 15 \cdot \frac{1}{6} - 3.5^2 = 2.916 \\
 & E(X^2) = 1^2 \times \frac{1}{6} + 2^2 \times \frac{1}{6} + 3^2 \times \frac{1}{6} + 4^2 \times \frac{1}{6} + 5^2 \times \frac{1}{6} + 6^2 \times \frac{1}{6} = 15 \\
 & \text{Var}(X) = E(X^2) - (E(X))^2 \\
 & (iii) E(X) = 1 \times \frac{6}{6} + 2 \times \frac{6}{6} + 3 \times \frac{6}{6} + 4 \times \frac{6}{6} + 5 \times \frac{6}{6} + 6 \times \frac{6}{6} = 3.5 \\
 & (ii) P(Y \leq 4) = \frac{6}{4} = 0.75 \\
 & (i)
 \end{aligned}$$

(b)

$P(Y=y)$	6	6	6	6	6	6
y	0	1	2	3	4	5

9. (10 marks)

(a)

The decrease in volume when the side has melted to 9 mm is 0.3 cm^3

$$\begin{aligned}
 \Delta V & \approx -0.3 \text{ cm}^3 \\
 \Delta V & \approx -0.3 \times 1^2 \times (-0.1) = -0.3 \\
 \Delta V & \approx -0.3 \times 1 \text{ cm}, x = 1 \text{ cm} \\
 \Delta V & \approx -3x^2 \times \Delta x \\
 \Delta V & \approx \frac{dx}{dx} \times \Delta x \\
 \Delta V & = 3x^2 \\
 V & = x^3
 \end{aligned}$$

8. (4 marks)

SECTION TWO

14. (9 marks)

$$\begin{aligned}
 & \therefore y = e^x \sin(x) + 1 \\
 & \text{At } (0,1) 1 = 0 + c \\
 & \int e^x (\sin(x) + \cos(x)) dx = e^x \sin(x) + c \\
 & \text{Hence} \\
 & f(x) = e^x \sin(x) \leftarrow f'(x) = e^x \sin(x) + e^x \cos(x) \\
 & (i) \quad \int_{\frac{\pi}{8}}^{\frac{\pi}{4}} \sin(2x) \left| \frac{1+2x}{\sqrt{2}} dx = 0.0981 \right. \wedge
 \end{aligned}$$

13. (5 marks)

$$\begin{aligned}
 & \text{At } t = 3 \text{ s, } x = 6.5. \quad \wedge \\
 & \text{At } t = 2 \text{ s, } x = 8 \\
 & x = 8 \text{ m} \quad \wedge \\
 & \text{Changes direction when } v = 0 \text{ i.e. at } t = 2 \text{ s} \quad \wedge \\
 & (d)
 \end{aligned}$$

$$\begin{aligned}
 & a = -3 \text{ m/s}^2 \quad \wedge \\
 & (b) \quad a = -3 \text{ m/s}^2 \quad \wedge \\
 & (c) \quad 2 = -\frac{3t^2}{2} + 6t + 2 \quad \Leftarrow t = 4 \text{ s} \\
 & (d) \quad x = -\frac{3t^2}{2} + 6t + 2 \quad \Leftarrow
 \end{aligned}$$

$$\begin{aligned}
 & x = -\frac{3t^2}{2} + 6t + 2 \\
 & \text{At } t = 0, x = 2 \Leftarrow c = 2 \\
 & x = -\frac{3t^2}{2} + 6t + c \\
 & x = \int (-3t + 6) dt
 \end{aligned}$$

(b)

Not a probability density function as you can't have a negative probability

(c)

Is a probability density function as the total is 1.

(d)

Probability.

(e)

Not a probability density function as you can't have a negative

(f)

Not a probability density function as the total is only 0.9.

(g)

Greater than 1.

(h)

Not a probability density function as you can't have a probability

(i)

Probability.

(j)

Not a probability density function as you can't have a negative

(k)

Probability.

(l)

Not a probability density function as you can't have a negative

(m)

Probability.

(n)

Probability.

(o)

Probability.

(p)

Probability.

(q)

Probability.

(r)

Probability.

(s)

Probability.

(t)

Probability.

(u)

Probability.

(v)

Probability.

(w)

Probability.

(x)

Probability.

(y)

Probability.

(z)

Probability.

(aa)

Probability.

(bb)

Probability.

(cc)

Probability.

(dd)

Probability.

(ee)

Probability.

(ff)

Probability.

(gg)

Probability.

(hh)

Probability.

(ii)

Probability.

(jj)

Probability.

(kk)

Probability.

(ll)

Probability.

(mm)

Probability.

(nn)

Probability.

(oo)

Probability.

(pp)

Probability.

(qq)

Probability.

(rr)

Probability.

(ss)

Probability.

(tt)

Probability.

(uu)

Probability.

(vv)

Probability.

(ww)

Probability.

(xx)

Probability.

(yy)

Probability.

(zz)

Probability.

(aa)

Probability.

(bb)

Probability.

(cc)

Probability.

(dd)

Probability.

(ee)

Probability.

(ff)

Probability.

(gg)

Probability.

(hh)

Probability.

(ii)

Probability.

(jj)

Probability.

(kk)

Probability.

(ll)

Probability.

(mm)

Probability.

(nn)

Probability.

(oo)

Probability.

(pp)

Probability.

(qq)

Probability.

(rr)

Probability.

(ss)

Probability.

(tt)

Probability.

(uu)

Probability.

(vv)

Probability.

(ww)

Probability.

(xx)

Probability.

(yy)

Probability.

(zz)

Probability.

(aa)

Probability.

(bb)

Probability.

(cc)

Probability.

(dd)

Probability.

(ee)

Probability.

(ff)

Probability.

(gg)

Probability.

(hh)

Probability.

(ii)

Probability.

(jj)

Probability.

(kk)

Probability.

(ll)

Probability.

(mm)

Probability.

(nn)

Probability.

(oo)

Probability.

(pp)

Probability.

(qq)

Probability.

(rr)

Probability.

(ii)

Probability.

(jj)

Probability.

(kk)

Probability.

(ll)

Probability.

(mm)

Probability.

(nn)

Probability.

(oo)

Probability.

(pp)

Probability.

(qq)

Probability.

(rr)

Probability.

(ii)

Probability.

(jj)

Probability.

(kk)

Probability.

(ll)

Probability.

(mm)

Probability.

(nn)

Probability.

(oo)

Probability.

(pp)

Probability.

(qq)

Probability.

(rr)

Probability.

(ii)

Probability.

(jj)

Probability.

(kk)

Probability.

(ll)

Probability.

(mm)

Probability.

(nn)

Probability.

(oo)

Probability.

(pp)

Probability.

(qq)

Probability.

(rr)

Probability.

(ii)

Probability.

(jj)

Probability.

(kk)

Probability.

(ll)

Probability.

(mm)

Probability.

(nn)

Probability.

(oo)

Probability.

(pp)

Probability.

(qq)

Probability.

(rr)

Probability.

(ii)

Probability.

(jj)

10. (8 marks)

(a) $P = \pi r + 2r + 2l$ (1)

(b) $A = \frac{\pi r^2}{2} + 2rl$
But $P = 4$
 $4 = \pi r + 2r + 2l \Rightarrow l = \frac{4 - \pi r - 2r}{2}$
 $\therefore A = \frac{\pi r^2}{2} + 2r\left(\frac{4 - \pi r - 2r}{2}\right)$

$$\begin{aligned}A &= \frac{\pi r^2}{2} + 4r - \pi r^2 - 2r^2 \\A &= 4r - \frac{\pi r^2}{2} - 2r^2\end{aligned}$$

(c) $A = 4r - \frac{\pi r^2}{2} - 2r^2$

Maximum area occurs when $A'(r) = 0$ and $A''(r) < 0$

$A'(r) = 4 - \pi r - 4r$

$A''(r) = -\pi - 4$

If $A'(r) = 0$ then $0 = 4 - \pi r - 4r \Rightarrow r = \frac{4}{\pi + 4}$

$r = 0.5600991535$


$r \approx 0.56$

$A''\left(\frac{4}{\pi + 4}\right) = -\pi - 4 < 0$ so maximum

At $r = 0.56$, $A = 1.1202 \text{ m}^2$

The maximum area of the window is 1.12 m^2 .

(5)



(2)

11. (6 marks)

(a) $t = 2s$ ✓ (1)


(b) $a > 0 \quad \forall t \text{ s.t. } t \geq 0.$
 $x = (t - 2)^2 + 2$
 $v = \frac{dx}{dt} = 2(t - 2)$
 $a = \frac{d^2x}{dt^2} = 2$ which is always positive! (2)

(c) $v = 2(t - 2)$
At $t = 3$, $v = 2 \text{ m/s.}$ ✓ (1)


(d) Distance travelled for $1 \leq t \leq 4 = ?$
 $x(1) = 3, x(2) = 2, x(4) = 6$
Distance travelled = $1 + 4 = 5 \text{ m}$ (2)

12. (7 marks)

(a) (i) (3)



(ii) Point shows where the maximum gradient is. (1)



(b) If $f(x) = \cos(x)$, then $f'(x) = -\sin(x)$ ✓ (1)

(c) $\lim_{h \rightarrow 0} \left(\frac{e^h - 1}{h} \right) = 1 \quad \checkmark \checkmark$ (2)