

further.  
room. If you have any unauthorised material with you, hand it to the supervisor before reading any  
do not have any unauthorised notes or other items of a non-personal nature in the examination  
No other items may be taken into the examination room. It is your responsibility to ensure that you

### **Important note to candidates**

conditions set by the Curriculum Council for this course.  
and up to three calculators, CAs, graphic or scientific, which satisfy the  
Special items: drawing instruments, templates, notes on up to two unfolded sheets of A4 paper,

Standard items: pens, pencils, pencil sharpener, highlighter, eraser, ruler.

### **To be provided by the candidate**

Formula sheet.

Question/answer booklet for Section Two.  
To be provided by the supervisor

### **Material required/recommended for this section**

Working time for paper: 100 minutes  
Reading time before commencing work: 10 minutes

### **Time allowed for this section**

---

Your name \_\_\_\_\_  
(Calculator Assumed)  
Section Two

### **MATHEMATICS 3C**

Question/Answer Booklet  
Semester 1 Examination 2010

Insert School Name	Insert School Name	Logo
--------------------	--------------------	------

**Structure of this examination**

	Number of questions	Working time (minutes)	Marks available
Section One Calculator Free	7	50	40
<b>This Section (Section 2) Calculator Assumed</b>	<b>12</b>	<b>100</b>	<b>80</b>
Total marks			120

DO NOT WRITE IN THIS AREA

**Instructions to candidates**

1. The rules for the conduct of WACE external examinations are detailed in the booklet *WACE Examinations Handbook*. Sitting this examination implies that you agree to abide by these rules.
2. Answer the questions in the spaces provided.
3. Spare answer pages are provided at the end of this booklet. If you need to use them, indicate in the original answer space where the answer is continued i.e. give the page number.
4. Show all working clearly. Any question, or part question, worth more than 2 marks requires valid working or justification to receive full marks. If you repeat an answer to any question, ensure that you cancel the answer you do not wish to have marked.

(2 marks)

(d) Julie is the only centre in the team?

Julie is a centre player and is chosen in the team. If the other players are selected at random, what is the probability that:

(3 marks)

(c) the team includes at most 2 centres?

(2 marks)

(b) the team must consist of 2 guards, 1 centre and 2 forwards?

(1 mark)

(a) there are no restrictions?

In how many ways can this starting team be chosen if:

A basketball training squad consists of 4 guards, 2 centres and 3 forwards. A team of 5 is to be chosen to start the game.

(8 marks)

### Question 8

Suggested working time for this section is 100 minutes.

DO NOT WRITE IN THIS AREA

number of the question(s) that you are continuing to answer at the top of the page.

original answer space where the answer is continue, i.e. give the page number. Fill in the

continuing an answer: If you need to use the space to continue an answer, indicate in the

Planning: If you use the spare pages for planning, indicate this clearly at the top of the page.

Space pages are included at the end of this booklet. They can be used for planning your responses and/or as additional space if required to continue an answer.

This section has twelve (12) questions. Answer all questions. Write your answers in the space provided.

Section Two: Calculator-assumed  
(80 Marks)

**Question 9**

- (a) The composite function  $f(g(x)) = e^{2x-6}$ . Determine two different pairs of equations for functions  $f(x)$  and  $g(x)$ . (2 marks)

- (b) If  $f(x) = 3x^2 - 2$  and  $h(x) = \frac{3}{1-x}$  find  $h(f(x))$ . (1 mark)

- (c) A composite function is defined by the equation  $h(f(x)) = \sqrt{\xi-3} - 4$ . Determine the domain and range of this function for  $x$  real. (2 marks)

DO NOT WRITE IN THIS AREA

See Next Page

(5 marks)


**Question 19**

The cross section of land can be modeled by the equation

$$H = 0.00003d(d - 30)(d - 50)(100 - d)$$

where  $H$  and  $d$  are, respectively, the height (in metres) above a fixed horizontal level and the distance (in metres) from a fixed point.

The cross section has been shown in the diagram below.



DO NOT WRITE IN THIS AREA

A tunnel, 10m high, will be constructed through the two hills. Show how the cross sectional area of soil removed can be determined using integrals and mensuration (measurement) formula. There is no need to evaluate your answer.

$x$  intercepts at 30 m, 50 m and 100 m ✓

(5 marks)

Solve for  $0.00003d(d - 30)(d - 50)(100 - d) = 10$   
 $d = 2.64, 23.03, 55.33, 99.00$  ✓ ✓


Area =  $\int_0^{2.64} H + \int_{23.03}^{55.33} H + \int_{50}^{99.00} H - \int_{99.00}^{100} H$   
 $\checkmark \checkmark$

See Next Page

CALCULATOR ASSUMED  
MATHEMATICS 3C

Two competing cyclists are riding with constant speed. At 12 midday cyclist X is 40 metres north of a judge and is riding east at 9 m/s, while cyclist Y is 70 metres east of the judge and is riding north at 7 m/s.

(a) Show diagrammatically this situation (a scale diagram is not required) (1 mark)



(b) If the distance between the cyclist  $t$  seconds later is D metres, show that

$$D^2 = 6500 - 1820t + 130t^2 \quad (3 \text{ marks})$$

$$\begin{aligned} &= 6500 - 1260t + 81t^2 + 1600 - 560t + 49t^2 \\ &= 4900 - 1260t + (81t^2 + 49t^2) \\ &= (70 - 9t)^2 + (40 - 7t)^2 \quad \checkmark \quad \checkmark \end{aligned}$$

(c) Determine the time the cyclists are closest together and determine the minimum distance between them.

at  $t = 7 \text{ sec}$  ✓ the minimum distance is 130 metres. ✓

(2 marks)

(c) Determine the approximate change in volume as it increases from 3 to 3.01 sec. (3 marks)

(b) Determine the exact volume of the balloon 3 seconds after first being inflated. (1 mark)

$$(a) \text{ Show that the relationship between volume } (V \text{ cm}^3) \text{ and time is given by } V = \frac{4\pi r^2 t}{3} \quad (1 \text{ mark})$$

$$r = -(t - 10)$$

In the first five seconds of inflation, the relationship between the radius (cm) and time (t sec) of a spherical party balloon are related by the formula

Question 10

**SECTION TWO****6****MATHEMATICS 3C  
CALCULATOR ASSUMED****Question 11****(5 marks)**

Consider the function  $f(x) = x^3 + ax^2 + 2x + b$  where  $a$  and  $b$  are constants

- (a) Find an expression for the gradient of the curve

(1 mark)

- (b) Given that the tangents at  $A(0, b)$  and  $B(2, 5)$  are parallel, find the value of  $a$  and  $b$ .

(4 marks)

**SECTION TWO****11****MATHEMATICS 3C  
CALCULATOR ASSUMED****Question 17****(8 marks)**

A manufacturer produces 100mm metal rods for sale to another firm. Investigations revealed that the rod lengths are normally distributed with a mean of 100.2mm and a variance of 0.16mm.

The receiving firm will only accept a rod if its length is between 99mm and 101mm

- (a) What proportion of rods will be accepted by the receiving firm?

(1 marks)

$$P(99 < X < 101) \approx 0.9759 \text{ (with the SD at 0.4)} \quad \checkmark$$

- (b) If a rod has been accepted by the receiving firm, what is the probability it was less than 100mm?

(2 marks)

$$P(X < 100 | 99 < X < 101) = \frac{P(99 < X < 100)}{0.9759} \approx \frac{0.3072}{0.9759} \quad \checkmark$$

$$\approx 0.3148 \quad \checkmark$$

- (c) What is the length exceeded by 95% of all rod lengths produced by the manufacturer?

(2 marks)

$$P(X > k) = 0.95$$

$$k \approx 99.5421 \quad \checkmark \quad \checkmark$$

To reduce the number of rejected rods the manufacturer wishes to reduce the variability of the lengths.

- (d) If 99% of the rods must have a length within 0.5mm of the mean (100.2mm), determine the standard deviation length.

(3 marks)

$$P(-\infty < Z < k) = 0.995 \quad \checkmark$$

$$k \approx 2.575829 \quad \checkmark$$

$$\frac{100.7 - 100.2}{\sigma} = 2.575829$$

$$\sigma \approx 0.1941 \quad \checkmark$$

DO NOT WRITE IN THIS AREA

**SECTION TWO**

**MATHEMATICS 3C**

**10**

**CALCULATOR ASSUMED  
MATHEMATICS 3C**

**8 marks)**

- (a) At 9.00am a patient is injected with the serum. At what time, are half of the infected cells expected to be exposed to the serum? Justify your answer.
- The markers of a new serum have reported that when injected into damaged cells the serum doubles every hour until all damaged cells have been exposed to the serum after 24 hours.

- (b) Medical instruments cannot identify any visible change in the damaged cells when less than 1% of the damaged cells have been exposed to the serum. If the serum was injected into the damaged cells at 9.00am, at what time would medical instruments detect the exposure of the serum on the damaged cells? (6 marks)

$$\text{The exponential growth function is } G = G_0 e^{kt}$$

$$\text{Because it doubles every hour } 2 = e^k \quad \checkmark$$

$$t \approx 17.36 \text{ hours} \quad \checkmark$$

At approximately 2.22 am the following day.  $\checkmark$

When 1% of the damaged cells have been exposed to the serum

$$\frac{I_0 e^{0.6931 \cdot 24}}{I_0 e^{0.6931 t}} = 0.01 \quad \checkmark$$

When 1% of the damaged cells have been exposed to the serum

- (a) Poorly behaved (Group C consisting of 5 students)  
 Students could be classified as  
 Well behaved (Group A consisting of 15 students)  
 Moderately behaved (Group B consisting of 10 students)  
 Poorly behaved (Group C consisting of 5 students)  
 She also mentioned that when there is a full moon on any particular lunar cycle the probability that a student will misbehave one or more times is 0.05, 0.15 and 0.3 for a randomly selected student from Group A, B and C respectively.  
 (a) What is the probability that a randomly selected student will misbehave at least once within a lunar cycle?

- (b) Poorly behaved (Group C consisting of 5 students)  
 Students could be classified as  
 Well behaved (Group A consisting of 15 students)  
 Moderately behaved (Group B consisting of 10 students)  
 Poorly behaved (Group C consisting of 5 students)  
 She also mentioned that when there is a full moon on any particular lunar cycle the probability that a student will misbehave one or more times is 0.05, 0.15 and 0.3 for a randomly selected student from Group A, B and C respectively.  
 (a) What is the probability that a randomly selected student will misbehave at least once within a

DO NOT WRITE IN THIS AREA

**Question 12**

**SECTION TWO**

**MATHEMATICS 3C**

**8 marks)**

- (a) At 9.00am half of them would have been exposed to the serum because all of them would have been exposed to the serum after 24 hours.  $\checkmark$
- If the serum reportedly doubles the number of exposed cells every hour then 23 hours after 9.00am half of them would have been exposed to the serum because all of them would have been exposed to the serum after 24 hours.  $\checkmark$

- (b) When 1% of the damaged cells have been exposed to the serum. If the serum was injected into the damaged cells at 9.00am, at what time would medical instruments detect the exposure of the serum on the damaged cells? (6 marks)

$$\frac{I_0 e^{0.6931 \cdot 24}}{I_0 e^{0.6931 t}} = 0.01 \quad \checkmark$$

At approximately 2.22 am the following day.  $\checkmark$

When 1% of the damaged cells have been exposed to the serum

**Question 16**

**SECTION TWO**

**MATHEMATICS 3C**

**10**

**CALCULATOR ASSUMED  
MATHEMATICS 3C**

**8 marks)**

- (a) At 9.00am a patient is injected with the serum. At what time, are half of the infected cells expected to be exposed to the serum? Justify your answer.
- The markers of a new serum have reported that when injected into damaged cells the serum doubles every hour until all damaged cells have been exposed to the serum after 24 hours.

- (b) Poorly behaved (Group C consisting of 5 students)  
 Students could be classified as  
 Well behaved (Group A consisting of 15 students)  
 Moderately behaved (Group B consisting of 10 students)  
 Poorly behaved (Group C consisting of 5 students)

**SECTION TWO****8****MATHEMATICS 3C  
CALCULATOR ASSUMED****Question 13****(8 marks)**

In a packet of multi-coloured Statice seeds (a flower) only one in fifteen seeds will germinate to have a yellow flower.

(a) In a trial 500 seeds germinated.

(i) How many might we expect (mean) to have a yellow flower? (1 mark)

(ii) How many yellow flowered germinated seeds might we expect within two standard deviation of the mean. (2 marks)

(b) Consider a home gardener who has germinated 20 seeds. What is the probability that

(i) exactly two of the 20 seeds will have yellow flowers? (1 mark)

(ii) more than three germinated seeds will have a yellow flower? (1 mark)

(c) How many germinated seeds are required if a home gardener wants to be at least 95% confident that at least one germinated seed will have a yellow flower? (3 marks)

**SECTION TWO****9****MATHEMATICS 3C  
CALCULATOR ASSUMED****Question 15****(8 marks)**

A function is defined as  $y = pxe^{qx}$  where  $p$  and  $q$  are constants.

(a) Determine  $\frac{\delta\psi}{\delta\xi}$  and  $\frac{\delta^2\psi}{\delta\xi^2}$ . (3 marks)

$$\frac{\delta\psi}{\delta\xi} = \pi e^{\theta\xi} + \pi\theta\xi e^{\theta\xi} \quad \checkmark$$

$$\frac{\delta^2\psi}{\delta\xi^2} = 2\pi\theta e^{\theta\xi} + \pi\theta^2\xi e^{\theta\xi} \quad \checkmark \quad \checkmark$$

(b) Using the results found in (a), determine the values for  $p$  and  $q$  so that  $y$  has a maximum of 1 when  $x = \frac{1}{2}$  (5 marks)

$$0.5pe^{0.5q} = 1 \quad \checkmark \text{ and } pe^{0.5q} + 0.5pqe^{0.5q} = 0 \quad \checkmark \quad \checkmark$$

Solving simultaneous equations

$$p \approx 5.4366 \quad q = -2 \quad \checkmark \quad \checkmark$$

DO NOT WRITE IN THIS AREA



**Question 15**

(8 marks)

A function is defined as  $y = pxe^{qx}$  where  $p$  and  $q$  are constants.

- (a) Determine  $\frac{\delta \psi}{\delta \xi}$  and  $\frac{\delta^2 \psi}{\delta \xi^2}$ . (3 marks)

- (b) Using the results found in (a), determine the values for  $p$  and  $q$  so that  $y$  has a maximum of 1

when  $x = \frac{1}{2}$  (5 marks)

DO NOT WRITE IN THIS AREA

$$2 \text{ SD is } 2 \times \sqrt{500 \cdot \frac{1}{14} \cdot \frac{15}{14}} \approx 11.16 \text{ either side of } 33$$

$$\text{binomialPDF}(2, 20, \frac{1}{14}) \approx 0.2439$$

$$\text{binomialCDF}(4, 20, 20, \frac{1}{14}) \approx 0.0405$$

## Question 16

(a) At 9.00am a patient is injected with the serum. At what time, are half of the infected cells

expected to be exposed to the serum? Justify your answer.

(b) Medical instruments cannot identify any visible change in the damaged cells when less than

1% of the damaged cells have been exposed to the serum. If the serum was injected into the

damaged cells at 9.00am, at what time would medical instruments detect the exposure of the

serum on the damaged cells?

DO NOT WRITE IN THIS AREA

(a) The marks of a new serum have reported that when injected into damaged cells the serum

spreads doubles every hour until all damaged cells have been exposed to the serum after 24 hours.

The marks of a new serum have reported that when injected into damaged cells the serum

spreads doubles every hour until all damaged cells have been exposed to the serum after 24 hours.

The marks of a new serum have reported that when injected into damaged cells the serum

spreads doubles every hour until all damaged cells have been exposed to the serum after 24 hours.

(b) The marks of a new serum have reported that when injected into damaged cells the serum

spreads doubles every hour until all damaged cells have been exposed to the serum after 24 hours.

(c) The marks of a new serum have reported that when injected into damaged cells the serum

spreads doubles every hour until all damaged cells have been exposed to the serum after 24 hours.

(d) Medical instruments can only detect the presence of damaged cells when less than

1% of the damaged cells have been exposed to the serum. If the serum was injected into the

damaged cells at 9.00am, at what time would medical instruments detect the exposure of the

serum on the damaged cells?

(2 marks)

(1 mark)

(1 mark)

(4 marks)

(1 mark)

**SECTION TWO****12****MATHEMATICS 3C  
CALCULATOR ASSUMED****Question 17****(8 marks)**

A manufacturer produces 100mm metal rods for sale to another firm. Investigations revealed that the rod lengths are normally distributed with a mean of 100.2mm and a variance of 0.16mm.

The receiving firm will only accept a rod if its length is between 99mm and 101mm

- (a) What proportion of rods will be accepted by the receiving firm? (1 marks)

- (b) If a rod has been accepted by the receiving firm, what is the probability it was less than 100mm? (1 marks)

- (c) What is the length exceeded by 95% of all rod lengths produced by the manufacturer? (2 marks)

To reduce the number of rejected rods the manufacturer wishes to reduce the variability of the lengths.

- (d) If 99% of the rods must have a length within 0.5mm of the mean (100.2mm), determine the standard deviation length. (4 marks)

**SECTION TWO****5****MATHEMATICS 3C  
CALCULATOR ASSUMED****Question 12****(5 marks)**

A mathematics teacher, in conversation with a colleague explained that her Year 10 class of students could be classified as

Well behaved (Group A consisting of 15 students)

Moderately behaved (Group B consisting of 10 students)

Poorly behaved (Group C consisting of 5 students)

She also mentioned that when there is a full moon on any particular lunar cycle the probability that a student will misbehave one or more times is 0.05, 0.15 and 0.3 for a randomly selected student from Group A, B and C respectively.

- (a) What is the probability that a randomly selected student will misbehave at least once within a lunar cycle?

A tree diagram would assist.

(2 marks)

$$\begin{aligned} & \frac{1}{2} \cdot 0.05 + \frac{1}{3} \cdot 0.15 + \frac{1}{6} \cdot 0.3 \quad \checkmark \\ & = \frac{1}{8} = 0.125 \quad \checkmark \end{aligned}$$

- (b) If a randomly selected student had misbehaved at least once during a lunar cycle, what is the probability the student was from Group C? (3 marks)

$$\Pi(X | M) = \frac{\Pi(X \cap M)}{\Pi(M)} \quad \checkmark$$

$$\begin{aligned} & \frac{\frac{1}{6} \cdot 0.3}{\frac{1}{8}} \quad \checkmark \\ & = 0.4 \quad \checkmark \end{aligned}$$

$b=5 \quad \checkmark$  $2^3 - 3 \cdot 2^2 + 2 \cdot 2 + b = 5 \quad \checkmark$ 

$$\begin{aligned} a &= -3 \quad \checkmark \\ f(2) &= 5 \end{aligned}$$

$$\begin{aligned} 2 &= 12 + 4a + 2 \quad \checkmark \\ f(0) &= f(2) \end{aligned}$$

$f(x) = 3x^2 + 2ax + 2 \quad \checkmark$

$$\begin{aligned} \text{Given that the tangents at } A(0, b) \text{ and } B(2, 5) \text{ are parallel, find the value of } a \text{ and } b. \\ \text{Find an expression for the gradient of the curve} \end{aligned}$$

- (b) Given that the tangents at  $A(0, b)$  and  $B(2, 5)$  are parallel, find the value of  $a$  and  $b$ .
- (4 marks)

(a) Find an expression for the gradient of the curve

$$f(x) = x^3 + ax^2 + 2x + b$$

where  $a$  and  $b$  are constants

Consider the function

**Question 18**

**CALCULATOR ASSUMED**

**MATHEMATICS 3C**

- Two competing cyclists are riding with constant speed. At 12 midday cyclist X is 40 metres north of a judge and is riding east at 9 m/s, while cyclist Y is 70 metres east of the judge and is riding north at 7 m/s.
- (a) Show diagrammatically this situation (a scale diagram is not required) (1 mark)
- (b) If the distance between the cyclist  $t$  seconds later is  $D$  metres, show that
- $$D^2 = 6500 - 1820t + 130t^2$$
- (3 marks)
- (c) Determine the time the cyclists are closest together and determine the minimum distance between them. (2 marks)


**Question 19**

The cross section of land can be modeled by the equation

$$H = 0.00003d(d - 30)(d - 50)(100 - d)$$

where  $H$  and  $d$  are, respectively, the height (in metres) above a fixed horizontal level and the distance (in metres) from a fixed point.

The cross section has been shown in the diagram below.



A tunnel, 10m high, will be constructed through the two hills. Show how the cross sectional area of soil removed can be determined using integrals and mensuration (measurement) formula. There is no need to evaluate your answer.

(5 marks)

**Question 10**

In the first five seconds of inflation, the relationship between the radius ( $r$  cm) and time ( $t$  sec) of a spherical party balloon are related by the formula

$$r = -t(t - 10)$$

(a) Show that the relationship between volume ( $V$   $\text{cm}^3$ ) and time is given by  $V = \frac{4\pi(10t - t^2)^3}{3}$

$$r = (10t - t^2) \text{ and } V = \frac{4\pi r^3}{3} = \frac{4\pi(10t - t^2)^3}{3} \quad \checkmark \quad (1 \text{ mark})$$

(b) Determine the exact volume of the balloon 3 seconds after first being inflated. (1 mark)

$$12348\pi \text{ cm}^3 \quad \checkmark$$

(c) Determine the approximate change in volume as  $t$  increases from 3 to 3.01 sec. (3 marks)

$$\delta V = \frac{\partial V}{\partial t} \cdot \delta t$$

$$= 4\pi(10t - t^2)^2(10 - 2t) \times 0.01 \text{ (at } t = 3) \quad \checkmark \quad \checkmark$$

$$= 70.56\pi \text{ cm}^3 \quad \checkmark$$

DO NOT WRITE IN THIS AREA

Question 9

Question 9

$$f(x) = e^x \text{ and } g(x) = 2x - 6$$

$$f(x) = e^x \text{ and } g(x) = 2x - 6$$

$$(b) \quad \text{If } f(x) = 3x^2 - 2 \text{ and } h(x) = \frac{1}{3} - \frac{5}{x}, \text{ find } h(f(x)).$$

$f(x) = e^{-x}$  and  $g(x) = x$   $\vee$  (Alternatives possible)

functions  $f(x)$  and  $g(x)$ . (2 marks)

(a) The composite function  $f(g(x)) = e^{2x-6}$ . Determine two different pairs of equations for

(4 marks)

DO NOT WRITE IN THIS AREA

Domain  $x \geq 3$  ✓ Range  $y \geq -4$  ✓

and range of this function for  $x$  real.

(c) A composite function is defined by the equation  $h(f(x)) = \sqrt[3]{x} - 3$ .  
 4. Determine the domain and range of this function for  $x$  real.

$$\wedge \frac{c^5 - 1}{1} = ((x)_f)h$$

(b) If  $f(x) = 3x^2 - 2$  and  $h(x) = \frac{1}{3} \ln(3f(x))$ .  
 (1 mark)

$f(x) = e^{-x}$  and  $g(x) = x$   $\vee$  (Alternatives possible)

$$f(x) = e^x \text{ and } g(x) = 2x - 6$$

functions  $f(x)$  and  $g(x)$ . (2 marks)

(a) The composite function  $f(g(x)) = e^{2x-6}$ . Determine two different pairs of equations for

(4 marks)

$$\text{Domain } x \geq 3 \quad \checkmark \quad \text{Range } y \geq -4 \quad \checkmark$$

and range of this function for  $x$  real.

(c) A composite function is defined by the equation  $h(f(x)) = \sqrt[3]{x} - 3$ .  
 4. Determine the domain and range of this function for  $x$  real.

(8 marks)

A basketball training squad consists of 4 guards, 2 centres and 3 forwards. A team of 5 is to be chosen to start the game.

In how many ways can this starting team be chosen if:

- (a) there are no restrictions? (1 mark)

$${}^9X_5 = 126 \quad \checkmark$$

- (b) the team must consist of 2 guards, 1 centre and 2 forwards? (2 marks)

$${}^4X_2 \cdot {}^2X_1 \cdot {}^3X_2 = 36 \quad \checkmark \quad \checkmark$$

- (c) the team includes at most 2 centres? (3 marks)

$${}^2X_0 \cdot {}^7X_5 + {}^2X_1 \cdot {}^7X_4 + {}^2X_2 \cdot {}^7X_3 = 126 \quad \checkmark \quad \checkmark \quad \checkmark$$

Julie is a centre player and is chosen in the team. If the other players are selected at random, what is the probability that

- (d) Julie is the only centre in the team? (2 marks)

$$\frac{{}^1X_1 \cdot {}^7X_4}{{}^9X_5} = \frac{5}{18} \quad \checkmark \quad \checkmark$$

DO NOT WRITE IN THIS AREA