

SOLUTIONS

2021

**MATHEMATICS
METHODS
UNITS 3 & 4**

SEMESTER TWO



Copyright for test papers and marking guides remains with West Australian Test Papers.
Test papers may only be reproduced within the purchasing school according to the advertised Conditions of Sale.
Test papers should be withdrawn after use and stored securely in the school until 14th October 2021.

Calculator-free Solutions

1. (a)
- $(2\pi, 0)$

$$\frac{dy}{dx} = \frac{2\tan(x)}{x} + \frac{2\ln x}{\cos^2(x)}$$

✓

$$\text{When } x = 2\pi, m = 0 + 2\ln 2\pi$$

✓

$$0 = 2\ln 2\pi(2\pi) + c \therefore c = -4\pi \ln 2\pi$$

$$y = (2\ln 2\pi)x - 4\pi \ln 2\pi$$

✓

$$(b) 2t \ln(t^4) \tan(t^2)$$

✓✓

[5]

$$2. \quad (a) \quad \frac{1+2+3+4+5}{k} = 1$$

✓

$$k = 15$$

$$(b) \quad \frac{12}{15} = \frac{4}{5}$$

✓

$$(c) \quad \frac{1+4+9+16+25}{15} = \frac{55}{15} = \frac{11}{3}$$

✓

$$(d) \quad (i) \quad \text{Expected value } \left(\frac{1}{10}\right)\left(\frac{11}{3}\right) + 2 = \frac{71}{30}$$

✓

$$\frac{1+8+27+64+125}{15} - \frac{121}{9} = \frac{14}{9}$$

Variance of Y :

$$\frac{\sqrt{14}}{3}$$

$$\text{Standard deviation of } Y = \frac{\sqrt{14}}{3}$$

✓

$$\frac{\sqrt{14}}{30}$$

$$\text{Standard deviation of } X = \frac{\sqrt{14}}{30}$$


✓

$$(ii) \quad P\left(X \leq \frac{11}{5}\right) = P(Y \leq 2) = \frac{3}{15}$$

✓✓

[8]

3. (a) $g''(x) = -2x + 1$ when $x > \frac{1}{2}$ $g(x)$ is concave down
 (b) Maximum
 (c)



[3]



$$k = 8 \quad n = \frac{1}{4}$$



$$y = 8x^{\frac{1}{4}}$$

$$\log_2 y = \log_2 8x^{0.25}$$



$$\log_2 y = \frac{4}{1} \log_2 x + 3 \log_2 2$$

6.

$$\log_2 y = \frac{4}{1} \log_2 x + 3$$

[4]



$$\frac{0.5}{0.16} = 0.32$$



(c)



(b)

0.34



(a)

0.16

[7]



$$\therefore x = 8$$



$$(x - 8)(x + 2) = 0$$



$$x^2 - 6x - 16 = 0$$



$$\log_6(x^2 - 6x) = 2$$



(c)



(b)

= 3

$$= 3\log_6 5 + \log_6 2 - \frac{3}{1}(3)\log_6 2$$



$$2a - 1 = 0 \quad \therefore a = \frac{1}{2}$$



$$p(x) = 2a^x - 1 \quad \text{given } (1, 0)$$



$$0 = \log_6 x \quad \therefore x = 1 \quad \leftarrow C(1, 0)$$



$$b = -1$$



$$c_{-1} = 2 \quad \therefore c = \frac{1}{2}$$

4.

$$(a) \quad -1 = \log_6(2)$$

[5]



Shape showing local minimum and maximum

End of Questions

End of Questions

$$m'(x) = \frac{(x^2 - 1)(2xe^{x^2-1}) - (e^{x^2-1})(2x)}{(x^2 - 1)^2}$$

✓

$$= \frac{2xe^{x^2-1}(x^2 - 2)}{(x^2 - 1)^2}$$

Stationary points occur when $2xe^{x^2-1}(x^2 - 2) = 0$

✓

$$\therefore x = 0, \pm\sqrt{2}$$

✓

$$\therefore \left(0, -\frac{1}{e}\right) (\sqrt{2}, e) (-\sqrt{2}, e)$$

✓

[4]

$$p(x) = \begin{cases} \frac{1}{20} & 0 \leq x \leq 20 \\ 0 & x > 20 \end{cases}$$

✓

$$\frac{2}{20} = \frac{2}{17}$$

- (b) $\frac{17}{20}$
(c) 10 minutes

✓✓

✓

[4]

$$\int_{-1}^1 k(1-x^2) - 2k(x^2-1) dx = 3k \int_{-1}^1 (1-x^2) dx$$

✓

$$3k \left[x - \frac{x^3}{3} \right]_{-1}^1 = 8$$

✓

$$4k = 8$$

$$k = 2$$

✓

[3]

$$10. (a) 0 < x \leq 1$$


✓✓

$$(b) \left(\frac{4}{3}\right)^p = \frac{16}{9}$$

✓

$$(c) p = 2$$

✓



$$21. (a) p = \frac{28}{80}$$

$$(b) (i) (0.2455, 0.4545)$$

✓

✓✓

$$1.960 \sqrt{\frac{28 \left(\frac{52}{80} \right)}{80}} = 0.1045$$

✓

$$(ii) ME =$$

(iii) 95% the width is 0.209
99% the width is 0.2747

$$0.209x = 0.2747$$

$$x = 1.3144$$

Therefore an increase of 31.45%

$$1.960 \sqrt{\frac{28 \left(\frac{52}{80} \right)}{80}} = 0.02$$

✓

$$(c) n = 2184.9$$

✓

Therefore sample size of 2185

✓

$$(d) 0.95 \times 40 = 38$$

∴ Approximately 38 would contain the true proportion

✓

[10]

$$22. (a) -192\cos(4t) - 80\sin(4t) = 0$$

$$t = 1.27679 \text{ (max)}$$

✓

$$v(t) = 20\cos(4t) - 48\sin(4t) + c$$

$$v(t) = 20 \text{ when } t = 0 \text{ therefore } c = 0$$

✓

$$v(1.27679) = 52 \text{ units/sec}$$

✓

$$(b) v(2) = -50.4 \text{ units/sec}$$

✓

$$a(2) = -51.2 \text{ units/sec}^2$$

✓

The velocity and the acceleration are both negative, therefore the particle is moving faster to the left.

∴ Increasing speed

✓

$$(c) x(t) = 5\sin 4t + 12\cos 4t + c$$

✓

$$x(0) = 12 \therefore c = 0$$

✓

$$x(2) = 3.2008 \text{ units}$$

✓

$$(d) \int_4^5 20\cos(4t) - 48\sin(4t) dt = 22.393 \text{ units}$$

✓✓

$$(e) \int_0^5 |20\cos(4t) - 48\sin(4t)| dt = 160.538 \text{ units}$$

✓✓

[12]

¹⁴ mature trees can be used for luxury furniture.

18. Over - estimate = $0.2(0.127+0.172+0.184+0.184+0.181) = 0.1696$
Under - estimate = $0.2(0+0.127+0.172+0.181+0.173) = 0.1306$
Average = 0.1501 units^2

19. (a) (i) $n > 30$ (b) $np = 300 \times 0.12 = 36$ (c) $npq = 300 \times 0.12 \times 0.88 = 31.68$
 $np > 10$, $npq > 10$ therefore a normal distribution can apply.

10. (c) (iii) $\log_{\frac{1}{4}} x = \ln \frac{3}{x}$

[3]

(iii) $P(p > 0.125) = 0.3949$
 Calculator-Assumed Solutions

11. (a) $M_p(t) = M^*(t) - M_o(t) = -0.4t + 8 - 0.3t - 2$
 $P(p < k) = 0.25 \quad k = 0.1073$
 10.7%
 $\int_{18}^{15} \frac{1}{x} dx = \frac{13}{15}$
 $\text{Expected waiting time} = 12.5 \text{ days}$

(b) $M^*(t) = -0.7t + 6$
 $M_p(t) = -0.7t + 6 = 0$
 $\text{Maximum profit occurs when } M_p(t) = 0$
 $\text{Therefore } t = 8.57 \text{ years}$
 $\int_0^{8.57} -0.7t + 6 dt = 25.714285$
 $\text{Maximum profit} = \$25.714285 - \$12000000 = \13.714285

[4]

Normal distribution $p \sim N(0.5, 0.07906^2)$

$$P(p < 0.7) = 0.9943 \quad \text{(iii)}$$

$$\begin{aligned} \text{(b) (i)} & \quad pH = -\log_{10} [H_3O^+] \\ & \quad [H_3O^+] = 10^{-4.5} = 0.00003162 \text{ moles per litre} \\ & \quad \text{Hydronium ions range from } 10^{-14} \text{ to } 10^{-1 \text{ to } 14} \\ & \quad \text{Therefore the maximum value of } n = 70 \\ & \quad 0.987^n \leq 0.4 \quad \therefore n \leq 70.025 \\ & \quad P(Y = 0) \geq 0.4 \\ & \quad 1 - P(X = 0) \leq 0.6 \\ & \quad Y \sim \text{Bin}(n, 0.013) \quad P(Y \geq 1) \leq 0.6 \\ & \quad \text{Standard deviation} = 3.6142 \\ & \quad \text{Variance} = 275 \times 0.05 \times 0.95 = 13.0625 \\ & \quad \therefore 13 \text{ drivers.} \end{aligned}$$

$$[\text{H}_3\text{O}^+] = 10^{-4.7} = 0.000019953 \text{ moles per litre}$$

13. Point of intersection:

$$x^3 - 4x^2 + 3x + 1 = x^2 - 3x + 1$$

 $x = 0$ or 2 or 3

Area of shaded part:

$$\int_0^3 |(x^3 - 4x^2 + 3x + 1) - (x^2 - 3x + 1)| dx = \frac{37}{12} \text{ units}^2$$

✓✓ Alternate

Shaded area under line:

$$\int_0^2 1 - x - (x^2 - 3x + 1) dx = \frac{4}{3} \text{ units}^2$$

$$\frac{16}{37}$$

Fraction: $\frac{16}{37}$

✓

[4]

14. (a) $AC = t - 30 \therefore p^2 = 30^2 - (t - 30)^2$

✓

$$p^2 = -t^2 + 60t$$

✓

$$\frac{1}{3}\pi r^2 h$$

Volume of cone =

$$V(t) = \frac{1}{3}\pi (-t^2 + 60t)(t) = -\frac{\pi t^3}{3} + 20\pi t^2$$

✓

(b) $V'(t) = 40\pi t - \pi t^2 = 0$

✓

$$t = 0 \text{ or } 40$$

✓

$$V''(t) = 40\pi - 2\pi t$$

✓

 $V''(40) = -40\pi \therefore$ maximum at $t = 40$ cm

✓

$$\frac{32000\pi}{3} = 33510.322 \text{ cm}^3$$

(c) Volume of cone at $t = 40$ cm is

✓

Volume of sphere = $113\ 097.336 \text{ cm}^3$

✓

Percentage = 29.63%

✓

(d) $\frac{\delta r}{r} = -0.015 \quad \frac{dV}{dr} = 4\pi r^2 \quad \frac{\delta V}{\delta r} \approx \frac{dV}{dr}$

$$\frac{\delta V}{V} \approx \frac{4\pi r^2}{\frac{4}{3}\pi r^3} \times \delta r$$

✓

$$\frac{\delta V}{V} \approx \frac{3\delta r}{r}$$

✓

$$\frac{\delta V}{V} \approx 3(-0.015)$$

✓

Approximately 4.5% decrease in volume.

✓

[11]

$$50 = 100(1 - e^{3k})$$

$$k = -0.23105$$

✓

15. (a) $P = 100(1 - e^{5 \times -0.23105}) = 68.5\%$

✓

Percentage 5 minutes or longer = 31.5%

✓

(c) (i) t is a continuous random variable where $0 \leq t \leq 30$.The function in the domain $0 \leq t \leq 30$ is positive.The probability at $t = 0$ is 0 and $t = 30$ is 1, therefore is cumulative.

✓✓

(ii) $P(t) = 1 - e^{-0.23105(15)} = 0.96896$

✓

(iii) $\frac{0.68502}{0.90079} = 0.7605$

✓✓

[8]

16. $T(t) = 21e^{-kt} + 4$

$$9.8 = 21e^{-kt} + 4$$

✓

$$6.5 = 21e^{-k(t+15)} + 4$$

✓

$$k = 0.056104 \quad t = 22.9334$$

✓

The liquid was placed in the fridge at 11:37 am

✓

[4]

17. (a) 6 year old tree is growing at a rate of 17.69 cm/year

✓

50 year old tree is growing at a rate of 2.277 cm/year

✓

7.8 times faster

✓

(b) (i) Convenience sample: this sample may not be representative of all the six year old trees in the plantation.

✓✓

The sample is not large enough.

(ii) Stratified sample where a number of trees from each area according to the size of the area are chosen at random.

Or

Systematic and Array sample: Every n th tree starting at a randomly assigned tree as one walks down each row.

✓✓

(c) (i) $P(Z > z) = 0.0094 \therefore z = 2.3495$

✓

$$\frac{326 - \mu}{12} = 2.3495$$

✓

$$\mu = 297.8 \text{ cm}$$

✓

(ii) According to the model, the height of a six year old tree is 297.8 cm and the mean of the sample is the same, therefore the model is suitable.

✓

(iii) $X \sim (5, 0.0094) \quad P(X = 0) = 0.95388$

✓

(d) $\frac{132}{400} \pm 2.58 \sqrt{\frac{\left(\frac{132}{400}\right)\left(\frac{268}{400}\right)}{400}} = (0.269, 0.391)$

✓✓

With a 99% confidence level, between 26.9% and 39.1% of the