


Copyright for test papers and marking guides remains with *West Australian Test Papers*.
Test papers may only be reproduced within the purchasing school according to the advertised Conditions of Sale.
Test papers should be withdrawn after use and stored securely in the school until Friday 14th June 2019.



MATHEMATICS METHODS UNIT 3

Semester One

2019

SOLUTIONS

Calculator-free Solutions

1. (a) $y = -(4x + 3)^{-\frac{1}{2}}$
 $\frac{dy}{dx} = \frac{2}{(4x + 3)^{\frac{3}{2}}}$ ✓
 $m = 2 \quad n = \frac{3}{2}$ ✓
 $\frac{2}{(9)^{\frac{3}{2}}}$
 (b) $= \frac{2}{27}$ ✓ [3]
2. (a) $f(x) = x^4 - 4x^3$
 $f'(x) = 4x^3 - 12x^2$
 $4x^2(x - 3) = 0$ ✓
 $x = 0$ or $x = 3$ ✓
 Stationary points (0, 0) and (3, -27) ✓
 (b) $f''(x) = 12x^2 - 24x$ ✓
 $f''(0) = 0$ ∴ horizontal point of inflection occurs at (0, 0) ✓
 Since $f''(3) > 0$, then minimum occurs at (3, -27) ✓✓
- (c) (i) $f''(x) = 12x^2 - 24x$
 $12x(x - 2) = 0$
 $\therefore x = 0$ or $x = 2$ ✓
 Point of oblique inflection at (2, -16) ✓
 (ii) $m = -16$
 $y = -16x + c$ ✓
 $-16 = -16(2) + c$
 $c = 16$
 Tangent is $y = -16x + 16$ ✓ [11]
3. (a) $y = e^{-12x}$
 $\frac{dy}{dx} = -\frac{12}{e^{12x}}$ ✓✓
 (b) $-2\sin x e^{\cos x}$ ✓✓ [4]

4. (a) The values of $f(x)$ are all positive ✓

$$\frac{5}{15} + \frac{4}{15} + \frac{3}{15} + \frac{2}{15} + \frac{1}{15} = 1$$
 ✓
 \therefore Probability function

$$P(X < 3) = \frac{9}{15}$$
 ✓
 (b)
$$\frac{\frac{5}{15}}{\frac{14}{15}} = \frac{5}{14}$$
 ✓
 (c) ✓✓ [5]
5.
$$g(x) = -\cos x + \frac{1}{2} \sin 2x + c$$
 ✓

$$1 = -\cos\left(\frac{\pi}{2}\right) + \frac{1}{2} \sin \pi + c$$
 ✓

$$1 = c$$

$$g(x) = -\cos x + \frac{1}{2} \sin 2x + 1$$
 ✓ [3]
6. (a)
$$f'(x) = \frac{(1 - e^{2x})(-\sin x) - \cos x(-2e^{2x})}{(1 - e^{2x})^2}$$
 ✓✓
 (b)
$$\frac{dy}{dx} = 6x \sin^4(2x - 3) + 24x^2 \cos(2x - 3) \sin^3(2x - 3)$$
 ✓✓ [4]
7. (a) (i) -3 ✓
 (ii) $6 + 2 = 4$ ✓✓
 (iii) 5 units² ✓✓
 (b)
$$x = 2e^{2t} + \frac{2}{3}(t + 1)^{\frac{3}{2}} + c$$
 ✓

$$1 = 2 + \frac{2}{3} + c$$
 ✓

$$c = -\frac{5}{3}$$

$$x = 2e^{2t} + \frac{2}{3}(t + 1)^{\frac{3}{2}} - \frac{5}{3}$$
 ✓ [8]

$$(a) \quad E(X^2) - [E(X)]^2 = \frac{1}{2}$$

8.

$$2p + 4p - (2p + 2p)^2 = 6p - 16p^2 = \frac{1}{2}$$

✓

$$(32p^2 - 12p + 1) = 0$$

$$\therefore (8p - 1)(4p - 1) = 0$$

$$\therefore p = \frac{1}{8} \text{ or } \frac{1}{4}$$

✓

$$\therefore E(X) = \frac{1}{2} \text{ or } 1$$

✓✓

$$(b) \quad (i) \quad 21$$

✓

$$(ii) \quad \text{Standard deviation of } X = 3$$

$$|2| \times 3 = 6$$

✓✓

[7]

9.

$$(a) \quad \left[\pi \sqrt{2x+3} \right]_1^3$$

✓

$$= 3\pi - \pi = 2\pi$$

✓

$$y = -\frac{2}{\pi} \cos \frac{\pi x}{2} + x + c$$

(b)

✓

$$4 = 0 + 1 + c \quad \therefore c = 3$$

✓

$$\text{Its path is } y = -\frac{2}{\pi} \cos \frac{\pi x}{2} + x + 3$$

✓

[5]

Calculator-assumed Solutions

$$10. \quad (a) \quad A = \pi r^2 = \pi(3t + 1)^2$$

✓

$$(b) \quad \frac{dr}{dt} = 3 \text{ cm/s}$$

✓

$$(c) \quad \frac{dA}{dt} = 6\pi(3t + 1) \big|_{t=1}$$

$$= 24\pi \text{ cm}^2/\text{s}$$

✓

$$(d) \quad \delta A = \frac{dA}{dr} \times \delta r$$

$$= 2\pi(r)(0.05)$$

✓✓

$$\text{When } r = 4, \delta A = 0.4\pi \approx 1.26 \text{ cm}^2$$

✓

[6]

$$k + k + 2k + \frac{3}{2}(2k) + 3k + 4k = 1$$

11. (a) Let $P(X = 0) = k$ then

$$k = \frac{1}{14}$$

✓

X	0	1	2	3	4	5
$P(X = x)$	$\frac{1}{14}$	$\frac{1}{14}$	$\frac{2}{14}$	$\frac{3}{14}$	$\frac{3}{14}$	$\frac{4}{14}$
Frequency	23	23	46	69	69	92

✓✓✓✓

(b) mode = 5

✓

$$\text{mean} = \frac{\sum x \cdot P(X = x)}{7} = \frac{23}{7} \approx 3.2857$$


✓

(c) $\frac{5}{14}$

✓


[7]

12. (a)



✓✓✓✓

(b)



✓✓✓

(c) $F'(x) = 2(2x + 3) \sin(2x + 3)$

✓

[6]

13. (a) $f'(t) = \frac{2}{50} f(t) = 0.04 f(t)$
 $\therefore k = 0.04$ ✓
 $2A = Ae^{0.04t}$ ✓
 $t = 17.3$ days ✓
- (b) (i) $\int_0^8 2e^{2t-7} dt = 8103.083 \text{ m}^2$ ✓
- (ii) $\int_9^{10} 2e^{2t-7} dt = 382\,539.25 \text{ m}^2$ ✓
- (iii) The exponential growth of the area becomes too large too quickly for the model to be realistic. ✓ [6]
14. (a) $s'(t) = v(t) = -3t^2 + 27 = 0$ ✓
 $\therefore t = 3$ (discard -3) ✓
The maximum distance from O occurs at $t = 3$ seconds and then the particle turns back towards O.
- (b) $s(t) = 27t - t^3 = 0$
 $\therefore t = 3\sqrt{3}$ (discard $t = 0$ and $-3\sqrt{3}$) ✓
 $\therefore v(3\sqrt{3}) = -54$ ✓
Therefore speed when the particle returns to O is 54 cm/s. ✓ [5]
15. $\left[x^2 \sqrt{1-x^2} \right] \frac{1}{2} = t^2 \sqrt{1-t^2} - \frac{1}{4} \sqrt{\frac{3}{4}}$ ✓✓
- $A(t) = t^2 \sqrt{1-t^2} - \frac{\sqrt{3}}{8}$ ✓ [3]
16. (a) Profit = Revenue - Cost
 $C = 50 + 0.8n$ and $R = n(3.5 - 0.01n)$ ✓
 $P = 3.5n - 0.01n^2 - 50 - 0.8n = 2.7n - 0.01n^2 - 50$ ✓
- (b) $2.5 = 3.5 - 0.01n$ ✓
 $\therefore n = 100$ He will sell 100 figs ✓
 $P'(n) = 2.7 - 0.02n$ where $n = 100$ ✓
The marginal profit is $0.7 = 70$ cents ✓ [6]

17. (a) (i)

Y	0	1	2	3
$P(Y=y)$	$\frac{1}{56}$	$\frac{15}{56}$	$\frac{15}{28}$	$\frac{5}{28}$

✓✓

$$(ii) \quad E(Y) = \frac{105}{56} \approx 1.875$$

✓

$$\text{Var}(Y) = \frac{225}{448} \approx 0.50$$

✓

$$\frac{15}{56}$$

(iii)

✓

$$(b) \quad (i) \quad X \sim \text{Bin}(10, 0.7)$$

$$P(X = 7) = 0.2668$$

✓

$$(ii) \quad X \sim \text{Bin}(26, 0.7)$$

$$P(X \leq 13) = 0.0255$$

✓

$$(c) \quad E(X) = np = 12$$

✓

$$\text{Var}(x) = npq = np(1 - p) = 9$$

✓

$$n = 48 \quad \text{and} \quad p = \frac{1}{4}$$

✓

[10]

18. (a) The amount of water cannot be a negative amount.
If no water flows in or out the functions can equal zero.

✓✓

$$(b) \quad f(t) - w(t) \quad \text{megalitres}$$

✓

$$(c) \quad \int_{t_0}^{t_1} f(t) \, dt$$

✓

$$(d) \quad \int_0^{30} \left(10 - \frac{1}{2}t - 2\sin 2\pi t \right) dt = 75$$


(d)

✓

75 megalitres of water.

✓

(e)



✓✓ [8]

19. (a) (i) Each washing machine is an independent trial. ✓
 There are two outcomes: Success (a washing machine works) and Failure (a washing machine is broken.) ✓
 (ii) $E(X) = p = 0.2$ ✓
 $\text{Var}(X) = p(1 - p) = 0.2 \times 0.8 = 0.16$ ✓
 Standard deviation = 0.4 ✓
 (b) (i) This means that the first mobile phone was selected and found to be defective. (0 non-defective mobile phones were found) ✓
 $P(X = 0) = 0.03$ ✓
 (ii) This means that the first two mobile phones selected were found to be non-defective but the third mobile phone selected was defective. ✓
 $P(X = 2) = 0.97 \times 0.97 \times 0.03 = 0.028227$ ✓
 (iii) $0.97^5 \times 0.03 = 0.02576$ ✓✓
 (iv) $1 - [P(0) + P(1) + P(2)] = 0.91267$ ✓✓ [12]

20. (a) $y = x(x - 1)(x - 3)$
 Graph cuts the x-axis at 0, 1 and 3. ✓
 $\int_0^1 (x^3 - 4x^2 + 3x) dx = \frac{5}{12}$ ✓
 $-\int_1^3 (x^3 - 4x^2 + 3x) dx = \frac{8}{3}$ ✓
 $\text{Area} = \frac{37}{12} = 3\frac{1}{12}$
 (b) $h'(x) = 4e^{2x^2 + x} + c$ ✓
 $h'(0) = 1 \therefore 1 = 4e^0 + c \quad c = -3$ ✓
 $h'(1) = 4e^3 - 3$ which is the gradient at $x = 1$ ✓
 (c) Graphs intersect at $x = 0.6349, 2.5067$ ✓
 $\int_0^{0.6349} \cos 2x - \frac{1}{2} \sin x \, dx + \int_{0.6349}^{2.5067} \frac{1}{2} \sin x - \cos 2x \, dx + \int_{2.5067}^{\pi} \cos 2x - \frac{1}{2} \sin x \, dx$
 $= 2.5203 \text{ units}^2$ ✓✓ [9]
 Or: $\int_0^{\pi} \left| \cos 2x - \frac{1}{2} \sin x \right| dx = 2.5203 \text{ units}^2$

21. In triangle $h = \sqrt{15^2 - x^2}$ therefore $A = \frac{1}{2}bh = x(\sqrt{15^2 - x^2})$ ✓
 $A'(x) = \frac{225 - 2x^2}{\sqrt{225 - x^2}} = 0$ ✓
 $x = \pm \frac{15\sqrt{2}}{2} \approx 10.6066$ (Discard negative value for x) ✓
 $A''(10.6066) = -3.999 \therefore A'' < 0$ therefore maximum ✓
Maximum area is 112.5 cm^2 ✓ [5]

22. (a) $v(t) = \int \cos t - 4\sin(2t) = \sin t + 2\cos 2t + c$ ✓
 $\sin 0 + 2\cos 0 + c = 2 \therefore c = 0$
 $v(t) = \sin t + 2\cos 2t = 0$ ✓
The particle changes direction when $t = 1.003 \text{ s}$ or $t = 2.139 \text{ s}$ ✓
(b) $\int_0^\pi |\sin t + 2\cos 2t| dt = 3.476$ m² ✓✓ [5]

23. (a)
- | $g(0)$ | $g(0.2)$ | $g(0.4)$ | $g(0.6)$ | $g(0.8)$ | $g(1)$ |
|--------|----------|----------|----------|----------|--------|
| 1 | 0.96 | 0.85 | 0.70 | 0.53 | 0.37 |

Area from left = $0.2(1 + 0.96 + 0.85 + 0.70 + 0.53) = 0.808$ ✓
Area from right = $0.2(0.96 + 0.85 + 0.70 + 0.53 + 0.37) = 0.682$ ✓
Average = $0.745 = 0.75 \text{ units}^2$ ✓

- (b) As the width of the rectangle tends to 0, the more accurate will be the area. ✓ [5]

24. (a) $t = 110 \text{ years}$
 $A = e^{-0.047069} = 0.95402$ grams ✓
Therefore 4.598 % has decayed. ✓
(b) $0.5 = e^{-0.0004279t}$ ✓
 $t = 1619.88 \approx 1620 \text{ years}$ ✓
(c) $0.5 = e^{k \times 3.8}$
 $\therefore k = -0.1824$ ✓
 $A = 10e^{-0.1824(15)}$ ✓
0.648 mg of radon remains ✓ [7]

END OF QUESTIONS