



## Examination Question/Answer Booklet

# CHEMISTRY

## STAGE 3AB

### Semester 2 examination

Please place your student identification label in this box

#### ***Time allowed for this paper***

Reading time before commencing work: Ten minutes  
Working time for paper: Three hours

#### **Materials required/recommended for this paper**

##### **To be provided by the supervisor**

Question/answer booklet  
Separate multiple-choice answer sheet  
Data sheet

##### **To be provided by the candidate**

Standard items: Pens, pencils, eraser, correction fluid, ruler, highlighters  
Special items: Scientific

#### ***Important note to candidates***

No other items may be taken into the examination room. It is **your** responsibility to ensure that you do not have any unauthorised notes or other items of a non-personal nature in the examination room. If you have any unauthorised material with you, hand it to the supervisor **before** reading any further.

Section	Out of	
Multiple Choice	/50	
Short Answers	/69	
Extended response	/65	
<b>Total</b>	<b>/184</b>	<b>%</b>

## Structure of this paper

Section	Suggested working time	Number of questions available	Number of questions to be attempted	% of paper	Marks
ONE Multiple choice	50 minutes	25	25	27	50
TWO Short response	60 minutes	13	13	38	69
THREE Extended response	70 minutes	6	6	35	65
Total				100	184

## Instructions to candidates

- The rules for the conduct of Curriculum Council examinations are detailed in the *Student Information Handbook*. Sitting this examination implies that you agree to abide by these rules.
- Answer the questions according to the following instructions.  
**Section One** Answer all questions on the separate multiple-choice answer sheet provided.  
**Section Two** Answer all questions in the spaces provided in this Question/Answer Booklet.  
**Section Three** Answer all questions in the spaces provided in this Question/Answer Booklet.
- A blue or black ballpoint or ink pen should be used.
- For full marks, chemical equations should refer only to those species consumed in the reaction and the new species produced. These species may be **ions** [for example  $\text{Ag}^+$  (aq)], **molecules** [for example  $\text{NH}_3(\text{g})$ ,  $\text{NH}_3(\text{aq})$ ,  $\text{CH}_3\text{COOH}(\text{l})$ ,  $\text{CH}_3\text{COOH}(\text{aq})$ ] or **solids** [for example  $\text{BaSO}_4(\text{s})$ ,  $\text{Cu}(\text{s})$ ,  $\text{Na}_2\text{SO}_4(\text{s})$ ].
- Additional information which may be necessary to answer questions is located on the separate Chemistry data sheet.

**SECTION ONE – MULTIPLE CHOICE****[50 marks]**

This section has **TWENTY FIVE (25)** questions. Attempt **ALL** questions.


Answer all questions in Section 1 on the separate Multiple-Choice Answer Sheet provided, using a blue or black pen. Each question in this part is worth 2 marks.

Suggested working time: 50 minutes


---

1. An atom has the first five successive ionisation energies  
1.1      2.4      4.6      6.2      37.8      MJ mol<sup>-1</sup>  
Which one of the following elements is it?
- a) Aluminium  
b) Argon  
c) Calcium  
d) Carbon
2. Which one of the following pairs of elements is most likely to form an ionic compound?
- a) H and O  
b) H and F  
c) C and F  
d) K and F
3. Which one of the following molecules is non-polar; that is, the molecule has no overall dipole moment?
- a) H<sub>2</sub>CO  
b) CO<sub>2</sub>  
c) H<sub>2</sub>O  
d) NH<sub>3</sub>
4. Which of the following best explain why hexane has a very low solubility in water?
- a) Hexane and water each have strong forces between their molecules.  
b) Like dissolves like, and these two liquids are unlike.  
c) Attractions between hexane and water molecules are much weaker than attractions between water molecules or between hexane molecules.  
d) The covalent bonds in a water molecule are exceptionally strong.
5. The following gaseous equilibrium is established at high temperatures in the presence of a finely divided nickel (Ni) catalyst.
- $$\text{CH}_4(\text{g}) + \text{H}_2\text{O}(\text{g}) \rightleftharpoons \text{CO}(\text{g}) + 3\text{H}_2(\text{g}); \quad \Delta H = +206 \text{ kJ mol}^{-1}$$
- A particular reaction is carried out using equal amounts of CH<sub>4</sub>(g) and H<sub>2</sub>O(g). Which one of the following sets of changes in conditions would lead to the greatest increase in the proportion of the reactant converted to products?
- | Volume of reaction vessel | Temperature |
|---------------------------|-------------|
| a) increased              | increased   |
| b) increased              | decreased   |
| c) decreased              | increased   |
| d) decreased              | decreased   |


6. Equal amounts of  $\text{CH}_4(\text{g})$  and  $\text{H}_2\text{O}(\text{g})$  are added to a reaction vessel and allowed to react. After 10 minutes, equilibrium has been reached. At that time, some  $\text{H}_2$  is added to the mixture and equilibrium is re-established. Which one of the following graphs best represents the changes in the amounts of  $\text{CH}_4$  and  $\text{H}_2$  in the reaction mixture?



7. Hydrogen iodide dissociates into its elements according to the following equation.



A mixture of  $\text{H}_2(\text{g})$ ,  $\text{I}_2(\text{g})$  and  $\text{HI}(\text{g})$  rapidly comes to equilibrium in a 2.0 L container. After the reaction has been at equilibrium for 10 minutes, the volume of the container is suddenly reduced to 1.3 L at constant temperature. Which one of the following graphs best represents the effect of this decrease in volume on the concentration of the gases in the mixture?



8. The two statements below give possible explanations for changes that occur when the temperature of a reaction mixture is increased.
- I At a higher temperature, particles move faster and the reactant particles collide more frequently.
  - II At a higher temperature, more particles have energy greater than the activation energy.


Which alternative below best explains why the observed reaction rate is greater at higher temperatures?

- a) I only
  - b) II only
  - c) I and II to an equal extent
  - d) I and II, but II to a greater extent than I
9. The anaesthetic, nitrous oxide,  $\text{N}_2\text{O}$ , decomposes to form an equilibrium mixture of  $\text{N}_2\text{O}$ ,  $\text{N}_2$  and  $\text{O}_2$  according to the following equation.



At  $25^\circ\text{C}$ ,  $K = 7.3 \times 10^{37}$  M and at  $40^\circ\text{C}$ ,  $K = 2.7 \times 10^{36}$  M  
What valid conclusion can be made from this?

- a) The equilibrium concentration of  $\text{N}_2$  and  $\text{O}_2$  are equal at  $25^\circ\text{C}$ .
  - b) The equilibrium concentration of  $\text{N}_2\text{O}$  is higher at  $25^\circ$  than at  $40^\circ\text{C}$ .
  - c)  $\text{N}_2\text{O}$  is less stable at the higher temperature.
  - d) The forward reaction is exothermic.
10. Which of the following changes will always shift this equilibrium reaction to the right?




- a) Adding a catalyst
  - b) Increasing the pressure
  - c) Increasing the temperature
  - d) Adding more of the reactant
11. A 2.45g sample of lawn fertiliser was analysed for its sulfate content. After filtration and drying, 2.18g of barium sulfate was recovered.

What is the % w/w of sulfate in the lawn fertiliser?


- a) 16.8
- b) 36.6
- c) 46.2
- d) 89.0

12. The apparatus shown is used in a first-hand investigation to determine and compare the heat of combustion of three different liquid alkanols.




Which is the independent variable?


- a) Type of alkanol used
  - b) Amount of water used
  - c) Amount of alkanol used
  - d) Temperature change in the water
13. The Winkler method is used to determine the amount of dissolved oxygen in a sample. In this procedure, oxygen reacts with  $\text{Mn}^{2+}$  under alkaline conditions to produce a precipitate of  $\text{MnO}(\text{OH})_2$ .



The precipitate is then dissolved in acid and reacted with iodide, forming iodine and  $\text{Mn}^{2+}$ .



Finally, the amount of iodine produced is determined by reaction with thiosulfate.



When a sample of water was analysed using the Winkler method, a total of 0.60 mol of thiosulfate was used in the reaction.

How many moles of oxygen were present in the original sample?

- a) 0.15
- b) 0.30
- c) 0.60
- d) 1.20

14. The sodium salt of propanoic acid (sodium propanoate) is used as a preservative in bread and other baked goods. It can be produced by reacting propanoic acid with sodium hydroxide. In a particular experiment 100 mL of 0.080 M NaOH was added to 100 mL of 0.16 M propanoic acid.  
Which of the following statements is/are correct?
- I The pH of the resulting solution will be less than that of the propanoic acid solution.
  - II The resulting solution contains equal numbers of moles of propanoic acid and its conjugate base.
  - III Before the NaOH was added there were no propanoate ions present.
- a) II only
  - b) III only
  - c) I and II only
  - d) I and III only
15. In a flask, 10.0 mL of a 0.100 M HCl solution is diluted to 1.00 L. In a second flask, 10.0 mL of a 0.100 M KOH solution is also diluted to 1.00 L.  
Which statement best describes the changes in pH in these flasks?



pH change of the HCl solution	pH change of the KOH solution
-------------------------------	-------------------------------

- |                   |                |
|-------------------|----------------|
| a) increases by 2 | decreases by 2 |
| b) increases by 2 | increases by 2 |
| c) decreases by 2 | increases by 2 |
| d) decreases by 2 | decreases by 2 |
16. Citric acid, the predominant acid in lemon juice, is a triprotic acid. A student titrated 25.0 mL samples of lemon juice with 0.550 mol L<sup>-1</sup> NaOH. The mean titration volume was 29.50 mL. The molar mass of citric acid is 192.12 g mol<sup>-1</sup>.

What was the concentration of citric acid in the lemon juice?

- a) 1.04 g L<sup>-1</sup>
  - b) 41.6 g L<sup>-1</sup>
  - c) 125 g L<sup>-1</sup>
  - d) 374 g L<sup>-1</sup>
17. Which of the following pairs would form a buffer solution?
- a) HCl(aq)/Cl<sup>-</sup>(aq)
  - b) H<sub>2</sub>PO<sub>4</sub><sup>-</sup>(aq)/PO<sub>4</sub><sup>3-</sup>(aq)
  - c) H<sub>2</sub>SO<sub>4</sub>(aq)/HSO<sub>4</sub><sup>-</sup>(aq)
  - d) CH<sub>3</sub>COOH(aq)/CH<sub>3</sub>COO<sup>-</sup>(aq)

18. The following two unbalanced equations represent processes which are part of the nitrogen cycle.




Which one of the following alternatives correctly describes the reactants in each of these processes?

In process I, $\text{NH}_3(\text{aq})$ is	In process II, the $\text{NH}_4^+(\text{aq})$ ion is
a) an acid	reduced
b) a base	reduced
c) an acid	oxidised
d) a base	oxidised

**Questions 19 and 20 refer to the following information.**

A rechargeable cell, used in laptop computers, contains a metal alloy (designated M) which has hydrogen atoms absorbed on its surface, and nickel in the form of  $\text{NiO}(\text{OH})$  (s) and  $\text{Ni}(\text{OH})_2$  (s).

The half reactions, written as reduction reactions, are




While this cell is generating electricity, the metal alloy acts as the anode.

19. When this cell is generating electricity
- $\text{NiO}(\text{OH})$  acts as the oxidant.
  - the concentration of  $\text{OH}^-$  ions in the cell increases as the cell discharges.
  - $\text{OH}^-$  ions produced at the negative electrode migrate to the positive electrode.
  - electrons flow in the external circuit from the positive to the negative electrode.
20. When the cell is recharged, which one of the following processes occurs at the electrode connected to the positive terminal of the external power source?
- reduction of  $\text{H}_2\text{O}(\text{l})$
  - reduction of  $\text{NiO}(\text{OH})(\text{s})$
  - oxidation of  $\text{Ni}(\text{OH})_2(\text{s})$
  - oxidation of H (adsorbed on M)




21. Many reactions occurring in plant and animal cells involve a chemical called nicotinamide adenine dinucleotide,  $\text{NAD}^+$ . One such reaction is



It has been suggested that this reaction could be used in biochemical fuel cells to power pacemakers used to control irregular heartbeats.

If this reaction were performed in a fuel cell, NADH would


- undergo oxidation at the anode.
  - undergo reduction at the cathode.
  - undergo reduction at the anode.
  - undergo oxidation at the cathode.
22. An electrochemical cell is set up as shown in the diagram.



What are two observations for this electrochemical cell?

	<i>Observation 1</i>	<i>Observation 2</i>
a)	A reading was shown on the voltmeter	In Beaker 2 the solution became a darker blue
b)	In Beaker 2 the blue solution faded	A reddish precipitate formed on the copper electrode
c)	A grey precipitate formed on the aluminium electrode	In Beaker 2 the solution became a darker blue
d)	Electrons moved through the voltmeter	A reddish precipitate formed on the copper electrode

23. Iron pipes are used to transport natural gas to cities. Corrosion occurs when water droplets sit on the outer surface of the iron pipe. Miniature galvanic cells are created, with regions such as those below, that act as anodes and cathodes.




The type of region and reaction occurring at X in the cell is

Region	Reaction
a) anode	$\text{Fe(s)} \rightarrow \text{Fe}^{2+}(\text{aq}) + 2\text{e}^-$
b) cathode	$\text{Fe(s)} \rightarrow \text{Fe}^{2+}(\text{aq}) + 2\text{e}^-$
c) anode	$\text{O}_2(\text{g}) + 2\text{H}_2\text{O(l)} + 4\text{e}^- \rightarrow 4\text{OH}^-(\text{aq})$
d) cathode	$\text{O}_2(\text{g}) + 2\text{H}_2\text{O(l)} + 4\text{e}^- \rightarrow 4\text{OH}^-(\text{aq})$

24. In which of the following alternatives are the three compounds listed in order of increasing boiling point?
- Pentane, butan-1-ol, propanoic acid
  - Propanoic acid, butan-1-ol, pentane
  - Propanoic acid, pentane, butan-1-ol
  - Butan-1-ol, propanoic acid, pentane

25. Terylene (polyester) is a condensation polymer. Part of the structure of the polymer is shown.



What are the two monomers that form this polymer?

	Monomer 1	Monomer 2
a)		
b)		
c)		
d)		

## SECTION TWO – SHORT RESPONSE

[69 marks]

Section Two contains **THIRTEEN (13)** questions. Attempt **ALL** questions in the spaces provided.

In this section, unless asked to write molecular equations, chemical equations should refer to those species consumed in the reaction and the new species produced.


These species may be **ions** [for example  $\text{Ag}^+(\text{aq})$ ], **molecules** [for example  $\text{NH}_3(\text{g})$ ,  $\text{NH}_3(\text{aq})$ ,  $\text{CH}_3\text{COOH}(\text{l})$ ,  $\text{CH}_3\text{COOH}(\text{aq})$ ] or **solids** [for example  $\text{BaSO}_4(\text{s})$ ,  $\text{Cu}(\text{s})$ ,  $\text{Na}_2\text{SO}_4(\text{s})$ ].

Suggested working time: 60 minutes

---

### Question 1 [4 marks]

Mayonnaise is an example of an oil-in-water emulsion. It is stabilised by the addition of egg yolk, which contains the emulsifier (wetting agent) lecithin (structure below).




In terms of its structure, explain why the lecithin molecule is able to act as an emulsifier.

(4 marks)

**Question 2 [5 marks]**

The diagram shows the wreck of an iron ship sitting on the bottom of the ocean at a great depth.



- a) Identify the gas dissolved in water that causes corrosion.

(1 mark)

- b) Outline the effect of temperature and pressure on the solubility of gases in water.

(2 marks)

- c) Microenvironments are localised areas where conditions are different from those in the surrounding areas.


Explain the effect of **each** microenvironment on the rate of corrosion of the iron ship. Include at least one balanced chemical equation in your answer.

(2 marks)

**Question 3 [2 marks]**

A fuel cell that can provide power for buses is the phosphoric acid fuel cell, PAFC. The electrolyte is concentrated phosphoric acid and the reactants are hydrogen and oxygen gases.

A simplified sketch of a phosphoric acid fuel cell is given below.



Give the equation for the half cell reaction that takes place at the

i. anode of this cell


(1 mark)

ii. cathode of this cell


(1 mark)

**Question 4 [3 marks]**

A rechargeable galvanic cell, based on nickel and cadmium (NiCd cell), has been commercially available for a number of years and has been used to power small appliances such as mobile phones. A simplified diagram of a NiCd cell is given below.



The overall cell reaction for the cell when discharging is



- a) Identify the cathode and the anode electrodes by writing their name in the circles provided in the diagram.

(1 mark)

- b) What features of this secondary cell enables it to be recharged?


(1 mark)

- c) Give the equation for the half reaction that takes place at the anode when the cell is discharging.

(1 mark)

**Question 5 [5 marks]**

The diagram represents three separate petri dishes each containing a mixture of agar, sodium chloride solution, phenolphthalein and an indicator which turns blue in the presence of  $\text{Fe}^{2+}$ . Nails are added to each dish.



a) Why does the mixture contain sodium chloride solution?

(1 mark)

b) Write two half equations for the reactions occurring in Dish B.

(2 marks)

c) In which dish would the nail be protected from corrosion? Explain your answer.

(2 marks)



**Question 6 [4 marks]**

- a) Calculate the pH of a  $0.2 \text{ mol L}^{-1}$  solution of hydrochloric acid.

(1 mark)

- b) Calculate the pH after 20mL of  $0.01 \text{ mol L}^{-1}$  sodium hydroxide is added to 50 mL of  $0.2 \text{ mol L}^{-1}$  hydrochloric acid. Include a balanced chemical equation in your answer.

(3 marks)

**Question 7 [5 marks]**

**Red cabbage indicator chart**

Colour	red		violet		purple		blue		green		yellow			
pH	1	2	3	4	5	6	7	8	9	10	11	12	13	14

- a) State what colour the red cabbage indicator would be in a  $0.005 \text{ mol L}^{-1}$  solution of  $\text{H}_2\text{SO}_4$ . Show your working.

(1 mark)

- b) Using the red cabbage indicator, what colour would the solution be if 10 mL of  $0.005 \text{ mol L}^{-1} \text{ H}_2\text{SO}_4$  was diluted to 100 mL?


(1 mark)

- c) What volume of  $0.005 \text{ mol L}^{-1} \text{ KOH}$  is required to neutralise 15 mL of the diluted solution of  $\text{H}_2\text{SO}_4$ ?

(3 marks)

**Question 8 [3 marks]**

A 'QwikCure' pack, used to treat sporting injuries, contains a bag of water inside a larger bag of finely powdered ammonium nitrate,  $\text{NH}_4\text{NO}_3$ . Squeezing the pack causes the bag of water to break and the  $\text{NH}_4\text{NO}_3$  to dissolve. The change of energy that occurs can be used to treat an injury.




- a) Suppose the activation energy of the **reverse reaction** is  $35 \text{ kJ mol}^{-1}$ .

- i. Explain the meaning of the term 'activation energy'.

(1 mark)

- ii. On the graph below, sketch an energy profile diagram showing the changes that occur in chemical energy as the  $\text{NH}_4\text{NO}_3$  powder dissolves.



Answer hidden

**Question 9** [9 marks]

Complete the table below.

For the dot diagrams, show all valence electron pairs as : or –

For the shapes, either a name or a diagram should be given.

For the intermolecular force, only the principal force operating in a pure sample of the substance in the solid state should be given.

Substance	Dot diagram	Shape	Intermolecular force
Propanone $\text{C}_3\text{H}_6\text{O}$			
Monochloroacetic acid $\text{C}_2\text{H}_3\text{ClO}_2$			
Dibromoethyne $\text{C}_2\text{Br}_2$			

### Question 10 [6 marks]

The discovery of element 113 was claimed by teams of Russian and American scientists in February 2004. Following international conventions, it has initially been given the name ununtrium and the symbol Uut, before a permanent name and symbol are given to it.

Uut undergoes rapid radioactive decay but atoms of Uut have been identified with a mass number of 283 and also with a mass number of 284.

- a) State the number of subatomic particles in an uncharged Uut atom of mass number 284.

Protons	Electrons	Neutrons	
			(1 mark)

- b) In what group and period is Uut located in the periodic table?

Group	Period	
		(2 marks)

- c) Give the symbol of the element that is expected to be most similar to Uut in chemical properties.

(1 mark)

- d) In terms of atomic structure, explain why the atomic radius of Uut is predicted to be smaller than that of Fr ( $Z = 87$ ).

(1 mark)

- e) In terms of atomic structure, explain why the first ionisation energy of Uut is predicted to be smaller than that of Al ( $Z = 13$ ).

(1 mark)

**Question 11 [9 marks]**

Write equations for any reactions that occur in the following procedures. If no reaction occurs write no reaction. In each case describe in full what you would observe, including any colours, odours, precipitates (give the colour), gases evolved (give the colour or describe as colourless). If no change is observed, you should state this.

Barium hydroxide solution is added to a solution of hydrochloric acid.

Equation:

Observation:

(3 marks)

A piece of copper metal is added to concentrated nitric acid.

Equation:

Observation:

(3 marks)

Ethanol is warmed with acidified acetic acid solution.

Equation:

Observation:

(3 marks)

**Question 12 [6 marks]**

The following table lists the pH of 0.10 M solutions of four different acids at 25°C.

Acid	pH
I	1.0
II	3.0
III	0.7
IV	2.1

- a) Which acid **must** have more than one acidic hydrogen per molecule? Give a reason for your answer.

(2 marks)

- b) Calculate the value of the  $[\text{OH}^-]$  present in the solution of acid III.

(2 marks)

- c) Samples of the solutions of acids I and IV are diluted by a factor of 10. The resulting **change in pH** units would be  
(Tick **one** of the following boxes.)

greater for acid i than for acid IV	
greater for acid iv than for acid I	
the same for both acids	

Give an explanation for your answer.

(2 marks)


**Question 13 [8 marks]**

The polymer, polyvinylchloride is composed of units of the monomer  $\text{C}_2\text{H}_3\text{Cl}$ .

a) Draw the structure of this monomer. (1 mark)

b) Draw a section of the polymer showing at least three repeating units. (1 mark)

A tripeptide may be represented by the structure shown below.



c) Draw structures to show the amino acids from which this polymer is formed. (2 marks)

d) Draw a section, with at least three silicon atoms, of the silicone that could be prepared from the silanol shown below. (1 mark)



Silicones have a useful range of physical properties including:

- Stable at extreme temperatures
- Water repellent
- Flexible

Choose one of these properties and explain it.

(3 marks)

### SECTION THREE – EXTENDED RESPONSE

[65 marks]

Section Three has **SIX (6)** questions. Attempt ALL questions in the spaces provided below.


In descriptive responses, marks are awarded for relevant chemical content, including equations, diagrams and illustrative examples of the chemistry you are describing.

Calculations are to be set out in detail. Marks will be awarded for correct equations and clear setting out, even if you cannot complete the calculation. Express numerical answers to three (3) significant figures and provide units where appropriate.

Suggested working time: 70 minutes

#### Question 1 [12 marks]

A particular industrial process involves the following steps.





- a) It is possible to alter the temperature and pressure at which reaction 2 occurs. In the table below, indicate what effect the following changes to temperature and pressure would have on the **rate**, **equilibrium yield** and value of the **equilibrium constant, *K***, for reaction 2.

	Would the <b>rate</b> of reaction 2 become <b>higher</b> , <b>lower</b> or <b>remain unchanged</b> ?	Would the <b>equilibrium</b> yield of reaction 2 become <b>higher</b> , <b>lower</b> or <b>remain unchanged</b> ?	Would the <b>value of the equilibrium constant, <i>K</i></b> , of reaction 2 become <b>higher</b> , <b>lower</b> or <b>remain unchanged</b> ?
The temperature of reaction 2 is lowered to 150°C			
The pressure of reaction 2 is increased to 5 atm by pumping more B(g) and C(g) into the reaction vessel, at constant temperature.			

(6 marks)

- b) The graph below shows the variation in concentration of reactant and products as a function of time for the following system.



Identify and explain each of the changes in conditions that have shaped the curves during the time the system was observed.

(6 marks)

## Question 2 [10 marks]

When assessing the alcohol content of wine for labelling purposes, the ethanol may be separated by distillation of the wine and then treated with excess acidified potassium permanganate solution. The reaction mixture is placed in a water bath for several minutes to ensure the complete oxidation of the alcohol to acetic acid.

Iron II sulfate solution is then added to reduce the excess permanganate ion. By this method, the amount of permanganate required to oxidise the alcohol and hence the amount of alcohol, may be determined.


The following results were obtained for a 750 mL bottle of wine, analysed as described above.

Mass of 750 mL wine = 750 g

Vol of 1.500 mol L<sup>-1</sup> KMnO<sub>4</sub> added to the distillate = 1250 mL

Vol of 0.5000 mol L<sup>-1</sup> Fe<sup>2+</sup> solution required to titrate the excess KMnO<sub>4</sub> = 447.5 mL

The half-equation for the oxidation of ethanol is:




- a) Calculate the number of moles of KMnO<sub>4</sub> added to the ethanol. (1mark)
  
  
  
  
  
  
  
  
  
  
- b) Calculate the number of moles of KMnO<sub>4</sub> in excess. (3 marks)
  
  
  
  
  
  
  
  
  
  
- c) Calculate the number of moles of KMnO<sub>4</sub> that combined with the ethanol. (2 marks)
  
  
  
  
  
  
  
  
  
  
- d) Calculate the concentration of ethanol in the wine as a percentage by mass. (4 marks)


### Question 3 [12 marks]

A student carried out a first-hand investigation to identify the relationship between the mass of a metal used in a reaction, and the volume of gas produced. The first-hand investigation was carried out at 25°C, and 100 kPa. In each experiment, 200 mL of hydrochloric acid was added to some zinc, and the volume of gas produced was recorded. The diagram shows the equipment used, and the table contains the student's results.

<i>Mass of zinc (g)</i>	<i>Volume of gas collected at 25°C and 100 kPa (mL)</i>
0.12	45
0.33	125
0.56	115
0.83	315
0.96	365
1.22	380
1.64	380
1.93	380



a) Graph the results from these experiments.



(3 marks)

- b) Predict the volume of gas that would be produced in this experiment if 3.00g of zinc had been used. Justify your answer.

(2 marks)

- c) Calculate the theoretical volume of gas produced at 25°C and 100 kPa by the reaction of 0.56 g of zinc with the 200 mL of hydrochloric acid.

(2 marks)

- d) A different student carried out the same experiment. She used a constant temperature of 15°C.  
Describe the results she would have obtained and explain why these results would vary from those shown above.


(5 marks)

#### Question 4 [10 marks]

The compound known as phenazine ( $C_{12}H_8N_2$ ) is important in the manufacture of a number of dyes that are used as biological staining agents in the science of histology. Phenazine is manufactured in a series of stages as shown by the equations below. The efficiencies of the stages are shown in brackets.

Stage 1 (87% efficient)

Concentrated sulphuric and nitric acids are heated together.




Stage 2 (40% efficient)

The hot acid mixture is boiled with benzene to form nitrobenzene.



Stage 3 (62% efficient)

The nitrobenzene is reacted with aminobenzene to form phenazine.



(a) If 120 tonnes ( $1.20 \times 10^8$  g) of nitrobenzene are consumed in the manufacturing process in one week of production, calculate the mass of phenazine formed. (3 marks)

(b) Calculate the volume of 8M nitric acid used in Stage 1 in the same week.

(4 marks)

(c) Draw the structures of nitrobenzene and aminobenzene and suggest a structure for phenazine.

(3 marks)



**Question 5 [9 marks]**

Analysis of a block of air freshener indicated that it contained several compounds that could be separated from each other by dissolving in a number of solvents. One of the ingredients, which dissolved in water, was found to contain only carbon, hydrogen oxygen and nitrogen.

A 1.255 g sample of the compound was burned in air and produced 1.171 g of carbon dioxide and 1.202 g of water.

A second 2.552 g sample was decomposed at high temperature and found to contain 0.7602 g of nitrogen.

A third sample, weighing 1.500 g was vapourised at 150°C and 101.3 kPa and the vapour occupied 1.108 L.

A fourth sample was dissolved water and tested with phenolphthalein. The indicator turned pink. The compound was also found to turn acidified potassium dichromate solution from orange to green.

(a) Determine the empirical formula of the compound. (4 marks)

(b) Determine the molecular formula of the compound.

(2 marks)

(c) Suggest a structure for the compound.

(1 mark)

(d) Write an equation to explain why the compound turned phenolphthalein pink.

(2 marks)

**Question 6****[12 marks]**

Consider the information presented in the table below.

Substance	Electrical conductivity of solid	Solubility in water	Solubility in petrol	Melting point (°C)
Candle wax $C_{25}H_{52}$	nil	insoluble	soluble	47
Methanol	nil	soluble	insoluble	-97
Sodium bromide	nil	soluble	insoluble	734
Lead iodide	nil	insoluble	insoluble	402
Propyl propanoate	nil	insoluble	soluble	-76

Discuss the similarities and differences in physical properties that are presented by these data.