



PERTH MODERN SCHOOL

Exceptional schooling. Exceptional students.

INDEPENDENT PUBLIC SCHOOL

Semester One Examination, 2020

Question/Answer booklet

MATHEMATICS METHODS UNIT 1

Section Two: Calculator-assumed

Your Name _____

Your Teacher's Name _____

Time allowed for this section

Reading time before commencing work: ten minutes
Working time: one hundred minutes

Materials required/recommended for this section

To be provided by the supervisor

This Question/Answer booklet
Formula sheet (retained from Section One)

To be provided by the candidate

Standard items: pens (blue/black preferred), pencils (including coloured), sharpener, correction fluid/tape, eraser, ruler, highlighters

Special items: drawing instruments, templates, notes on two unfolded sheets of A4 paper, and up to three calculators approved for use in this examination

Important note to candidates

No other items may be taken into the examination room. It is **your** responsibility to ensure that you do not have any unauthorised material. If you have any unauthorised material with you, hand it to the supervisor **before** reading any further.

Question	Marks	Max	Question	Marks	Max
9		9	17		7
10		8	18		11
11		5	19		6
12		8	20		8
13		7	21		6
14		10			
15		5			
16		10			

Structure of this paper

Section	Number of questions available	Number of questions to be answered	Working time (minutes)	Marks available	Percentage of examination
Section One: Calculator-free	8	8	50	50	33
Section Two: Calculator-assumed	13	13	100	100	67
Total					100

Instructions to candidates

1. The rules for the conduct of the Western Australian Certificate of Education ATAR course examinations are detailed in the *Year 12 Information Handbook 2020*. Sitting this examination implies that you agree to abide by these rules.
2. Write your answers in this Question/Answer booklet.
3. You must be careful to confine your answers to the specific questions asked and to follow any instructions that are specific to a particular question.
4. Additional pages for the use of planning your answer to a question or continuing your answer to a question have been provided at the end of this Question/Answer booklet. If you use the space to continue an answer, indicate in the original answer space where the answer is continued, i.e. give the page number.
5. **Show all your working clearly.** Your working should be in sufficient detail to allow your answers to be checked readily and for marks to be awarded for reasoning. Incorrect answers given without supporting reasoning cannot be allocated any marks. For any question or part question worth more than two marks, valid working or justification is required to receive full marks. If you repeat any question, ensure that you cancel the answer you do not wish to have marked.
6. It is recommended that you **do not use pencil**, except in diagrams.
7. The Formula sheet is **not** to be handed in with your Question/Answer booklet.

Section Two: Calculator-assumed

(100 Marks)

This section has **thirteen (13)** questions. Answer **all** questions. Write your answers in the spaces provided.

Spare pages are included at the end of this booklet. They can be used for planning your responses and/or as additional space if required to continue an answer.

- Planning: If you use the spare pages for planning, indicate this clearly at the top of the page.
- Continuing an answer: If you need to use the space to continue an answer, indicate in the original answer space where the answer is continued, i.e. give the page number. Fill in the number of the question that you are continuing to answer at the top of the page.

Working time: 100 minutes.


Question 9 {1.1.23, 1.1.24, 1.1.25}

(9 marks)


(a) Circle the diagrams that show functions.

(2 marks)


(i)




(ii)



(iii)



(iv)




Solution
(i)
(ii)
(iii)
Specific behaviours
✓ circles two functions
✓ circles three functions

(b) State the domain and range for the following functions.


(4 marks)

i)



Solution
Domain: $\{x \in R: -3 \leq x < 3\}$
Range: $\{y \in R: -5 \leq y \leq 5\}$
Specific behaviours
✓ correct domain ✓ correct range

ii)



Solution
Domain: $\{x \in R: x \neq 2\}$
Range: $\{y \in R: y \neq -3\}$
Specific behaviours
✓ correct domain ✓ correct range

(c) If $f(x) = x^2 + 2x$ find

(3 marks)

i) $f(-1) = -1$

ii) $f(a) = a^2 + 2a$

iii) $f(x+h) = (x+h)^2 + 2(x+h)$

Solution
$f(-1) = (-1)^2 + 2 \times (-1) = -1$ $f(a) = a^2 + 2a$ $f(x+h) = (x+h)^2 + 2(x+h)$
Specific behaviours
✓ correct value for $f(-1)$ ✓ correct expression for $f(a)$ ✓ correct expression for $f(x+h)$

Question 10 {1.3.13, 1.3.15, 1.3.16}

(8 marks)

In a school survey of 207 students in Year 11 and Year 12, it was observed that 77 of the 102 Year 12 students studied Mathematics and the 10 students in Year 11 did not study Mathematics.

If one student is selected at random from those surveyed, determine the probability that

(a) they were in Year 11.

(1 mark)

Solution
$P(\text{Year 11}) = \frac{207 - 102}{207} = \frac{105}{207}$
Specific behaviours
✓ correct probability

(b) they studied Mathematics.

(1 mark)

Solution
$P(\text{Maths}) = \frac{77 + (105 - 10)}{207} = \frac{172}{207}$
Specific behaviours
✓ correct probability

(c) they were in Year 12 or studied Mathematics.

(2 marks)

Solution
$P(\text{Year 12} \cup \text{Maths}) = \frac{102 + 172 - 77}{207} = \frac{197}{207}$
Specific behaviours
✓ uses inclusion-exclusion principle ✓ correct probability

(d) They studied Mathematics given they were in Year 11.

(2 marks)

Solution
$P(\text{Maths} \text{Year 11}) = \frac{95}{105}$
Specific behaviours
✓ correct numerator ✓ correct denominator

(e) Without calculating any further probabilities, is there any indication that studying Mathematics is independent of Year? Justify your answer.

(2 marks)


Solution
The choice of Mathematics is NOT independent of year. As $P(\text{Maths}) = \frac{172}{207} \approx 0.83$ but $P(\text{Maths} \cap \text{Year 11}) = \frac{95}{105} \approx 0.90$
Specific behaviours

- ✓ states 'not independent'
- ✓ uses the rule $P(A|B) = P(A)$

Question 11 {1.1.3, 1.1.4, 1.1.5}

(5 marks)

Consider the graph shown:



(a) Find the gradient of the line PQ.

(2 marks)

Solution
$m_{PQ} = \frac{3a - 0}{-5a - a} = \frac{3a}{-6a} = -\frac{1}{2}$
Specific behaviours
<ul style="list-style-type: none"> ✓ uses $\frac{\text{rise}}{\text{run}}$ ✓ correct gradient

(b) Find the equation of the line RS.
marks)

(2

Solution
$m_{RS} \times m_{PQ} = -1$ $\frac{-1}{2} m_{RS} = -1 \Rightarrow m_{RS} = 2$ $y = 2x + 6b$
Specific behaviours
<ul style="list-style-type: none"> ✓ determines gradient of line RS ✓ determines equation of line RS

(c) Find the coordinates of point S.
mark)

(1

Solution
$S(0, 6b)$
Specific behaviours
<ul style="list-style-type: none"> ✓ states the coordinates of S

Question 12 {1.1.7, 1.1.10, 1.1.12}

(8 marks)

Use your ClassPad to display parabolas $f(x)$, $g(x)$ and $h(x)$ whose equations are

$$f(x) = -0.5x(x+2)$$

$$g(x) = -2x(x-4)$$

$$h(x) = x(x-1)$$

(a) Write down a feature these three parabolas have in common.

(1 mark)

Solution
<i>They all pass through (0,0).</i>
Specific behaviours
✓ states parabolas pass through the origin

(b) Write down the turning point for each parabola and state its nature.

(3 marks)

Solution
For $f(x)$, maximum turning point $(-1, 0.5)$ For $g(x)$, maximum turning point $(2, 8)$ For $h(x)$, minimum turning point $(0.5, -0.25)$
Specific behaviours
✓ states maximum turning point and its coordinates for $f(x)$ ✓ states maximum turning point and its coordinates for $g(x)$ ✓ states minimum turning point and its coordinates for $h(x)$

(c) Write down the equation of $h(x)$ in turning point form and state the line of symmetry.

(2 marks)

Solution
$h(x) = (x - 0.5)^2 - 0.25$ $x = 0.5$
Specific behaviours
✓ expresses $h(x)$ in turning point form ✓ states equation for the line of symmetry

(d) If the graph of $h(x)$ is translated 2 units in the positive direction of the x axis and 2 units in the negative direction of the y axis, write down the new equation for $h(x)$ and state its new line of symmetry.

(2 marks)

Solution
$h(x-2) - 2 = [(x-2) - 0.5]^2 - 0.25 - 2$ $x = 2.5$
Specific behaviours
✓ writes equation for $h(x)$ after the translations ✓ states equation for the line of symmetry after the

translations


Question 13 {1.1.15}

(7 marks)

Sketch the graphs of the following functions. Draw and label the key features, including x- and y-intercepts, vertices, asymptotes (if relevant), showing correct general shape and behaviour.

(a) $y = \frac{4}{x} - 5$


(4 marks)



(b) $y = 3\sqrt{x}$

(3 marks)

Solution



Specific behaviours

- ✓ states at (0, 0)
- ✓ plots and labels at least one point on the graph (4, 6), (16, 12)
- ✓ correct shape with smooth curve

Question 14 {1.1.18}


(10 marks)

(a) Sketch the graph of each of the following function, labelling intercepts, point of inflection and showing the shape and behaviour as $x \rightarrow \infty$ and $x \rightarrow -\infty$.

i) $y = -x^3$

(2 marks)

Solution




Specific behaviours

- ✓ plots and label the point of inflection (0, 0)
- ✓ current shape with smooth curve

ii) $y = 2(3-x)^3 + 4$

(3 marks)


Solution



- (b) Express cubic function $y = \frac{-x^3}{2} + 2x^2 + 2x - 8$ in its factorised form and sketch its graph, labelling all intercepts and using arrows to show its shape and general behaviour as $x \rightarrow -\infty$ and $x \rightarrow \infty$.

(5 marks)

Solution
$y = \frac{-1}{2}(x+2)(x-2)(x-4)$

**Question 15 {1.1.13}****(5 marks)**

An online company that sells onesies has modelled their expected number of daily sales (S) against the average daily temperature (T) in $^{\circ}\text{C}$.

Their model has S as inversely proportional to $\sqrt{2T+16}$ for $-5 \leq T \leq 30$.

On a 10°C day in August they sold 20 onesies.

a) Determine the relationship between S and T.

(3 marks)

Solution
$S \propto \frac{1}{\sqrt{2T+16}} \Rightarrow S = \frac{k}{\sqrt{2T+16}}$ $20 = \frac{k}{\sqrt{2 \times 10 + 16}} \Rightarrow k = 120$ $S = \frac{120}{\sqrt{2T+16}}$
Specific behaviours
<ul style="list-style-type: none"> ✓ constructs equation for S ✓ determines k ✓ determines relationship between S and T

- b) Determine the **number of expected sales** tomorrow if the average forecasted temperature is 29. (2 marks)

Solution
$S = \frac{120}{\sqrt{2 \times 29 + 16}} = 13.9 (1 d. p)$ <p>Hence, the number of expected sales is 14</p>
Specific behaviours
<ul style="list-style-type: none"> ✓ substitutes $T = 29$ ✓ states the number of expected sales as an integer


Question 16 {1.1.14}

(10 marks)

- a) State all the asymptotes on each graph below then determine each equation of the rectangular hyperbola:

i)

(3 marks)



Solution

$$y = \frac{k}{x+1}$$

$$\frac{k}{0+1} = 3 \Rightarrow k = 3$$


$$y = \frac{3}{x+1}$$

Specific behaviours

- ✓ states asymptotes $x = -1$ & $y = 0$
- ✓ substitutes point and solves
- ✓ states equation

ii)

(3 marks)



Solution

$$y = \frac{k}{x-3} + 2$$

$$\frac{k}{0-3} + 2 = 4 \Rightarrow k = -6$$

$$y = \frac{-6}{x-3} + 2$$

Specific behaviours

- ✓ states asymptotes $x = 3$ & $y = 2$
- ✓ substitutes point and solves
- ✓ states equation

- b) Complete the table below.

c) (4 marks)

Solution				
Curve	x-intercept	y-intercept	Asymptote parallel to the	
			x-axis	y-axis
$y = \frac{6}{x-2} - 3$	(4, 0)	(0, -6)	$y = -3$	$x = 2$
$y = \frac{-9}{x+3}$	None	(0, -3)	$y = 0$	$x = -3$
Specific behaviours				
<ul style="list-style-type: none"> ✓ correct x-intercepts ✓ correct y-intercepts 				

- ✓ correct horizontal asymptotes
- ✓ correct vertical asymptotes

(7 marks)


A C-9 jet utilised by NASA enters a parabolic flight at an altitude of 9200m to simulate zero gravity. For the duration of the zero-gravity flight, its altitude A in metres can be represented as a function of time t in seconds as follows:

$$A(t)=at^2+bt+c$$

- a) Given that the function passes through the points $(2, 9400)$ and $(4, 9560)$, determine the values of a , b and c . (3 marks)

Solution

$A(0)=9200 \Rightarrow c=9200$



$$\begin{cases} 4a+2b+9200=9400 \\ 16a+4b+9200=9560 \end{cases} | a, b$$
$$\{a=-5, b=110\}$$

Specific behaviours

- ✓ states value for c
- ✓ states value for a
- ✓ states value for b

- b) When is the maximum altitude reached by the jet? State the maximum altitude. (2 marks)

Solution
$A(t) = -5t^2 + 110t + 9200$ $\text{When } t = \frac{-110}{2 \times (-5)} = 11 \text{ s}$ $A(11) = -5(11)^2 + 110(11) + 9200$ $= 9805$ <p>Hence, the maximum altitude is 9805m at 11s</p>
Specific behaviours
<ul style="list-style-type: none"> ✓ correct value for t at maximum altitude ✓ correct value for A at maximum altitude

- c) Given that the plane exits the parabolic flight at the same altitude it begins at, what is the duration of the zero-gravity simulation? (2 marks)

Solution
$A(t) = -5x^2 + 110x + 9200 = 9200$ $t(t - 22) = 0$ $\therefore t = 0 \vee 22$ <p>Hence, the duration is 22s</p>
Specific behaviours

- ✓ substitutes $A(t)$ by 9200 and solves for t
- ✓ states the duration

Question 18 {1.1.26, 1.1.27}

(11 marks)

(a) Let $f(x) = x^3$

- (i) What is the equation of the new function if $f(x)$ is translated 4 units to the right and 3 units down? What are the coordinates of the point of inflection?

(3 marks)

Solution
$y = (x - 4)^3 - 3$ <p>Point of inflection is $(4, -3)$</p>
Specific behaviours
<ul style="list-style-type: none"> ✓ writes equation for $f(x - 4) - 3$ ✓ states point of inflection for $f(x - 4) - 3$

- (ii) Find the coordinates of the point of inflection of the function $y = \frac{-1}{2}f(2x + 1)$.

(2 marks)

Solution
$\frac{-1}{2}f(2x + 1) = \frac{-1}{2}f\left[2\left(x + \frac{1}{2}\right)\right]$ <p>Since the point of inflection for $f(x)$ is $(0, 0)$</p> <p>$\left(\frac{-1}{2}, 0\right)$ is the point of inflection for $\frac{-1}{2}f(2x + 1)$.</p>
Specific behaviours
<ul style="list-style-type: none"> ✓ correct x value for the point of inflection for $\frac{-1}{2}f(2x + 1)$ ✓ correct y value for the point of inflection for $\frac{-1}{2}f(2x + 1)$.

- (b) State the sequence of graphical transformation(s) which occurs if:

- (i) $g(x) \rightarrow g(2x - 2)$

(2 marks)

Solution	
<p>If in factorised form $g\left[2\left(x - 1\right)\right]$:</p> <p>Horizontal dilation by scale factor $\frac{1}{2}$, then horizontal translation to the right by 1.</p>	<p>If in expanded form $g(2x - 2)$:</p> <p>Horizontal translation to the right by 2, then Horizontal dilation by scale factor $\frac{1}{2}$.</p>
Specific behaviours	
<ul style="list-style-type: none"> ✓ states horizontal dilation in correct order ✓ states horizontal translation in correct order 	<p><i>Incorrect answer if student gives "horizontal dilation by SF $\frac{1}{2}$" prior to "translate left by 2".</i></p>

- (ii) $h(x) \rightarrow \frac{5}{2}h\left(\frac{2}{3}x + \frac{\pi}{3}\right) - 1$

(4 marks)


Solution	
<p>If in factorised form $\frac{5}{2}h\left[\frac{2}{3}\left(x + \frac{\pi}{2}\right)\right] - 1$:</p>	<p>If in expanded form $\frac{5}{2}h\left(\frac{2}{3}x + \frac{\pi}{3}\right) - 1$</p>

<p>①Vertical dilation by scale factor $\frac{5}{2}$</p> <p>②Horizontal dilation by scale factor $\frac{3}{2}$</p> <p>③Horizontal translation to the left by $\frac{\pi}{2}$</p> <p>④Vertical translation down by 1</p>	<p>①Vertical dilation by scale factor $\frac{5}{2}$</p> <p>②Horizontal translation to the left by $\frac{\pi}{3}$</p> <p>③Horizontal dilation by scale factor $\frac{3}{2}$</p> <p>④Vertical translation down by 1</p>
Specific behaviours	
<ul style="list-style-type: none"> ✓ states vertical dilation in correct order ✓ states horizontal dilation in correct order ✓ states horizontal translation in correct order ✓ states vertical translation in correct order 	<p>For both versions, ①&② can switch, ③&④ can switch. Incorrect answer if student gives "horizontal dilation by SF $\frac{3}{2}$" prior to "translate left by $\frac{\pi}{3}$".</p>

Question 19 {1.1.26, 1.1.27}


(6 marks)

The graph shows the function $y = f(x)$.




(a) Sketch the graph of $y = 2f(x+2)$.


(3 marks)



(b) Sketch the graph of $y = f(-x) - 2$.

(3 marks)





Question 20 {1.3.11, 1.3.12, 1.3.17}

(8 marks)

1000 people were studied to see if they contracted an illness (I) and whether they were vaccinated against it (V). 86 people caught the illness and 348 were vaccinated, including 1 person who caught the illness despite being vaccinated.

- a) Given that the above data is representative of a population, find the probability that a person randomly selected from the population:

- i) is not vaccinated?

(2 marks)

Solution
$P(\bar{V}) = 1 - P(V) = 1 - \frac{348}{1000}$ $= \frac{652}{1000}$
Specific behaviours
✓ uses the rule $P(\bar{A}) = 1 - P(A)$ ✓ correct probability

- ii) has caught the illness or is vaccinated?

(2 marks)

Solution
$P(I \cup V) = P(I) + P(V) - P(I \cap V)$ $= \frac{86}{1000} + \frac{348}{1000} - \frac{1}{1000}$ $= \frac{433}{1000}$
Specific behaviours
✓ uses the rule $P(A \cup B) = P(A) + P(B) - P(A \cap B)$ ✓ correct probability

- b) Does the vaccination affect a person's probability of catching the illness? If it does, how is it affected? Justify your answer with calculations. (4 marks)

Solution
$P(I V) = \frac{1}{348} \approx 0.003$ $P(I \bar{V}) = \frac{85}{652} \approx 0.13$ <p>Yes. It decreases the probability of catching the illness.</p>
Specific behaviours
<ul style="list-style-type: none"> ✓ states $P(I V)$ ✓ states $P(I \bar{V})$ ✓ states "Yes" ✓ compares $P(I V)$ and $P(I \bar{V})$

Question 21 {1.1.20}



(6 marks)

It is found that the shape of a part of a roller coaster ride can be modelled by a cubic function for $x \in [0, 15]$. The coordinates of several points on the track are (2, 360.5), (6, 195.14), (10, 54.74) and (12, 87.5).

- (a) Determine the rule for the function in the form $f(x) = ax^3 + bx^2 + cx + d$ for Find the values of a , b , c and d . (4 marks)

Solution
<p>Define $f(x) = a \cdot x^3 + b \cdot x^2 + c \cdot x + d$ done</p> <div style="border: 1px solid black; padding: 5px; display: inline-block;"> $\begin{cases} f(2) = 360.5 \\ f(6) = 195.14 \\ f(10) = 54.74 \\ f(12) = 87.5 \end{cases} \quad a, b, c, d$ $\left\{ a = \frac{39}{50}, b = -\frac{663}{50}, c = \frac{1209}{50}, d = \frac{17947}{50} \right\}$ </div> <p>OR $\{a = 0.78, b = -13.26, c = 24.18, d = 358.94\}$</p>
Specific behaviours
<ul style="list-style-type: none"> ✓ correct value for a ✓ correct value for b ✓ correct value for c ✓ correct value for d

- (b) Find the coordinates of the point(s) on the track that is furthest from the ground. (2 marks)

Solution	
	(1, 370.64)
	(15, 370.64)
Specific behaviours	
<ul style="list-style-type: none"> ✓ obtains one point and states its coordinates ✓ obtains two points and states their coordinates 	

END OF QUESTIONS

Additional working space

Question number: _____

Question number: _____

Additional working space

Question number: _____