Australian Islamic College 2019

ATAR Chemistry Units 3 and 4

Task 6 (Weighting: 3%)

Redox and Electrochemistry Test

Test Time: 50 minutes

Please do not turn this page until instructed to do so.

Surname		
Teacher		

Mark / 52	Percentage

Equipment allowed: Pens, pencils, erasers, whiteout, correction tape, rulers and non-programmable calculators permitted by the Schools Curriculum and Standards Authority.

Special conditions:

2 marks will be deducted for failing to write your full name on this test paper.

Teacher help: Your teacher can only help you during your test in one situation.

If you believe there is a mistake in a question show your teacher and your teacher will tell you whether or not there is a mistake in the question and if appropriate, how to fix that mistake.

Spelling of Science words must be correct. Science words with more than one letter wrong (wrong letter and/or wrong place) will be marked wrong.

Equations must be written balanced and with correct state symbols or they will be marked wrong.

Questions must be answered in this booklet.

Total marks: 52

PART ONE: MULTIPLE CHOICE QUESTIONS (10 MARKS)

- 1. Which one of the following species listed below contains sulfur with the lowest oxidation state?
 - (a) SO₃
 - (b) SO_2
 - (c) $(NH_4)_2S$
 - (d) H_2SO_4
- 2. In which one of the following processes is the oxidation number of chlorine being increased by 2?
 - (a) $PC\ell_3 + C\ell_2 \rightarrow PC\ell_5$
 - (b) $C\ell_2 + H_2O \rightarrow C\ell^- + HC\ellO + H^+$
 - (c) $2C\ell^- + 2e^- \rightarrow C\ell_2$
 - (d) $ClO_3^- + H_2O_2 \rightarrow ClO_4^- + H_2O$
- 3. Which one of the following substances is capable of oxidising copper metal?
 - (a) Zn
 - (b) Br_2
 - (c) CrC_{l₃}
 - (d) Au
- 4. Which one of the following pairs of chemicals will undergo a spontaneous redox reaction?
 - (a) solid iodine and water
 - (b) solid lead and liquid bromine
 - (c) lead nitrate solution and potassium iodide solution.
 - (d) solid copper and a 1.0 mol L⁻¹ solution of hydrochloric acid.
- 5. The following half-reactions occur in a zinc-carbon cell during discharge.

Anode $Zn_{(s)} \rightarrow Zn^{2+}_{(aq)} + 2e^{-}$


Cathode $2MnO_{2(s)} + 2NH_4^+_{(aq)} + 2e^- \rightarrow Mn_2O_{3(s)} + 2NH_{3(aq)} + H_2O_{(l)}$

What is the oxidising agent in a zinc-carbon cell?

- (a) Zn.
- (b) Mn.
- (c) MnO_2 .
- (d) NH_4^+ .

The following diagram relates to questions 6, 7 and 8.

The following galvanic cell was set up. Platinum (Pt) is an inert electrode.



- 6. When the circuit is connected, which one of the following concentrations will decrease?
 - (a) $[Fe^{3+}]$
 - (b) [Fe²⁺]
 - (c) $[Sn^{2+}]$
 - (d) $[K^{\dagger}]$
- 7. When the circuit is connected, which one of the following will be the changes of masses of the electrodes?

	Mass of tin electrode	Mass of platinum electrode
(a)	increases	decreases
(b)	increases	no change
(c)	decreases	no change
(d)	no change	increases

- 8. Which one of the following would be the expected reading on the voltmeter?
 - (a) 0.63 V
 - (b) 0.90 V
 - (c) 1.68 V
 - (d) 0.30 V

Question 9 and 10 relate to the following information

Most car batteries consist of six cells. The cell reactions involve lead, lead(IV) oxide and lead sulfate. Each cell produces just over 2 volts when fully charged. The car battery is a secondary battery.

The following reactions occur in a lead-acid battery as it discharges (produces electricity).

Anodes
$$Pb(s) + SO_4^{2-}(aq) \rightarrow PbSO_4(s) + 2e^{-}$$

Cathodes
$$PbO_2(s) + SO_4^{2-}(aq) + 4H^+(aq) + 2e^- \rightarrow PbSO_4(s) + 2H_2O(\ell)$$

- 9. The car battery is described as secondary because
 - (a) it requires two separate reactions.
 - (b) the cells can be recharged.
 - (c) it contains more than one cell.
 - (d) it requires the reactants to be supplied to the cell during operation.
- 10. Which one of the following is the overall equation for the discharging of the cell?
 - (a) $Pb(s) + PbO_2(s) + 2SO_4^{2-}(aq) \rightarrow 2PbSO_4(s) + 2H_2O(\ell)$
 - (b) $Pb(s) + PbO_2(s) + 2SO_4^{2-}(aq) + 4H^+(aq) \rightarrow PbSO_4(s) + 2H_2O(\ell)$
 - (c) $Pb(s) + PbO_2(s) + H_2SO_4(aq) \rightarrow 2PbSO_4(s) + 2H_2O(\ell)$
 - (d) Pb(s) + PbO₂(s) + 2SO₄²⁻(aq) + 4H⁺(aq) \rightarrow 2PbSO₄(s) + 2H₂O(ℓ)

END OF MULTIPLE CHOICE SECTION

(19 marks)

This section has **2** questions. Answer both questions. Write your answers in the spaces provided.

Question 11 (4 marks)

The two initial reactions involved in the wet corrosion of iron are shown below. Both of these reactions are reversible.

Oxidation (anode) reaction: Fe(s) \rightarrow Fe²⁺(aq) + 2e⁻ Reduction (cathode) reaction: O₂(g) + 2H₂O(ℓ) + 4e⁻ \rightarrow 4OH⁻(aq)


(a)	Combine these two equations to write a redox reaction for the process.	(1 mark)

(b)	Use your knowledge of chemical equilibrium to predict whether iron will corrode more
	quickly in water that is slightly acidic or water that is slightly basic. Justify your
	reasoning.

	(3 marks)
_	
_	
_	
_	

Question 12 (15 marks)

Below is a representation of an electrochemical cell.



(a) Give the half-equation for the reactions occurring at the anode and cathode and write an overall redox equation for the reaction occurring in the cell. (3 marks)

Anode half-equation:
Cathode half-equation:
Overall equation:

- (b) On the diagram:
 - (i) draw an arrow that shows the direction of electron flow through the external circuit. (1 mark)
 - (ii) label the electrode which is the ANODE. (1 mark)
 - (iii) label the electrode which is POSITIVE (1 mark)
 - (iv) draw an arrow on the salt bridge to show the direction of the migration of POSITIVE IONS. (1 mark)

(c) Standard electrode potentials are measured at standard conditions as shown in the diagram in (a). According to theory, "increasing reactant concentration will increase the cell potential". Based on this statement, predict the effect on the value of the cell potential when each of the following changes were made to the cell. Explain your reasoning. (Equations are not required in your explanations).

(8 marks)


Change	Effect on cell potential (Increase, decrease or no change)	Reasoning
The pressure of hydrogen gas is increased to 200 kPa		
A larger copper electrode is used		
A few drops of silver nitrate solution are added to the copper(II) chloride solution		
The platinum electrode is replaced by a (inert) graphite electrode.		

END OF PART 2

Answer each of the following questions in the space provided.

Question 13

The following diagram represents some of the features of a alkali fuel cell, such as those used on the Apollo missions to the moon. The electrode is a hot solution of KOH soaking in an asbestos cloth. The electrodes are a porous nickel-based material. The electrodes are coated in platinum.



(1 mark)
(1 mark)
(1 mark)


	uming that an alkaline electrolyte is used in the hydrogen/oxygen fuel cell, write the ctions and the overall balanced chemical reaction for the fuel cell. (3 marks			
	Anode half-equation:			
	Cathode half-equation:			
	Overall equation:			
Questic	on 14			
reaction	Balance the following unbalanced redox reaction occurring in acidic medium using the half-reaction method. Write an oxidation half-equation, a reduction half-equation and an overall balanced equation. State symbols are not required.			
	$Sb_2S_3 + NO_3^- \rightarrow Sb_2O_5 + HSO_4^- + NO$ (6 marks)			
	Oxidation half-equation:			
	Reduction half-equation:			
	Overall equation:			

Question 15

State and explain two (2) advantages and two (2) disadvantages of fuel cells or conventional galvanic cells.	ver (4 marks)
Advantage 1:	
Advantage 2:	
Disadvantage 1:	
Disadvantage 2:	

Question 16

Refer to the diagram below to answer the questions. Note that 'chlorine water' is $Cl_2(aq)$. **[6 Marks]**





(b) Write down the balanced ionic equations for all the reactions. (2 marks including states)

(c) Write observation for all three test tubes (3 marks)

Blank Page for Student Use

END OF TEST