Copyright for test papers and marking guides remains with West Australian Test Papers.

Test papers may only be reproduced within the purchasing school according to the advertised Conditions of Sale.

Test papers should be withdrawn after use and stored securely in the school until Friday, 30th June 2017.



## **SEMESTER ONE**

## MATHEMATICS METHODS UNITS 1

2017

**SOLUTIONS** 

[6]

## **Calculator-free Solutions**

(b) (i) 
$$A \cap B \cap C = \{3\}$$

(ii) 
$$(A \cup B)' = \{8, 9, 10, 14, 15, 16, 18, 19\}$$
  
(iii)  $n(B \cup C) = 8$ 

2. (a) (i) 
$$x^{2} + y^{2} = 13$$
  $\checkmark \checkmark$ 

$$y = -\frac{6}{x}$$
(ii)  $y = -x - 1$   $\checkmark \checkmark$ 
(b)  $x = -3, -2, 2, 3$   $\checkmark \checkmark$ 
(c) (i)  $(-\frac{3}{2}, 3)$   $\checkmark \checkmark$ 
(iii)  $m = -2$ 

(iii) 
$$y = -x - 1$$
  
(b)  $x = -3, -2, 2, 3$ 

(c) (i) 
$$(-\frac{3}{2}, 3)$$
  $\checkmark \checkmark$  [11]

3. 
$$(x-1)^2 + (y-1)^2 = 25$$
  
Centre  $(1, 1)$   
 $\frac{4+x}{2} = 1$   $\frac{5+y}{2} = 1$   
M  $(-2, -3)$ 

4. (a) 
$$\sin 225^{\circ}$$
  $= -\sin 45^{\circ}$   $= -\frac{\sqrt{2}}{2}$ 

(b) 
$$\tan\left(\frac{3\pi}{12} + \frac{4\pi}{12}\right) = \tan\left(\frac{\pi}{4} + \frac{\pi}{3}\right)$$

$$= \frac{\tan\frac{\pi}{4} + \tan\frac{\pi}{3}}{1 - \left(\tan\frac{\pi}{4}\right)\left(\tan\frac{\pi}{3}\right)}$$

$$= \frac{1 + \sqrt{3}}{1 - \sqrt{3}}$$

$$= -2 - \sqrt{3}$$
5. (a) 
$$A = \frac{1}{2}(x)(x+3)(\sin 60)$$

$$= \frac{x}{2}(x+3)\left(\frac{\sqrt{3}}{2}\right)$$

$$= \frac{x}{4}(\sqrt{3}x + 3\sqrt{3}) \text{ cm}^2$$
(b)  $x^2 + (x+3)^2 - 2(x)(x+3)(\cos 60) = 49$ 

$$x^2 + 3x - 40 = 0$$

$$x^{2} + 3x - 40 = 0$$


$$(x + 8)(x - 5) = 0$$

$$(x = -8 \text{ discard}) \therefore x = 5$$
(c) 
$$A = \frac{5\sqrt{3}}{4} (5 + 3)$$


(c) 
$$A = \frac{A}{4} (5 + 3)$$
  $\checkmark$  [7]

- 6. (a) 60%
  - (b) 40%
  - (c) 70%

(d)



Or



**√** [5]

- - (b)  $R_{k(x)} = \{y : -6 \le y \le 2, y \in \mathbb{R}\}$
  - (c) For every x value there is only one y value Vertical line through graph
- 8. (a)  $\theta = 210^{\circ} \text{ or } 330^{\circ}$ 
  - (b)  $\cos^2 \theta = \frac{3}{4}$  $\cos \theta = \pm \frac{\sqrt{3}}{2}$

 $\theta = \frac{\pi}{6}, \frac{11\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}$ 

. .

✓

[5]

[8]


## **Calculator-assumed Solutions**

9. (a) (i)  $V = \frac{k}{P}$   $k = 2.84 \times 11.5$  k = 32.66(ii) The volume doubles.

(b) s = 0.0325c is a linear relationship in the form y = kx where k = 0.0325 and is therefore directly proportional.

**√** [5]

10. (a)



(b) (i) 0.3 (ii) 0.24

(iii) 0.46

(iv) 0.348 or  $\frac{16}{46}$ 

**√ √ √** 

**√** 

✓✓

 $\checkmark\checkmark$ 

[9]

[8]

[9]


[10]

- y = 4 + b(a) 11.
  - (0, -c-2)(b)
  - $\{x: x \neq a-3; x \in \mathbb{R}\}$ (c) [5]

 $\sin \theta = -$ 12. (a)  $\cos 2\theta = (\cos \theta)(\cos \theta) - (\sin \theta)(\sin \theta)$ 

 $=\frac{64}{225}-\frac{161}{225}$  $=-\frac{97}{225}$ 

 $AC = \frac{123}{\sin 43^{\circ}}$ (b)  $\angle$  CAD = 12° and  $\angle$ ADC = 31°  $\frac{CD}{\sin 12^{\circ}} = \frac{AC}{\sin 31^{\circ}}$  $\therefore CD = AC \frac{\sin 12^{\circ}}{\cos 12^{\circ}}$ sin 31° sin 12° sin 43° × sin 31° = 73 m



13. (a)


MS 0.175 0.075 0.05 0.7

- (b) 0.075
- (c) 133 0.075
- = 0.60.125 (d)
- $P(MS \cap L) = P(MS) \times P(L)$  to be independent. (e)
  - $0.075 \neq 0.25 \times 0.125$

Therefore not independent or  $P(M \mid L) \neq P(M)$  : Not independent  $\checkmark$ 

14. (i) 3 (a)

- 0 (ii) (iii)
- Domain of p(x) is the same as m(x). For both:  $x \ge 0$ ,  $x \in \mathbb{R}$ (b)
- $q(x) = -\sqrt{x+2} + 1$ (c)
- (d)



15. (a) 
$$-x^2 + 3x + 4 = 2x + q$$
 $-x^2 + x + 4 - q = 0$ 
 $b^2 - 4ac = 0$  for one solution
 $1.17 - 4q = 0$ 
 $color q = 17$ 
 $color q = 17$ 
(b)  $17 - 4q < 0$ 
 $color q = 17$ 
 $color q = 17$ 
(color q =  $17$ 
 $color q = 17$ 
 $color q = 17$ 

(i) (ii) (c)


1 m 5.5m

1.77 m or 28.23 m away (iii)

[9]

19. (a) 
$$y = 2 \sin \left(x - \frac{\pi}{3}\right) - 1$$





(c) 
$$x = -\frac{2\pi}{3} \text{ or } \frac{\pi}{3}$$

20. (a)

	H	H	
G	0.4	0.3	0.7
G'	0.2	0.1	0.3
	0.6	0.4	1

(b) (i) (ii) 0.3

0.6 (iii)

0.9 2 (iv)

Not mutually exclusive:  $P(G \cap H) = 0.4$  Should be 0 (c)

[10]

21. (a) 
$$C^{\left(\frac{3}{2}, \frac{5}{2}\right)}$$
  $r = 3$ 

(i)  $y = -(x + a)(x - b)^{2}$ (ii) y = x(x + a)(x - b)(iii)  $y = (x - a)^{3} + b$ (b)

[5]